

**SCHAUM'S®**  
outlines

# **Mathematical Handbook of Formulas and Tables**

---

**Fifth Edition**

**Murray R. Spiegel, PhD**

*Former Professor and Chairman  
Mathematics Department  
Rensselaer Polytechnic Institute  
Hartford Graduate Center*

**Seymour Lipschutz, PhD**

*Mathematics Department  
Temple University*

**John Liu, PhD**

*Mathematics Department  
University of Maryland*

---

**Schaum's Outline Series**



New York Chicago San Francisco  
Athens London Madrid Mexico City  
Milan New Delhi Singapore Sydney Toronto

Copyright © 2018 by McGraw-Hill Education. All rights reserved. Except as permitted under the United States Copyright Act of 1976, no part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written permission of the publisher.

ISBN: 978-1-26-001054-1  
MHID: 1-26-001054-6

The material in this eBook also appears in the print version of this title: ISBN: 978-1-26-001053-4,  
MHID: 1-26-001053-8.

eBook conversion by codeMantra  
Version 1.0

All trademarks are trademarks of their respective owners. Rather than put a trademark symbol after every occurrence of a trademarked name, we use names in an editorial fashion only, and to the benefit of the trademark owner, with no intention of infringement of the trademark. Where such designations appear in this book, they have been printed with initial caps.

McGraw-Hill Education eBooks are available at special quantity discounts to use as premiums and sales promotions or for use in corporate training programs. To contact a representative, please visit the Contact Us page at [www.mhprofessional.com](http://www.mhprofessional.com).

**SEYMORE LIPSCHUTZ** is on the faculty of Temple University and formally taught at the Polytechnic Institute of Brooklyn. He received his PhD in 1960 at Courant Institute of Mathematical Sciences of New York University. He is one of Schaum's most prolific authors. In particular, he has written, among others, *Linear Algebra*, *Probability*, *Discrete Mathematics*, *Set Theory*, *Finite Mathematics*, and *General Topology*.

**JOHN LIU** is presently a professor of mathematics at University of Maryland, and he formerly taught at Temple University. He received his PhD from the University of California, and he has held visiting positions at New York University, Princeton University, and Berkeley. He has published many papers in applied mathematics, including the areas of partial differential equations and numerical analysis.

The late **MURRAY R. SPIEGEL** received the MS degree in physics and the PhD degree in mathematics from Cornell University. He had positions at Harvard University, Columbia University, Oak Ridge, and Rensselaer Polytechnic Institute, and served as a mathematical consultant at several large companies. His last position was Professor and Chairman of Mathematics at the Rensselaer Polytechnic Institute, Hartford Graduate Center. He was interested in most branches of mathematics, especially those that involve applications to physics and engineering problems. He was the author of numerous journal articles and 14 books on various topics in mathematics.

#### TERMS OF USE

This is a copyrighted work and McGraw-Hill Education and its licensors reserve all rights in and to the work. Use of this work is subject to these terms. Except as permitted under the Copyright Act of 1976 and the right to store and retrieve one copy of the work, you may not decompile, disassemble, reverse engineer, reproduce, modify, create derivative works based upon, transmit, distribute, disseminate, sell, publish or sublicense the work or any part of it without McGraw-Hill Education's prior consent. You may use the work for your own noncommercial and personal use; any other use of the work is strictly prohibited. Your right to use the work may be terminated if you fail to comply with these terms.

THE WORK IS PROVIDED "AS IS." McGRAW-HILL EDUCATION AND ITS LICENSORS MAKE NO GUARANTEES OR WARRANTIES AS TO THE ACCURACY, ADEQUACY OR COMPLETENESS OF OR RESULTS TO BE OBTAINED FROM USING THE WORK, INCLUDING ANY INFORMATION THAT CAN BE ACCESSED THROUGH THE WORK VIA HYPERLINK OR OTHERWISE, AND EXPRESSLY DISCLAIM ANY WARRANTY, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. McGraw-Hill Education and its licensors do not warrant or guarantee that the functions contained in the work will meet your requirements or that its operation will be uninterrupted or error free. Neither McGraw-Hill Education nor its licensors shall be liable to you or anyone else for any inaccuracy, error or omission, regardless of cause, in the work or for any damages resulting therefrom. McGraw-Hill Education has no responsibility for the content of any information accessed through the work. Under no circumstances shall McGraw-Hill Education and/or its licensors be liable for any indirect, incidental, special, punitive, consequential or similar damages that result from the use of or inability to use the work, even if any of them has been advised of the possibility of such damages. This limitation of liability shall apply to any claim or cause whatsoever whether such claim or cause arises in contract, tort or otherwise.

# Preface

This handbook supplies a collection of mathematical formulas and tables which will be valuable to students and research workers in the fields of mathematics, physics, engineering, and other sciences. Care has been taken to include only those formulas and tables which are most likely to be needed in practice, rather than highly specialized results which are rarely used. It is a “user-friendly” handbook with material mostly rooted in university mathematics and scientific courses. In fact, the first edition can already be found in many libraries and offices, and it most likely has moved with the owners from office to office since their college times. Thus, this handbook has survived the test of time (while most other college texts have been thrown away).

This new edition maintains the same spirit as previous editions, with the following changes. First of all, we have deleted some out-of-date tables which can now be easily obtained from a simple calculator, and we have deleted some rarely used formulas. The main change is that sections on Probability and Random Variables have been expanded with new material. These sections appear in both the physical and social sciences, including education. There are also two new sections: Section XIII on Turing Machines and Section XIV on Mathematical Finance.

Topics covered range from elementary to advanced. Elementary topics include those from algebra, geometry, trigonometry, analytic geometry, probability and statistics, and calculus. Advanced topics include those from differential equations, numerical analysis, and vector analysis, such as Fourier series, gamma and beta functions, Bessel and Legendre functions, Fourier and Laplace transforms, and elliptic and other special functions of importance. This wide coverage of topics has been adopted to provide, within a single volume, most of the important mathematical results needed by student and research workers, regardless of their particular field of interest or level of attainment.

The book is divided into two main parts. Part A presents mathematical formulas together with other material, such as definitions, theorems, graphs, diagrams, etc., essential for proper understanding and application of the formulas. Part B presents the numerical tables. These tables include basic statistical distributions (normal, Student's *t*, chi-square, etc.), advanced functions (Bessel, Legendre, elliptic, etc.), and financial functions (compound and present value of an amount, and annuity).

McGraw-Hill Education wishes to thank the various authors and publishers—for example, the Literary Executor of the late Sir Ronald A. Fisher, F.R.S., Dr. Frank Yates, F.R.S., and Oliver and Boyd Ltd., Edinburgh, for Table III of their book *Statistical Tables for Biological, Agricultural and Medical Research*—who gave their permission to adapt data from their books for use in several tables in this handbook. Appropriate references to such sources are given below the corresponding tables.

Finally, I wish to thank the staff of McGraw-Hill Education Schaum's Outline Series, especially Diane Grayson, for their unfailing cooperation.

SEYMOUR LIPSCHUTZ  
Temple University

# **Contents**

---

<b>Part A</b>	<b>FORMULAS</b>	<b>1</b>
<b>Section I</b>	<b>Elementary Constants, Products, Formulas</b>	<b>3</b>
1.	Greek Alphabet and Special Constants	3
2.	Special Products and Factors	5
3.	The Binomial Formula and Binomial Coefficients	7
4.	Complex Numbers	10
5.	Solutions of Algebraic Equations	13
6.	Conversion Factors	15
<b>Section II</b>	<b>Geometry</b>	<b>16</b>
7.	Geometric Formulas	16
8.	Formulas from Plane Analytic Geometry	22
9.	Special Plane Curves	28
10.	Formulas from Solid Analytic Geometry	34
11.	Special Moments of Inertia	41
<b>Section III</b>	<b>Elementary Transcendental Functions</b>	<b>43</b>
12.	Trigonometric Functions	43
13.	Exponential and Logarithmic Functions	53
14.	Hyperbolic Functions	56
<b>Section IV</b>	<b>Calculus</b>	<b>62</b>
15.	Derivatives	62
16.	Indefinite Integrals	67
17.	Tables of Special Indefinite Integrals	71
18.	Definite Integrals	108
<b>Section V</b>	<b>Differential Equations and Vector Analysis</b>	<b>116</b>
19.	Basic Differential Equations and Solutions	116
20.	Formulas from Vector Analysis	119
<b>Section VI</b>	<b>Series</b>	<b>134</b>
21.	Series of Constants	134
22.	Taylor Series	138
23.	Bernoulli and Euler Numbers	142
24.	Fourier Series	144

<b>Section VII</b>	<b>Special Functions and Polynomials</b>	<b>149</b>
25.	The Gamma Function	149
26.	The Beta Function	152
27.	Bessel Functions	153
28.	Legendre and Associated Legendre Functions	164
29.	Hermite Polynomials	169
30.	Laguerre and Associated Laguerre Polynomials	171
31.	Chebyshev Polynomials	175
32.	Hypergeometric Functions	178
<b>Section VIII</b>	<b>Laplace and Fourier Transforms</b>	<b>180</b>
33.	Laplace Transforms	180
34.	Fourier Transforms	193
<b>Section IX</b>	<b>Elliptic and Miscellaneous Special Functions</b>	<b>198</b>
35.	Elliptic Functions	198
36.	Miscellaneous and Riemann Zeta Functions	203
<b>Section X</b>	<b>Inequalities and Infinite Products</b>	<b>205</b>
37.	Inequalities	205
38.	Infinite Products	207
<b>Section XI</b>	<b>Probability and Statistics</b>	<b>208</b>
39.	Descriptive Statistics	208
40.	Probability	217
41.	Random Variables	223
<b>Section XII</b>	<b>Numerical Methods</b>	<b>231</b>
42.	Interpolation	231
43.	Quadrature	235
44.	Solution of Nonlinear Equations	237
45.	Numerical Methods for Ordinary Differential Equations	239
46.	Numerical Methods for Partial Differential Equations	241
47.	Iteration Methods for Linear Systems	244
<b>Section XIII</b>	<b>Turing Machines</b>	<b>246</b>
48.	Basic Definitions, Expressions	246
49.	Pictures	247
50.	Quintuple, Turing Machine	248
51.	Computing with a Turing Machine	250
52.	Examples	252
<b>Section XIV</b>	<b>Mathematical Finance</b>	<b>254</b>
53.	Basic Probability	254
54.	Interest Rates	256
55.	Arbitrage Theorem and Options	257

<b>56.</b> Arbitrage Theorem <b>57.</b> Black-Scholes Formula <b>58.</b> The Delta Hedging Arbitrage Strategy	258 259 260
<b>Part B      TABLES</b>	<b>263</b>
<b>Section I    Logarithmic, Trigonometric, Exponential Functions</b>	<b>265</b>
<b>1.</b> Four Place Common Logarithms $\log_{10} N$ or $\log N$ <b>2.</b> Sin $x$ ( $x$ in Degrees and Minutes) <b>3.</b> Cos $x$ ( $x$ in Degrees and Minutes) <b>4.</b> Tan $x$ ( $x$ in Degrees and Minutes) <b>5.</b> Conversion of Radians to Degrees, Minutes, and Seconds or Fractions of Degrees <b>6.</b> Conversion of Degrees, Minutes, and Seconds to Radians <b>7.</b> Natural or Napierian Logarithms $\log_e x$ or $\ln x$ <b>8.</b> Exponential Functions $e^x$ <b>9.</b> Exponential Functions $e^{-x}$ <b>10.</b> Exponential, Sine, and Cosine Integrals	265 267 268 269 270 271 272 274 275 276
<b>Section II   Factorial and Gamma Function, Binomial Coefficients</b>	<b>277</b>
<b>11.</b> Factorial $n$ <b>12.</b> Gamma Function <b>13.</b> Binomial Coefficients	277 278 279
<b>Section III   Bessel Functions</b>	<b>281</b>
<b>14.</b> Bessel Functions $J_0(x)$ <b>15.</b> Bessel Functions $J_1(x)$ <b>16.</b> Bessel Functions $Y_0(x)$ <b>17.</b> Bessel Functions $Y_1(x)$ <b>18.</b> Bessel Functions $I_0(x)$ <b>19.</b> Bessel Functions $I_1(x)$ <b>20.</b> Bessel Functions $K_0(x)$ <b>21.</b> Bessel Functions $K_1(x)$ <b>22.</b> Bessel Functions $\text{Ber}(x)$ <b>23.</b> Bessel Functions $\text{Bei}(x)$ <b>24.</b> Bessel Functions $\text{Ker}(x)$ <b>25.</b> Bessel Functions $\text{Kei}(x)$ <b>26.</b> Values for Approximate Zeros of Bessel Functions	281 281 282 282 283 283 284 284 285 285 286 286 287
<b>Section IV   Legendre Polynomials</b>	<b>288</b>
<b>27.</b> Legendre Polynomials $P_n(x)$ <b>28.</b> Legendre Polynomials $P_n(\cos \theta)$	288 289
<b>Section V   Elliptic Integrals</b>	<b>290</b>
<b>29.</b> Complete Elliptic Integrals of First and Second Kinds <b>30.</b> Incomplete Elliptic Integral of the First Kind <b>31.</b> Incomplete Elliptic Integral of the Second Kind	290 291 291

<b>Section VI</b>	<b>Financial Tables</b>	<b>292</b>
32.	Compound Amount: $(1 + r)^n$	292
33.	Present Value of an Amount: $(1 + r)^{-n}$	293
34.	Amount of an Annuity: $\frac{(1+r)^n - 1}{r}$	294
35.	Present Value of an Annuity: $\frac{1 - (1+r)^{-n}}{r}$	295
<b>Section VII</b>	<b>Probability and Statistics</b>	<b>296</b>
36.	Areas Under the Standard Normal Curve from $-\infty$ to $x$	296
37.	Ordinates of the Standard Normal Curve	297
38.	Percentile Values ( $t_p$ ) for Student's $t$ Distribution	298
39.	Percentile Values ( $\chi_p^2$ ) for $\chi^2$ (Chi-Square) Distribution	299
40.	95th Percentile Values for the $F$ Distribution	300
41.	99th Percentile Values for the $F$ Distribution	301
42.	Random Numbers	302
	<b>Index of Special Symbols and Notations</b>	<b>303</b>
	<b>Index</b>	<b>305</b>

**PART A**

# **FORMULAS**

## Section I: Elementary Constants, Products, Formulas

# 1

## GREEK ALPHABET and SPECIAL CONSTANTS

### Greek Alphabet

Greek name	Greek letter	
	Lower case	Capital
Alpha	$\alpha$	A
Beta	$\beta$	B
Gamma	$\gamma$	$\Gamma$
Delta	$\delta$	$\Delta$
Epsilon	$\varepsilon$	E
Zeta	$\zeta$	Z
Eta	$\eta$	H
Theta	$\theta$	$\Theta$
Iota	$\iota$	I
Kappa	$\kappa$	K
Lambda	$\lambda$	$\Lambda$
Mu	$\mu$	M

Greek name	Greek letter	
	Lower case	Capital
Nu	$\nu$	N
Xi	$\xi$	$\Xi$
Omicron	$\o$	O
Pi	$\pi$	$\Pi$
Rho	$\rho$	R
Sigma	$\sigma$	$\Sigma$
Tau	$\tau$	T
Upsilon	$\upsilon$	$\Upsilon$
Phi	$\phi$	$\Phi$
Chi	$\chi$	X
Psi	$\psi$	$\Psi$
Omega	$\omega$	$\Omega$

### Special Constants

1.1.  $\pi = 3.14159\ 26535\ 89793 \dots$

1.2.  $e = 2.71828\ 18284\ 59045 \dots = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$

= natural base of logarithms

1.3.  $\gamma = 0.57721\ 56649\ 01532\ 86060\ 6512 \dots = Euler's\ constant$

$= \lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} - \ln n\right)$

1.4.  $e^\gamma = 1.78107\ 24179\ 90197\ 9852 \dots$  [see 1.3]

**1.5.**  $\sqrt{e} = 1.64872\ 12707\ 00128\ 1468 \dots$

**1.6.**  $\sqrt{\pi} = \Gamma(\frac{1}{2}) = 1.77245\ 38509\ 05516\ 02729\ 8167 \dots$   
where  $\Gamma$  is the *gamma function* [see 25.1].

**1.7.**  $\Gamma(\frac{1}{3}) = 2.67893\ 85347\ 07748 \dots$

**1.8.**  $\Gamma(\frac{1}{4}) = 3.62560\ 99082\ 21908 \dots$

**1.9.** 1 radian =  $180^\circ/\pi = 57.29577\ 95130\ 8232 \dots^\circ$

**1.10.**  $1^\circ = \pi/180$  radians =  $0.01745\ 32925\ 19943\ 29576\ 92 \dots$  radians

# 2

# SPECIAL PRODUCTS and FACTORS

$$2.1. \quad (x+y)^2 = x^2 + 2xy + y^2$$

$$2.2. \quad (x-y)^2 = x^2 - 2xy + y^2$$

$$2.3. \quad (x+y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$2.4. \quad (x-y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

$$2.5. \quad (x+y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

$$2.6. \quad (x-y)^4 = x^4 - 4x^3y + 6x^2y^2 - 4xy^3 + y^4$$

$$2.7. \quad (x+y)^5 = x^5 + 5x^4y + 10x^3y^2 + 10x^2y^3 + 5xy^4 + y^5$$

$$2.8. \quad (x-y)^5 = x^5 - 5x^4y + 10x^3y^2 - 10x^2y^3 + 5xy^4 - y^5$$

$$2.9. \quad (x+y)^6 = x^6 + 6x^5y + 15x^4y^2 + 20x^3y^3 + 15x^2y^4 + 6xy^5 + y^6$$

$$2.10. \quad (x-y)^6 = x^6 - 6x^5y + 15x^4y^2 - 20x^3y^3 + 15x^2y^4 - 6xy^5 + y^6$$

The results 2.1 to 2.10 above are special cases of the *binomial formula* [see 3.3].

$$2.11. \quad x^2 - y^2 = (x-y)(x+y)$$

$$2.12. \quad x^3 - y^3 = (x-y)(x^2 + xy + y^2)$$

$$2.13. \quad x^3 + y^3 = (x+y)(x^2 - xy + y^2)$$

$$2.14. \quad x^4 - y^4 = (x-y)(x+y)(x^2 + y^2)$$

$$2.15. \quad x^5 - y^5 = (x-y)(x^4 + x^3y + x^2y^2 + xy^3 + y^4)$$

$$2.16. \quad x^5 + y^5 = (x+y)(x^4 - x^3y + x^2y^2 - xy^3 + y^4)$$

$$2.17. \quad x^6 - y^6 = (x-y)(x+y)(x^2 + xy + y^2)(x^2 - xy + y^2)$$

$$\mathbf{2.18.} \quad x^4 + x^2y^2 + y^4 = (x^2 + xy + y^2)(x^2 - xy + y^2)$$

$$\mathbf{2.19.} \quad x^4 + 4y^4 = (x^2 + 2xy + 2y^2)(x^2 - 2xy + 2y^2)$$

Some generalizations of the above are given by the following results where  $n$  is a positive integer.

$$\begin{aligned}\mathbf{2.20.} \quad x^{2n+1} - y^{2n+1} &= (x - y)(x^{2n} + x^{2n-1}y + x^{2n-2}y^2 + \dots + y^{2n}) \\ &= (x - y) \left( x^2 - 2xy \cos \frac{2\pi}{2n+1} + y^2 \right) \left( x^2 - 2xy \cos \frac{4\pi}{2n+1} + y^2 \right) \\ &\quad \dots \left( x^2 - 2xy \cos \frac{2n\pi}{2n+1} + y^2 \right)\end{aligned}$$

$$\begin{aligned}\mathbf{2.21.} \quad x^{2n+1} + y^{2n+1} &= (x + y)(x^{2n} - x^{2n-1}y + x^{2n-2}y^2 - \dots + y^{2n}) \\ &= (x + y) \left( x^2 + 2xy \cos \frac{2\pi}{2n+1} + y^2 \right) \left( x^2 + 2xy \cos \frac{4\pi}{2n+1} + y^2 \right) \\ &\quad \dots \left( x^2 + 2xy \cos \frac{2n\pi}{2n+1} + y^2 \right)\end{aligned}$$

$$\begin{aligned}\mathbf{2.22.} \quad x^{2n} - y^{2n} &= (x - y)(x + y)(x^{n-1} + x^{n-2}y + x^{n-3}y^2 + \dots)(x^{n-1} - x^{n-2}y + x^{n-3}y^2 - \dots) \\ &= (x - y)(x + y) \left( x^2 - 2xy \cos \frac{\pi}{n} + y^2 \right) \left( x^2 - 2xy \cos \frac{2\pi}{n} + y^2 \right) \\ &\quad \dots \left( x^2 - 2xy \cos \frac{(n-1)\pi}{n} + y^2 \right)\end{aligned}$$

$$\begin{aligned}\mathbf{2.23.} \quad x^{2n} + y^{2n} &= \left( x^2 + 2xy \cos \frac{\pi}{2n} + y^2 \right) \left( x^2 + 2xy \cos \frac{3\pi}{2n} + y^2 \right) \\ &\quad \dots \left( x^2 + 2xy \cos \frac{(2n-1)\pi}{2n} + y^2 \right)\end{aligned}$$

# 3

# THE BINOMIAL FORMULA and BINOMIAL COEFFICIENTS

## Factorial $n$

For  $n = 1, 2, 3, \dots$ , factorial  $n$  or  $n$  factorial is denoted and defined by

$$3.1. \quad n! = n(n-1) \cdots 3 \cdot 2 \cdot 1$$

Zero factorial is defined by

$$3.2. \quad 0! = 1$$

Alternately,  $n$  factorial can be defined recursively by

$$0! = 1 \quad \text{and} \quad n! = n \cdot (n-1)!$$

**EXAMPLE:**  $4! = 4 \cdot 3 \cdot 2 \cdot 1 = 24,$   
 $5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 5 \cdot 4! = 5(24) = 120,$   
 $6! = 6 \cdot 5! = 6(120) = 720$

## Binomial Formula for Positive Integral $n$

For  $n = 1, 2, 3, \dots$ ,

$$3.3. \quad (x+y)^n = x^n + nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 + \frac{n(n-1)(n-2)}{3!}x^{n-3}y^3 + \cdots + y^n$$

This is called the *binomial formula*. It can be extended to other values of  $n$ , and also to an infinite series [see 22.4].

**EXAMPLE:**

(a)  $(a-2b)^4 = a^4 + 4a^3(-2b) + 6a^2(-2b)^2 + 4a(-2b)^3 + (-2b)^4 = a^4 - 8a^3b + 24a^2b^2 - 32ab^3 + 16b^4$

Here  $x = a$  and  $y = -2b$ .

(b) See Fig. 3-1a.

## Binomial Coefficients

Formula 3.3 can be rewritten in the form

$$3.4. \quad (x+y)^n = x^n + \binom{n}{1}x^{n-1}y + \binom{n}{2}x^{n-2}y^2 + \binom{n}{3}x^{n-3}y^3 + \cdots + \binom{n}{n}y^n$$

where the coefficients, called *binomial coefficients*, are given by

$$3.5. \quad \binom{n}{k} = \frac{n(n-1)(n-2)\cdots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!} = \binom{n}{n-k}$$

$$\text{EXAMPLE: } \binom{9}{4} = \frac{9 \cdot 8 \cdot 7 \cdot 6}{1 \cdot 2 \cdot 3 \cdot 4} = 126, \quad \binom{12}{5} = \frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8}{1 \cdot 2 \cdot 3 \cdot 4 \cdot 5} = 792, \quad \binom{10}{7} = \binom{10}{3} = \frac{10 \cdot 9 \cdot 8}{1 \cdot 2 \cdot 3} = 120$$

Note that  $\binom{n}{r}$  has exactly  $r$  factors in both the numerator and the denominator.

The binomial coefficients may be arranged in a triangular array of numbers, called Pascal's triangle, as shown in Fig. 3-1b. The triangle has the following two properties:

- (1) The first and last number in each row is 1.
- (2) Every other number in the array can be obtained by adding the two numbers appearing directly above it. For example

$$10 = 4 + 6, \quad 15 = 5 + 10, \quad 20 = 10 + 10$$

Property (2) may be stated as follows:

$$3.6. \quad \binom{n}{k} + \binom{n}{k+1} = \binom{n+1}{k+1}$$




Fig. 3-1

### Properties of Binomial Coefficients

The following lists additional properties of the binomial coefficients:

$$3.7. \quad \binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \cdots + \binom{n}{n} = 2^n$$

$$3.8. \quad \binom{n}{0} - \binom{n}{1} + \binom{n}{2} - \cdots - (-1)^n \binom{n}{n} = 0$$

$$3.9. \quad \binom{n}{n} + \binom{n+1}{n} + \binom{n+2}{n} + \cdots + \binom{n+m}{n} = \binom{n+m+1}{n+1}$$

**3.10.**  $\binom{n}{0} + \binom{n}{2} + \binom{n}{4} + \cdots = 2^{n-1}$

**3.11.**  $\binom{n}{1} + \binom{n}{3} + \binom{n}{5} + \cdots = 2^{n-1}$

**3.12.**  $\binom{n}{0}^2 + \binom{n}{1}^2 + \binom{n}{2}^2 + \cdots + \binom{n}{n}^2 = \binom{2n}{n}$

**3.13.**  $\binom{m}{0} \binom{n}{p} + \binom{m}{1} \binom{n}{p-1} + \cdots + \binom{m}{p} \binom{n}{0} = \binom{m+n}{p}$

**3.14.**  $(1) \binom{n}{1} + (2) \binom{n}{2} + (3) \binom{n}{3} + \cdots + (n) \binom{n}{n} = n2^{n-1}$

**3.15.**  $(1) \binom{n}{1} - (2) \binom{n}{2} + (3) \binom{n}{3} - \cdots - (-1)^{n+1} (n) \binom{n}{n} = 0$

### Multinomial Formula

---

Let  $n_1, n_2, \dots, n_r$  be nonnegative integers such that  $n_1 + n_2 + \cdots + n_r = n$ . Then the following expression, called a *multinomial coefficient*, is defined as follows:

**3.16.**  $\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!}$

**EXAMPLE:**  $\binom{7}{2, 3, 2} = \frac{7!}{2!3!2!} = 210, \quad \binom{8}{4, 2, 2, 0} = \frac{8!}{4!2!2!0!} = 420$

The name multinomial coefficient comes from the following formula:

**3.17.**  $(x_1 + x_2 + \cdots + x_p)^n = \sum \binom{n}{n_1, n_2, \dots, n_r} x_1^{n_1} x_2^{n_2} \cdots x_r^{n_r}$

where the sum, denoted by  $\Sigma$ , is taken over all possible multinomial coefficients.

# 4 COMPLEX NUMBERS

## Definitions Involving Complex Numbers

A complex number  $z$  is generally written in the form

$$z = a + bi$$

where  $a$  and  $b$  are real numbers and  $i$ , called the *imaginary unit*, has the property that  $i^2 = -1$ . The real numbers  $a$  and  $b$  are called the *real* and *imaginary parts* of  $z = a + bi$ , respectively.

The *complex conjugate* of  $z$  is denoted by  $\bar{z}$ ; it is defined by

$$\overline{a+bi} = a - bi$$

Thus,  $a + bi$  and  $a - bi$  are conjugates of each other.

## Equality of Complex Numbers

4.1.  $a + bi = c + di$  if and only if  $a = c$  and  $b = d$

## Arithmetic of Complex Numbers

Formulas for the addition, subtraction, multiplication, and division of complex numbers follow:

4.2.  $(a + bi) + (c + di) = (a + c) + (b + d)i$

4.3.  $(a + bi) - (c + di) = (a - c) + (b - d)i$

4.4.  $(a + bi)(c + di) = (ac - bd) + (ad + bc)i$

4.5.  $\frac{a+bi}{c+di} = \frac{a+bi}{c+di} \cdot \frac{c-di}{c-di} = \frac{ac+bd}{c^2+d^2} + \left( \frac{bc-ad}{c^2+d^2} \right) i$

Note that the above operations are obtained by using the ordinary rules of algebra and replacing  $i^2$  by  $-1$  wherever it occurs.

**EXAMPLE:** Suppose  $z = 2 + 3i$  and  $w = 5 - 2i$ . Then

$$\begin{aligned} z + w &= (2 + 3i) + (5 - 2i) = 2 + 5 + 3i - 2i = 7 + i \\ zw &= (2 + 3i)(5 - 2i) = 10 + 15i - 4i - 6i^2 = 16 + 11i \\ \bar{z} &= \overline{2+3i} = 2 - 3i \text{ and } \bar{w} = \overline{5-2i} = 5 + 2i \\ \frac{w}{z} &= \frac{5-2i}{2+3i} = \frac{(5-2i)(2-3i)}{(2+3i)(2-3i)} = \frac{4-19i}{13} = \frac{4}{13} - \frac{19}{13}i \end{aligned}$$

## Complex Plane

Real numbers can be represented by the points on a line, called the *real line*, and, similarly, complex numbers can be represented by points in the plane, called the *Argand diagram* or *Gaussian plane* or, simply, the *complex plane*. Specifically, we let the point  $(a, b)$  in the plane represent the complex number  $z = a + bi$ . For example, the point  $P$  in Fig. 4-1 represents the complex number  $z = -3 + 4i$ . The complex number can also be interpreted as a vector from the origin  $O$  to the point  $P$ .

The *absolute value* of a complex number  $z = a + bi$ , written  $|z|$ , is defined as follows:

$$4.6. \quad |z| = \sqrt{a^2 + b^2} = \sqrt{zz}$$

We note  $|z|$  is the distance from the origin  $O$  to the point  $z$  in the complex plane.




Fig. 4-1




Fig. 4-2

## Polar Form of Complex Numbers

The point  $P$  in Fig. 4-2 with coordinates  $(x, y)$  represents the complex number  $z = x + iy$ . The point  $P$  can also be represented by *polar coordinates*  $(r, \theta)$ . Since  $x = r \cos \theta$  and  $y = r \sin \theta$ , we have

$$4.7. \quad z = x + iy = r(\cos \theta + i \sin \theta)$$

called the *polar form* of the complex number. We often call  $r = |z| = \sqrt{x^2 + y^2}$  the *modulus* and  $\theta$  the *amplitude* of  $z = x + iy$ .

## Multiplication and Division of Complex Numbers in Polar Form

$$4.8. \quad [r_1(\cos \theta_1 + i \sin \theta_1)][r_2(\cos \theta_2 + i \sin \theta_2)] = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$$

$$4.9. \quad \frac{r_1(\cos \theta_1 + i \sin \theta_1)}{r_2(\cos \theta_2 + i \sin \theta_2)} = \frac{r_1}{r_2} [\cos(\theta_1 - \theta_2) + i \sin(\theta_1 - \theta_2)]$$

## De Moivre's Theorem

For any real number  $p$ , De Moivre's theorem states that

$$4.10. \quad [r(\cos \theta + i \sin \theta)]^p = r^p (\cos p\theta + i \sin p\theta)$$

### Roots of Complex Numbers

---

Let  $p = 1/n$  where  $n$  is any positive integer. Then 4.10 can be written

$$4.11. \quad [r(\cos\theta + i\sin\theta)]^{\frac{1}{n}} = r^{\frac{1}{n}} \left( \cos \frac{\theta + 2k\pi}{n} + i\sin \frac{\theta + 2k\pi}{n} \right)$$

where  $k$  is any integer. From this formula, all the  $n$ th roots of a complex number can be obtained by putting  $k = 0, 1, 2, \dots, n - 1$ .

# 5

## SOLUTIONS of ALGEBRAIC EQUATIONS

### Quadratic Equation: $ax^2 + bx + c = 0$

#### 5.1. Solutions:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

If  $a, b, c$  are real and if  $D = b^2 - 4ac$  is the *discriminant*, then the roots are

- (i) real and unequal if  $D > 0$
- (ii) real and equal if  $D = 0$
- (iii) complex conjugate if  $D < 0$

5.2. If  $x_1, x_2$  are the roots, then  $x_1 + x_2 = -b/a$  and  $x_1 x_2 = c/a$ .

### Cubic Equation: $x^3 + a_1x^2 + a_2x + a_3 = 0$

Let

$$\begin{aligned} Q &= \frac{3a_2 - a_1^2}{9}, & R &= \frac{9a_1a_2 - 27a_3 - 2a_1^3}{54}, \\ S &= \sqrt[3]{R + \sqrt{Q^3 + R^2}}, & T &= \sqrt[3]{R - \sqrt{Q^3 + R^2}} \end{aligned}$$

where  $ST = -Q$ .

#### 5.3. Solutions:

$$\begin{cases} x_1 = S + T - \frac{1}{3}a_1 \\ x_2 = -\frac{1}{2}(S + T) - \frac{1}{3}a_1 + \frac{1}{2}i\sqrt{3}(S - T) \\ x_3 = -\frac{1}{2}(S + T) - \frac{1}{3}a_1 - \frac{1}{2}i\sqrt{3}(S - T) \end{cases}$$

If  $a_1, a_2, a_3$  are real and if  $D = Q^3 + R^2$  is the *discriminant*, then

- (i) one root is real and two are complex conjugate if  $D > 0$
- (ii) all roots are real and at least two are equal if  $D = 0$
- (iii) all roots are real and unequal if  $D < 0$ .

If  $D < 0$ , computation is simplified by use of trigonometry.

#### 5.4. Solutions:

$$\text{if } D < 0 : \begin{cases} x_1 = 2\sqrt{-Q} \cos\left(\frac{1}{3}\theta\right) - \frac{1}{3}a_1 \\ x_2 = 2\sqrt{-Q} \cos\left(\frac{1}{3}\theta + 120^\circ\right) - \frac{1}{3}a_1 \\ x_3 = 2\sqrt{-Q} \cos\left(\frac{1}{3}\theta + 240^\circ\right) - \frac{1}{3}a_1 \end{cases}$$

where  $\cos\theta = R/\sqrt{-Q^3}$

**5.5.**  $x_1 + x_2 + x_3 = -a_1$ ,  $x_1x_2 + x_2x_3 + x_3x_1 = a_2$ ,  $x_1x_2x_3 = -a_3$   
where  $x_1, x_2, x_3$  are the three roots.

---

**Quartic Equation:**  $x^4 + a_1x^3 + a_2x^2 + a_3x + a_4 = 0$

---

Let  $y_1$  be a real root of the following cubic equation:

$$\mathbf{5.6.} \quad y^3 - a_2y^2 + (a_1a_3 - 4a_4)y + (4a_2a_4 - a_3^2 - a_1^2a_4) = 0$$

The four roots of the quartic equation are the four roots of the following equation:

$$\mathbf{5.7.} \quad z^2 + \frac{1}{2}(a_1 \pm \sqrt{a_1^2 - 4a_2 + 4y_1})z + \frac{1}{2}(y_1 \mp \sqrt{y_1^2 - 4a_4}) = 0$$

Suppose that all roots of 5.6 are real; then computation is simplified by using the particular real root that produces all real coefficients in the quadratic equation 5.7.

$$\mathbf{5.8.} \quad \begin{cases} x_1 + x_2 + x_3 + x_4 = -a_1 \\ x_1x_2 + x_2x_3 + x_3x_4 + x_4x_1 + x_1x_3 + x_2x_4 = a_2 \\ x_1x_2x_3 + x_2x_3x_4 + x_1x_2x_4 + x_1x_3x_4 = -a_3 \\ x_1x_2x_3x_4 = x_4 \end{cases}$$

where  $x_1, x_2, x_3, x_4$  are the four roots.

# 6 CONVERSION FACTORS

<b>Length</b>	1 kilometer (km) = 1000 meters (m)	1 inch (in) = 2.540 cm
	1 meter (m) = 100 centimeters (cm)	1 foot (ft) = 30.48 cm
	1 centimeter (cm) = $10^{-2}$ m	1 mile (mi) = 1.609 km
	1 millimeter (mm) = $10^{-3}$ m	1 millimeter = $10^{-3}$ in
	1 micron ( $\mu$ ) = $10^{-6}$ m	1 centimeter = 0.3937 in
	1 millimicron ( $m\mu$ ) = $10^{-9}$ m	1 meter = 39.37 in
	1 angstrom ( $\text{\AA}$ ) = $10^{-10}$ m	1 kilometer = 0.6214 mi
<b>Area</b>	1 square meter ( $\text{m}^2$ ) = 10.76 ft <sup>2</sup>	1 square mile ( $\text{mi}^2$ ) = 640 acres
	1 square foot ( $\text{ft}^2$ ) = 929 cm <sup>2</sup>	1 acre = 43,560 ft <sup>2</sup>
<b>Volume</b>	1 liter ( $l$ ) = 1000 cm <sup>3</sup> = 1.057 quart (qt) = 61.02 in <sup>3</sup> = 0.03532 ft <sup>3</sup>	
	1 cubic meter ( $\text{m}^3$ ) = 1000 $l$ = 35.32 ft <sup>3</sup>	
	1 cubic foot (ft <sup>3</sup> ) = 7.481 U.S. gal = 0.02832 m <sup>3</sup> = 28.32 $l$	
	1 U.S. gallon (gal) = 231 in <sup>3</sup> = 3.785 $l$ ; 1 British gallon = 1.201 U.S. gallon = 277.4 in <sup>3</sup>	
<b>Mass</b>	1 kilogram (kg) = 2.2046 pounds (lb) = 0.06852 slug; 1 lb = 453.6 gm = 0.03108 slug	
	1 slug = 32.174 lb = 14.59 kg	
<b>Speed</b>	1 km/hr = 0.2778 m/sec = 0.6214 mi/hr = 0.9113 ft/sec	
	1 mi/hr = 1.467 ft/sec = 1.609 km/hr = 0.4470 m/sec	
<b>Density</b>	1 gm/cm <sup>3</sup> = $10^3$ kg/m <sup>3</sup> = 62.43 lb/ft <sup>3</sup> = 1.940 slug/ft <sup>3</sup>	
	1 lb/ft <sup>3</sup> = 0.01602 gm/cm <sup>3</sup> ; 1 slug/ft <sup>3</sup> = 0.5154 gm/cm <sup>3</sup>	
<b>Force</b>	1 newton (nt) = $10^5$ dynes = 0.1020 kgwt = 0.2248 lbwt	
	1 pound weight (lbwt) = 4.448 nt = 0.4536 kgwt = 32.17 poundals	
	1 kilogram weight (kgwt) = 2.205 lbwt = 9.807 nt	
	1 U.S. short ton = 2000 lbwt; 1 long ton = 2240 lbwt; 1 metric ton = 2205 lbwt	
<b>Energy</b>	1 joule = 1 nt m = $10^7$ ergs = 0.7376 ft lbwt = 0.2389 cal = $9.481 \times 10^{-4}$ Btu	
	1 ft lbwt = 1.356 joules = 0.3239 cal = $1.285 \times 10^{-3}$ Btu	
	1 calorie (cal) = 4.186 joules = 3.087 ft lbwt = $3.968 \times 10^{-3}$ Btu	
	1 Btu (British thermal unit) = 778 ft lbwt = 1055 joules = 0.293 watt hr	
	1 kilowatt hour (kw hr) = $3.60 \times 10^6$ joules = 860.0 kcal = 3413 Btu	
	1 electron volt (ev) = $1.602 \times 10^{-19}$ joule	
<b>Power</b>	1 watt = 1 joule/sec = $10^7$ ergs/sec = 0.2389 cal/sec	
	1 horsepower (hp) = 550 ft lbwt/sec = 33,000 ft lbwt/min = 745.7 watts	
	1 kilowatt (kw) = 1.341 hp = 737.6 ft lbwt/sec = 0.9483 Btu/sec	
<b>Pressure</b>	1 nt/m <sup>2</sup> = 10 dynes/cm <sup>2</sup> = $9.869 \times 10^{-6}$ atmosphere = $2.089 \times 10^{-2}$ lbwt/ft <sup>2</sup>	
	1 lbwt/in <sup>2</sup> = 6895 nt/m <sup>2</sup> = 5.171 cm mercury = 27.68 in water	
	1 atm = $1.013 \times 10^5$ nt/m <sup>2</sup> = $1.013 \times 10^6$ dynes/cm <sup>2</sup> = 14.70 lbwt/in <sup>2</sup>	
	= 76 cm mercury = 406.8 in water	

## Section II: Geometry

# 7

## GEOMETRIC FORMULAS

### Rectangle of Length $b$ and Width $a$

7.1. Area =  $ab$

7.2. Perimeter =  $2a + 2b$




Fig. 7-1

### Parallelogram of Altitude $h$ and Base $b$

7.3. Area =  $bh = ab \sin \theta$

7.4. Perimeter =  $2a + 2b$




Fig. 7-2

### Triangle of Altitude $h$ and Base $b$

7.5. Area =  $\frac{1}{2}bh = \frac{1}{2}ab \sin \theta$

$$= \sqrt{s(s-a)(s-b)(s-c)}$$

where  $s = \frac{1}{2}(a+b+c)$  = semiperimeter




Fig. 7-3

7.6. Perimeter =  $a + b + c$

### Trapezoid of Altitude $h$ and Parallel Sides $a$ and $b$

7.7. Area =  $\frac{1}{2}h(a + b)$

7.8. Perimeter =  $a + b + h\left(\frac{1}{\sin \theta} + \frac{1}{\sin \phi}\right)$   
 $= a + b + h(\csc \theta + \csc \phi)$




Fig. 7-4

**Regular Polygon of  $n$  Sides Each of Length  $b$** 

7.9. Area =  $\frac{1}{4}nb^2 \cot \frac{\pi}{n} = \frac{1}{4}nb^2 \frac{\cos(\pi/n)}{\sin(\pi/n)}$

7.10. Perimeter =  $nb$




Fig. 7-5

**Circle of Radius  $r$** 

7.11. Area =  $\pi r^2$




Fig. 7-6

7.12. Perimeter =  $2\pi r$




Fig. 7-7

**Radius of Circle Inscribed in a Triangle of Sides  $a, b, c$** 

7.15.  $r = \frac{\sqrt{s(s-a)(s-b)(s-c)}}{s}$

where  $s = \frac{1}{2}(a + b + c)$  = semiperimeter.




Fig. 7-8

**Radius of Circle Circumscribing a Triangle of Sides  $a, b, c$** 

7.16.  $R = \frac{abc}{4\sqrt{s(s-a)(s-b)(s-c)}}$

where  $s = \frac{1}{2}(a + b + c)$  = semiperimeter.




Fig. 7-9

### Regular Polygon of $n$ Sides Inscribed in Circle of Radius $r$

7.17. Area =  $\frac{1}{2}nr^2 \sin \frac{2\pi}{n} = \frac{1}{2}nr^2 \sin \frac{360^\circ}{n}$

7.18. Perimeter =  $2nr \sin \frac{\pi}{n} = 2nr \sin \frac{180^\circ}{n}$




Fig. 7-10

### Regular Polygon of $n$ Sides Circumscribing a Circle of Radius $r$

7.19. Area =  $nr^2 \tan \frac{\pi}{n} = nr^2 \tan \frac{180^\circ}{n}$

7.20. Perimeter =  $2nr \tan \frac{\pi}{n} = 2nr \tan \frac{180^\circ}{n}$




Fig. 7-11

### Segment of Circle of Radius $r$

7.21. Area of shaded part =  $\frac{1}{2}r^2(\theta - \sin \theta)$




Fig. 7-12

### Ellipse of Semi-major Axis $a$ and Semi-minor Axis $b$

7.22. Area =  $\pi ab$

7.23. Perimeter =  $4a \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta$   
 $= 2\pi \sqrt{\frac{1}{2}(a^2 + b^2)}$  [approximately]

where  $k = \sqrt{a^2 - b^2}/a$ . See Table 29 for numerical values.




Fig. 7-13

### Segment of a Parabola

7.24. Area =  $\frac{2}{3}ab$

7.25. Arc length  $ABC = \frac{1}{2}\sqrt{b^2 + 16a^2} + \frac{b^2}{8a} \ln \left( \frac{4a + \sqrt{b^2 + 16a^2}}{b} \right)$




Fig. 7-14


**Rectangular Parallelepiped of Length  $a$ , Height  $b$ , Width  $c$** 7.26. Volume =  $abc$ 7.27. Surface area =  $2(ab + ac + bc)$ 

Fig. 7-15


**Parallelepiped of Cross-sectional Area  $A$  and Height  $h$** 7.28. Volume =  $Ah = abc \sin\theta$ 

Fig. 7-16


**Sphere of Radius  $r$** 7.29. Volume =  $\frac{4}{3}\pi r^3$ 7.30. Surface area =  $4\pi r^2$ 

Fig. 7-17


**Right Circular Cylinder of Radius  $r$  and Height  $h$** 7.31. Volume =  $\pi r^2 h$ 7.32. Lateral surface area =  $2\pi r h$ 

Fig. 7-18


**Circular Cylinder of Radius  $r$  and Slant Height  $l$** 7.33. Volume =  $\pi r^2 h = \pi r^2 l \sin \theta$ 7.34. Lateral surface area =  $2\pi r l = \frac{2\pi r h}{\sin \theta} = 2\pi r h \csc \theta$ 

Fig. 7-19

**Cylinder of Cross-sectional Area  $A$  and Slant Height  $l$** 

7.35. Volume =  $Ah = Al \sin\theta$

7.36. Lateral surface area =  $ph = pl \sin\theta$

Note that formulas 7.31 to 7.34 are special cases of formulas 7.35 and 7.36.




Fig. 7-20

**Right Circular Cone of Radius  $r$  and Height  $h$** 

7.37. Volume =  $\frac{1}{3}\pi r^2 h$

7.38. Lateral surface area =  $\pi r \sqrt{r^2 + h^2} = \pi r l$




Fig. 7-21

**Pyramid of Base Area  $A$  and Height  $h$** 

7.39. Volume =  $\frac{1}{3} Ah$




Fig. 7-22

**Spherical Cap of Radius  $r$  and Height  $h$** 

7.40. Volume (shaded in figure) =  $\frac{1}{3}\pi h^2(3r - h)$

7.41. Surface area =  $2\pi rh$




Fig. 7-23

**Frustum of Right Circular Cone of Radii  $a, b$  and Height  $h$** 

7.42. Volume =  $\frac{1}{3}\pi h(a^2 + ab + b^2)$

7.43. Lateral surface area =  $\pi(a + b)\sqrt{h^2 + (b - a)^2}$   
=  $\pi(a + b)l$




Fig. 7-24

**Spherical Triangle of Angles  $A, B, C$  on Sphere of Radius  $r$** 

7.44. Area of triangle  $ABC = (A + B + C - \pi)r^2$




Fig. 7-25

**Torus of Inner Radius  $a$  and Outer Radius  $b$** 

7.45. Volume =  $\frac{1}{4}\pi^2(a + b)(b - a)^2$

7.46. Surface area =  $\pi^2(b^2 - a^2)$




Fig. 7-26

**Ellipsoid of Semi-axes  $a, b, c$** 

7.47. Volume =  $\frac{4}{3}\pi abc$




Fig. 7-27

**Paraboloid of Revolution**

7.48. Volume =  $\frac{1}{2}\pi b^2a$




Fig. 7-28

# 8

## FORMULAS from PLANE ANALYTIC GEOMETRY

### Distance $d$ Between Two Points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$

$$8.1. \quad d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$




Fig. 8-1

### Slope $m$ of Line Joining Two Points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$

$$8.2. \quad m = \frac{y_2 - y_1}{x_2 - x_1} = \tan \theta$$

### Equation of Line Joining Two Points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$

$$8.3. \quad \frac{y - y_1}{x - x_1} = \frac{y_2 - y_1}{x_2 - x_1} = m \quad \text{or} \quad y - y_1 = m(x - x_1)$$

$$8.4. \quad y = mx + b$$

where  $b = y_1 - mx_1 = \frac{x_2 y_1 - x_1 y_2}{x_2 - x_1}$  is the *intercept* on the  $y$  axis, i.e., the  $y$  *intercept*.

### Equation of Line in Terms of $x$ Intercept $a \neq 0$ and $y$ Intercept $b \neq 0$

$$8.5. \quad \frac{x}{a} + \frac{y}{b} = 1$$




Fig. 8-2

### Normal Form for Equation of Line

8.6.  $x \cos \alpha + y \sin \alpha = p$

where  $p$  = perpendicular distance from origin  $O$  to line  
and  $\alpha$  = angle of inclination of perpendicular  
with positive  $x$  axis.




Fig. 8-3

### General Equation of Line

8.7.  $Ax + By + C = 0$

### Distance from Point $(x_1, y_1)$ to Line $Ax + By + C = 0$

8.8. 
$$\frac{Ax_1 + By_1 + C}{\pm\sqrt{A^2 + B^2}}$$

where the sign is chosen so that the distance is nonnegative.

### Angle $\psi$ Between Two Lines Having Slopes $m_1$ and $m_2$

8.9. 
$$\tan \psi = \frac{m_2 - m_1}{1 + m_1 m_2}$$

Lines are parallel or coincident if and only if  $m_1 = m_2$ .  
Lines are perpendicular if and only if  $m_2 = -1/m_1$ .




Fig. 8-4

### Area of Triangle with Vertices at $(x_1, y_1), (x_2, y_2), (x_3, y_3)$

8.10. 
$$\text{Area} = \pm \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$
  

$$= \pm \frac{1}{2} (x_1 y_2 + y_1 x_3 + y_3 x_2 - y_2 x_3 - y_1 x_2 - x_1 y_3)$$

where the sign is chosen so that the area is nonnegative.  
If the area is zero, the points all lie on a line.




Fig. 8-5

### Transformation of Coordinates Involving Pure Translation

$$8.11. \quad \begin{cases} x = x' + x_0 \\ y = y' + y_0 \end{cases} \quad \text{or} \quad \begin{cases} x' = x - x_0 \\ y' = y - y_0 \end{cases}$$

where  $(x, y)$  are old coordinates (i.e., coordinates relative to  $xy$  system),  $(x', y')$  are new coordinates (relative to  $x', y'$  system), and  $(x_0, y_0)$  are the coordinates of the new origin  $O'$  relative to the old  $xy$  coordinate system.




Fig. 8-6

### Transformation of Coordinates Involving Pure Rotation

$$8.12. \quad \begin{cases} x = x' \cos \alpha - y' \sin \alpha \\ y = x' \sin \alpha + y' \cos \alpha \end{cases} \quad \text{or} \quad \begin{cases} x' = x \cos \alpha + y \sin \alpha \\ y' = y \cos \alpha - x \sin \alpha \end{cases}$$

where the origins of the old  $[xy]$  and new  $[x'y']$  coordinate systems are the same but the  $x'$  axis makes an angle  $\alpha$  with the positive  $x$  axis.




Fig. 8-7

### Transformation of Coordinates Involving Translation and Rotation

$$8.13. \quad \begin{cases} x = x' \cos \alpha - y' \sin \alpha + x_0 \\ y = x' \sin \alpha + y' \cos \alpha + y_0 \end{cases} \quad \text{or} \quad \begin{cases} x' = (x - x_0) \cos \alpha + (y - y_0) \sin \alpha \\ y' = (y - y_0) \cos \alpha - (x - x_0) \sin \alpha \end{cases}$$

where the new origin  $O'$  of  $x'y'$  coordinate system has coordinates  $(x_0, y_0)$  relative to the old  $xy$  coordinate system and the  $x'$  axis makes an angle  $\alpha$  with the positive  $x$  axis.




Fig. 8-8

### Polar Coordinates $(r, \theta)$

A point  $P$  can be located by rectangular coordinates  $(x, y)$  or polar coordinates  $(r, \theta)$ . The transformation between these coordinates is as follows:

$$8.14. \quad \begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases} \quad \text{or} \quad \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1}(y/x) \end{cases}$$




Fig. 8-9

**Equation of Circle of Radius  $R$ , Center at  $(x_0, y_0)$** 

8.15.  $(x - x_0)^2 + (y - y_0)^2 = R^2$




Fig. 8-10

**Equation of Circle of Radius  $R$  Passing Through Origin**

8.16.  $r = 2R \cos(\theta - \alpha)$

where  $(r, \theta)$  are polar coordinates of any point on the circle and  $(R, \alpha)$  are polar coordinates of the center of the circle.




Fig. 8-11

**Conics (Ellipse, Parabola, or Hyperbola)**

If a point  $P$  moves so that its distance from a fixed point (called the *focus*) divided by its distance from a fixed line (called the *directrix*) is a constant  $\epsilon$  (called the *eccentricity*), then the curve described by  $P$  is called a *conic* (so-called because such curves can be obtained by intersecting a plane and a cone at different angles).

If the focus is chosen at origin  $O$ , the equation of a conic in polar coordinates  $(r, \theta)$  is, if  $OQ = p$  and  $LM = D$  (see Fig. 8-12),

8.17.  $r = \frac{p}{1 - \epsilon \cos \theta} = \frac{\epsilon D}{1 - \epsilon \cos \theta}$

The conic is

- (i) an ellipse if  $\epsilon < 1$
- (ii) a parabola if  $\epsilon = 1$
- (iii) a hyperbola if  $\epsilon > 1$




Fig. 8-12

**Ellipse with Center  $C(x_0, y_0)$  and Major Axis Parallel to  $x$  Axis**

**8.18.** Length of major axis  $A'A = 2a$

**8.19.** Length of minor axis  $B'B = 2b$

**8.20.** Distance from center  $C$  to focus  $F$  or  $F'$  is

$$c = \sqrt{a^2 - b^2}$$

$$\text{8.21. Eccentricity } \epsilon = \frac{c}{a} = \frac{\sqrt{a^2 - b^2}}{a}$$




Fig. 8-13

**8.22.** Equation in rectangular coordinates:

$$\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} = 1$$

$$\text{8.23. Equation in polar coordinates if } C \text{ is at } O: r^2 = \frac{a^2 b^2}{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$$

$$\text{8.24. Equation in polar coordinates if } C \text{ is on } x \text{ axis and } F' \text{ is at } O: r = \frac{a(1 - \epsilon^2)}{1 - \epsilon \cos \theta}$$

**8.25.** If  $P$  is any point on the ellipse,  $PF + PF' = 2a$

If the major axis is parallel to the  $y$  axis, interchange  $x$  and  $y$  in the above or replace  $\theta$  by  $\frac{1}{2}\pi - \theta$  (or  $90^\circ - \theta$ ).

**Parabola with Axis Parallel to  $x$  Axis**

If vertex is at  $A(x_0, y_0)$  and the distance from  $A$  to focus  $F$  is  $a > 0$ , the equation of the parabola is

$$\text{8.26. } (y - y_0)^2 = 4a(x - x_0) \quad \text{if parabola opens to right (Fig. 8-14)}$$

$$\text{8.27. } (y - y_0)^2 = -4a(x - x_0) \quad \text{if parabola opens to left (Fig. 8-15)}$$

If focus is at the origin (Fig. 8-16), the equation in polar coordinates is

$$\text{8.28. } r = \frac{2a}{1 - \cos \theta}$$




Fig. 8-14




Fig. 8-15




Fig. 8-16

In case the axis is parallel to the  $y$  axis, interchange  $x$  and  $y$  or replace  $\theta$  by  $\frac{1}{2}\pi - \theta$  (or  $90^\circ - \theta$ ).


**Hyperbola with Center  $C(x_0, y_0)$  and Major Axis Parallel to  $x$  Axis**

Fig. 8-17

**8.29.** Length of major axis  $A'A = 2a$

**8.30.** Length of minor axis  $B'B = 2b$

**8.31.** Distance from center  $C$  to focus  $F$  or  $F' = c = \sqrt{a^2 + b^2}$

**8.32.** Eccentricity  $\epsilon = \frac{c}{a} = \frac{\sqrt{a^2 + b^2}}{a}$

**8.33.** Equation in rectangular coordinates:  $\frac{(x - x_0)^2}{a^2} - \frac{(y - y_0)^2}{b^2} = 1$

**8.34.** Slopes of asymptotes  $G'H$  and  $GH' = \pm \frac{b}{a}$

**8.35.** Equation in polar coordinates if  $C$  is at  $O$ :  $r^2 = \frac{a^2 b^2}{b^2 \cos^2 \theta - a^2 \sin^2 \theta}$

**8.36.** Equation in polar coordinates if  $C$  is on  $x$  axis and  $F'$  is at  $O$ :  $r = \frac{a(\epsilon^2 - 1)}{1 - \epsilon \cos \theta}$

**8.37.** If  $P$  is any point on the hyperbola,  $PF - PF' = \pm 2a$  (depending on branch)

If the major axis is parallel to the  $y$  axis, interchange  $x$  and  $y$  in the above or replace  $\theta$  by  $\frac{1}{2}\pi - \theta$  (or  $90^\circ - \theta$ ).

# 9

## SPECIAL PLANE CURVES

### Lemniscate

9.1. Equation in polar coordinates:

$$r^2 = a^2 \cos 2\theta$$

9.2. Equation in rectangular coordinates:

$$(x^2 + y^2)^2 = a^2(x^2 - y^2)$$

9.3. Angle between  $AB'$  or  $A'B$  and  $x$  axis =  $45^\circ$

9.4. Area of one loop =  $a^2$




Fig. 9-1

### Cycloid

9.5. Equations in parametric form:

$$\begin{cases} x = a(\phi - \sin \phi) \\ y = a(1 - \cos \phi) \end{cases}$$

9.6. Area of one arch =  $3\pi a^2$

9.7. Arc length of one arch =  $8a$

This is a curve described by a point  $P$  on a circle of radius  $a$  rolling along  $x$  axis.




Fig. 9-2

### Hypocycloid with Four Cusps

9.8. Equation in rectangular coordinates:

$$x^{2/3} + y^{2/3} = a^{2/3}$$

9.9. Equations in parametric form:

$$\begin{cases} x = a \cos^3 \theta \\ y = a \sin^3 \theta \end{cases}$$

9.10. Area bounded by curve =  $\frac{3}{8}\pi a^2$

9.11. Arc length of entire curve =  $6a$

This is a curve described by a point  $P$  on a circle of radius  $a/4$  as it rolls on the inside of a circle of radius  $a$ .




Fig. 9-3

### Cardioid

**9.12.** Equation:  $r = 2a(1 + \cos \theta)$

**9.13.** Area bounded by curve =  $6\pi a^2$

**9.14.** Arc length of curve =  $16a$

This is the curve described by a point  $P$  of a circle of radius  $a$  as it rolls on the outside of a fixed circle of radius  $a$ . The curve is also a special case of the limacon of Pascal (see 9.32).




Fig. 9-4

### Catenary

**9.15.** Equation:  $y = \frac{a}{2}(e^{x/a} + e^{-x/a}) = a \cosh \frac{x}{a}$

This is the curve in which a heavy uniform chain would hang if suspended vertically from fixed points  $A$  and  $B$ .




Fig. 9-5

### Three-Leaved Rose

**9.16.** Equation:  $r = a \cos 3\theta$

The equation  $r = a \sin 3\theta$  is a similar curve obtained by rotating the curve of Fig. 9-6 counterclockwise through  $30^\circ$  or  $\pi/6$  radians.

In general,  $r = a \cos n\theta$  or  $r = a \sin n\theta$  has  $n$  leaves if  $n$  is odd.




Fig. 9-6

### Four-Leaved Rose

**9.17.** Equation:  $r = a \cos 2\theta$

The equation  $r = a \sin 2\theta$  is a similar curve obtained by rotating the curve of Fig. 9-7 counterclockwise through  $45^\circ$  or  $\pi/4$  radians.

In general,  $r = a \cos n\theta$  or  $r = a \sin n\theta$  has  $2n$  leaves if  $n$  is even.




Fig. 9-7

## Epicycloid

9.18. Parametric equations:

$$\begin{cases} x = (a+b) \cos \theta - b \cos\left(\frac{a+b}{b}\right)\theta \\ y = (a+b) \sin \theta - b \sin\left(\frac{a+b}{b}\right)\theta \end{cases}$$

This is the curve described by a point  $P$  on a circle of radius  $b$  as it rolls on the outside of a circle of radius  $a$ .

The cardioid (Fig. 9-4) is a special case of an epicycloid.




Fig. 9-8

## General Hypocycloid

9.19. Parametric equations:

$$\begin{cases} x = (a-b) \cos \phi + b \cos\left(\frac{a-b}{b}\right)\phi \\ y = (a-b) \sin \phi - b \sin\left(\frac{a-b}{b}\right)\phi \end{cases}$$

This is the curve described by a point  $P$  on a circle of radius  $b$  as it rolls on the inside of a circle of radius  $a$ .

If  $b = a/4$ , the curve is that of Fig. 9-3.




Fig. 9-9

## Trochoid

9.20. Parametric equations:  $\begin{cases} x = a\phi - b \sin \phi \\ y = a - b \cos \phi \end{cases}$

This is the curve described by a point  $P$  at distance  $b$  from the center of a circle of radius  $a$  as the circle rolls on the  $x$  axis.

If  $b < a$ , the curve is as shown in Fig. 9-10 and is called a *curtate cycloid*.

If  $b > a$ , the curve is as shown in Fig. 9-11 and is called a *prolate cycloid*.

If  $b = a$ , the curve is the cycloid of Fig. 9-2.




Fig. 9-10




Fig. 9-11

### Tractrix

9.21. Parametric equations:  $\begin{cases} x = a(\ln \cot \frac{1}{2}\phi - \cos \phi) \\ y = a \sin \phi \end{cases}$

This is the curve described by endpoint  $P$  of a taut string  $PQ$  of length  $a$  as the other end  $Q$  is moved along the  $x$  axis.




Fig. 9-12

### Witch of Agnesi

9.22. Equation in rectangular coordinates:  $y = \frac{8a^3}{x^2 + 4a^2}$

9.23. Parametric equations:  $\begin{cases} x = 2a \cot \theta \\ y = a(1 - \cos 2\theta) \end{cases}$

In Fig. 9-13 the variable line  $QA$  intersects  $y = 2a$  and the circle of radius  $a$  with center  $(0, a)$  at  $A$  and  $B$ , respectively. Any point  $P$  on the “witch” is located by constructing lines parallel to the  $x$  and  $y$  axes through  $B$  and  $A$ , respectively, and determining the point  $P$  of intersection.




Fig. 9-13

### Folium of Descartes

9.24. Equation in rectangular coordinates:

$$x^3 + y^3 = 3axy$$

9.25. Parametric equations:

$$\begin{cases} x = \frac{3at}{1+t^3} \\ y = \frac{3at^2}{1+t^3} \end{cases}$$

9.26. Area of loop =  $\frac{3}{2}a^2$




Fig. 9-14

9.27. Equation of asymptote:  $x + y + a = 0$

### Involute of a Circle

9.28. Parametric equations:

$$\begin{cases} x = a(\cos \phi + \phi \sin \phi) \\ y = a(\sin \phi - \phi \cos \phi) \end{cases}$$

This is the curve described by the endpoint  $P$  of a string as it unwinds from a circle of radius  $a$  while held taut.




Fig. 9-15

### Evolute of an Ellipse

9.29. Equation in rectangular coordinates:

$$(ax)^{2/3} + (by)^{2/3} = (a^2 - b^2)^{2/3}$$

9.30. Parametric equations:

$$\begin{cases} ax = (a^2 - b^2) \cos^3 \theta \\ by = (a^2 - b^2) \sin^3 \theta \end{cases}$$

This curve is the envelope of the normals to the ellipse  $x^2/a^2 + y^2/b^2 = 1$  shown dashed in Fig. 9-16.




Fig. 9-16

### Ovals of Cassini

9.31. Polar equation:  $r^4 + a^4 - 2a^2r^2 \cos 2\theta = b^4$

This is the curve described by a point  $P$  such that the product of its distance from two fixed points (distance  $2a$  apart) is a constant  $b^2$ .

The curve is as in Fig. 9-17 or Fig. 9-18 according as  $b < a$  or  $b > a$ , respectively.

If  $b = a$ , the curve is a *lemniscate* (Fig. 9-1).




Fig. 9-17




Fig. 9-18

### Limaçon of Pascal

9.32. Polar equation:  $r = b + a \cos \theta$

Let  $OQ$  be a line joining origin  $O$  to any point  $Q$  on a circle of diameter  $a$  passing through  $O$ . Then the curve is the locus of all points  $P$  such that  $PQ = b$ .

The curve is as in Fig. 9-19 or Fig. 9-20 according as  $2a > b > a$  or  $b < a$ , respectively. If  $b = a$ , the curve is a *cardioid* (Fig. 9-4). If  $b \geq 2a$ , the curve is convex.




Fig. 9-19




Fig. 9-20

### Cissoid of Diocles

9.33. Equation in rectangular coordinates:

$$y^2 = \frac{x^2}{2a - x}$$

9.34. Parametric equations:

$$\begin{cases} x = 2a \sin^2 \theta \\ y = \frac{2a \sin^3 \theta}{\cos \theta} \end{cases}$$

This is the curve described by a point  $P$  such that the distance  $OP = \text{distance } RS$ . It is used in the problem of *duplication of a cube*, i.e., finding the side of a cube which has twice the volume of a given cube.




Fig. 9-21

### Spiral of Archimedes

9.35. Polar equation:  $r = a\theta$




Fig. 9-22

# 10

## FORMULAS from SOLID ANALYTIC GEOMETRY

### Distance $d$ Between Two Points $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$

$$10.1. \quad d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$




Fig. 10-1

### Direction Cosines of Line Joining Points $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$

$$10.2. \quad l = \cos \alpha = \frac{x_2 - x_1}{d}, \quad m = \cos \beta = \frac{y_2 - y_1}{d}, \quad n = \cos \gamma = \frac{z_2 - z_1}{d}$$

where  $\alpha, \beta, \gamma$  are the angles that line  $P_1P_2$  makes with the positive  $x, y, z$  axes, respectively, and  $d$  is given by 10.1 (see Fig. 10-1).

### Relationship Between Direction Cosines

$$10.3. \quad \cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1 \quad \text{or} \quad l^2 + m^2 + n^2 = 1$$

### Direction Numbers

Numbers  $L, M, N$ , which are proportional to the direction cosines  $l, m, n$ , are called *direction numbers*. The relationship between them is given by

$$10.4. \quad l = \frac{L}{\sqrt{L^2 + M^2 + N^2}}, \quad m = \frac{M}{\sqrt{L^2 + M^2 + N^2}}, \quad n = \frac{N}{\sqrt{L^2 + M^2 + N^2}}$$

### Equations of Line Joining $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$ in Standard Form

$$10.5. \quad \frac{x - x_1}{x_2 - x_1} = \frac{y - y_1}{y_2 - y_1} = \frac{z - z_1}{z_2 - z_1} \quad \text{or} \quad \frac{x - x_1}{l} = \frac{y - y_1}{m} = \frac{z - z_1}{n}$$

These are also valid if  $l, m, n$  are replaced by  $L, M, N$ , respectively.

**Equations of Line Joining  $P_1(x_1, y_1, z_1)$  and  $P_2(x_2, y_2, z_2)$  in Parametric Form**

---

**10.6.**  $x = x_1 + lt, y = y_1 + mt, z = z_1 + nt$

These are also valid if  $l, m, n$  are replaced by  $L, M, N$ , respectively.

**Angle  $\phi$  Between Two Lines with Direction Cosines  $l_1, m_1, n_1$  and  $l_2, m_2, n_2$** 

---

**10.7.**  $\cos\phi = l_1l_2 + m_1m_2 + n_1n_2$

**General Equation of a Plane**

---

**10.8.**  $Ax + By + Cz + D = 0$

( $A, B, C, D$  are constants)

**Equation of Plane Passing Through Points  $(x_1, y_1, z_1), (x_2, y_2, z_2), (x_3, y_3, z_3)$** 

---

**10.9.** 
$$\begin{vmatrix} x - x_1 & y - y_1 & z - z_1 \\ x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ x_3 - x_1 & y_3 - y_1 & z_3 - z_1 \end{vmatrix} = 0$$

or

---

**10.10.** 
$$\begin{vmatrix} y_2 - y_1 & z_2 - z_1 \\ y_3 - y_1 & z_3 - z_1 \end{vmatrix} (x - x_1) + \begin{vmatrix} z_2 - z_1 & x_2 - x_1 \\ z_3 - z_1 & x_3 - x_1 \end{vmatrix} (y - y_1) + \begin{vmatrix} x_2 - x_1 & y_2 - y_1 \\ x_3 - x_1 & y_3 - y_1 \end{vmatrix} (z - z_1) = 0$$

**Equation of Plane in Intercept Form**

---

**10.11.** 
$$\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$$

where  $a, b, c$  are the intercepts on the  $x, y, z$  axes, respectively.




Fig. 10-2

**Equations of Line Through  $(x_0, y_0, z_0)$  and Perpendicular to Plane  $Ax + By + Cz + D = 0$** 

---

**10.12.** 
$$\frac{x - x_0}{A} = \frac{y - y_0}{B} = \frac{z - z_0}{C} \quad \text{or} \quad x = x_0 + At, y = y_0 + Bt, z = z_0 + Ct$$

Note that the direction numbers for a line perpendicular to the plane  $Ax + By + Cz + D = 0$  are  $A, B, C$ .

**Distance from Point  $(x_0, y_0, z_0)$  to Plane  $Ax + By + Cz + D = 0$** 

$$10.13. \quad \frac{Ax_0 + By_0 + Cz_0 + D}{\pm\sqrt{A^2 + B^2 + C^2}}$$

where the sign is chosen so that the distance is nonnegative.

**Normal Form for Equation of Plane**

$$10.14. \quad x \cos \alpha + y \cos \beta + z \cos \gamma = p$$

where  $p$  = perpendicular distance from  $O$  to plane at  $P$   
and  $\alpha, \beta, \gamma$  are angles between  $OP$  and positive  $x, y, z$  axes.




Fig. 10-3

**Transformation of Coordinates Involving Pure Translation**

$$10.15. \quad \begin{cases} x = x' + x_0 \\ y = y' + y_0 \\ z = z' + z_0 \end{cases} \quad \text{or} \quad \begin{cases} x' = x - x_0 \\ y' = y - y_0 \\ z' = z - z_0 \end{cases}$$

where  $(x, y, z)$  are old coordinates (i.e., coordinates relative to  $xyz$  system),  $(x', y', z')$  are new coordinates (relative to  $x'y'z'$  system) and  $(x_0, y_0, z_0)$  are the coordinates of the new origin  $O'$  relative to the old  $xyz$  coordinate system.




Fig. 10-4

**Transformation of Coordinates Involving Pure Rotation**

$$10.16. \quad \begin{cases} x = l_1 x' + l_2 y' + l_3 z' \\ y = m_1 x' + m_2 y' + m_3 z' \\ z = n_1 x' + n_2 y' + n_3 z' \end{cases}$$

or  $\begin{cases} x' = l_1 x + m_1 y + n_1 z \\ y' = l_2 x + m_2 y + n_2 z \\ z' = l_3 x + m_3 y + n_3 z \end{cases}$

where the origins of the  $xyz$  and  $x'y'z'$  systems are the same and  $l_1, m_1, n_1; l_2, m_2, n_2; l_3, m_3, n_3$  are the direction cosines of the  $x', y', z'$  axes relative to the  $x, y, z$  axes, respectively.




Fig. 10-5

### Transformation of Coordinates Involving Translation and Rotation

**10.17.** 
$$\begin{cases} x = l_1x' + l_2y' + l_3z' + x_0 \\ y = m_1x' + m_2y' + m_3z' + y_0 \\ z = n_1x' + n_2y' + n_3z' + z_0 \end{cases}$$

or 
$$\begin{cases} x' = l_1(x - x_0) + m_1(y - y_0) + n_1(z - z_0) \\ y' = l_2(x - x_0) + m_2(y - y_0) + n_2(z - z_0) \\ z' = l_3(x - x_0) + m_3(y - y_0) + n_3(z - z_0) \end{cases}$$

where the origin  $O'$  of the  $x'y'z'$  system has coordinates  $(x_0, y_0, z_0)$  relative to the  $xyz$  system and

$$l_1, m_1, n_1; l_2, m_2, n_2; l_3, m_3, n_3$$

are the direction cosines of the  $x', y', z'$  axes relative to the  $x, y, z$  axes, respectively.




Fig. 10-6

### Cylindrical Coordinates $(r, \theta, z)$

A point  $P$  can be located by cylindrical coordinates  $(r, \theta, z)$  (see Fig. 10-7) as well as rectangular coordinates  $(x, y, z)$ .

The transformation between these coordinates is

**10.18.** 
$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases} \quad \text{or} \quad \begin{cases} r = \sqrt{x^2 + y^2} \\ \theta = \tan^{-1}(y/x) \\ z = z \end{cases}$$




Fig. 10-7

### Spherical Coordinates $(r, \theta, \phi)$

A point  $P$  can be located by spherical coordinates  $(r, \theta, \phi)$  (see Fig. 10-8) as well as rectangular coordinates  $(x, y, z)$ .

The transformation between those coordinates is

**10.19.** 
$$\begin{cases} x = r \sin \theta \cos \phi \\ y = r \sin \theta \sin \phi \\ z = r \cos \theta \end{cases}$$

or 
$$\begin{cases} r = \sqrt{x^2 + y^2 + z^2} \\ \theta = \tan^{-1}(y/x) \\ \phi = \cos^{-1}(z/\sqrt{x^2 + y^2 + z^2}) \end{cases}$$




Fig. 10-8

**Equation of Sphere in Rectangular Coordinates**

---

**10.20.**  $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = R^2$

where the sphere has center  $(x_0, y_0, z_0)$  and radius  $R$ .




Fig. 10-9

**Equation of Sphere in Cylindrical Coordinates**

---

**10.21.**  $r^2 - 2r_0 r \cos(\theta - \theta_0) + r_0^2 + (z - z_0)^2 = R^2$

where the sphere has center  $(r_0, \theta_0, z_0)$  in cylindrical coordinates and radius  $R$ .

If the center is at the origin the equation is

**10.22.**  $r^2 + z^2 = R^2$

**Equation of Sphere in Spherical Coordinates**

---

**10.23.**  $r^2 + r_0^2 - 2r_0 r \sin \theta \sin \theta_0 \cos(\phi - \phi_0) = R^2$

where the sphere has center  $(r_0, \theta_0, \phi_0)$  in spherical coordinates and radius  $R$ .

If the center is at the origin the equation is

**10.24.**  $r = R$

**Equation of Ellipsoid with Center  $(x_0, y_0, z_0)$  and Semi-axes  $a, b, c$** 

---

**10.25.**  $\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} + \frac{(z - z_0)^2}{c^2} = 1$




Fig. 10-10

**Elliptic Cylinder with Axis as  $z$  Axis**

$$10.26. \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

where  $a, b$  are semi-axes of elliptic cross-section.

If  $b = a$  it becomes a circular cylinder of radius  $a$ .




Fig. 10-11

**Elliptic Cone with Axis as  $z$  Axis**

$$10.27. \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$




Fig. 10-12

**Hyperboloid of One Sheet**

$$10.28. \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$




Fig. 10-13

**Hyperboloid of Two Sheets**

$$10.29. \quad \frac{x^2}{a^2} - \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$$

Note orientation of axes in Fig. 10-14.




Fig. 10-14

**Elliptic Paraboloid**

$$10.30. \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z}{c}$$




Fig. 10-15

**Hyperbolic Paraboloid**

$$10.31. \quad \frac{x^2}{a^2} - \frac{y^2}{b^2} = \frac{z}{c}$$

Note orientation of axes in Fig. 10-16.




Fig. 10-16

# 11 SPECIAL MOMENTS of INERTIA

The table below shows the moments of inertia of various rigid bodies of mass  $M$ . In all cases it is assumed the body has uniform (i.e., constant) density.

TYPE OF RIGID BODY	MOMENT OF INERTIA
<b>11.1.</b> Thin rod of length $a$	$\frac{1}{12}Ma^2$ $\frac{1}{3}Ma^2$
(a) about axis perpendicular to the rod through the center of mass (b) about axis perpendicular to the rod through one end	
<b>11.2.</b> Rectangular parallelepiped with sides $a, b, c$	$\frac{1}{12}M(a^2 + b^2)$ $\frac{1}{12}M(4a^2 + b^2)$
(a) about axis parallel to $c$ and through center of face $ab$ (b) about axis through center of face $bc$ and parallel to $c$	
<b>11.3.</b> Thin rectangular plate with sides $a, b$	$\frac{1}{12}M(a^2 + b^2)$ $\frac{1}{12}Ma^2$
(a) about axis perpendicular to the plate through center (b) about axis parallel to side $b$ through center	
<b>11.4.</b> Circular cylinder of radius $a$ and height $h$	$\frac{1}{2}Ma^2$ $\frac{1}{12}M(h^2 + 3a^2)$ $\frac{1}{12}M(4h^2 + 3a^2)$
(a) about axis of cylinder (b) about axis through center of mass and perpendicular to cylindrical axis (c) about axis coinciding with diameter at one end	
<b>11.5.</b> Hollow circular cylinder of outer radius $a$ , inner radius $b$ and height $h$	
(a) about axis of cylinder (b) about axis through center of mass and perpendicular to cylindrical axis (c) about axis coinciding with diameter at one end	$\frac{1}{2}M(a^2 + b^2)$ $\frac{1}{12}M(3a^2 + 3b^2 + h^2)$ $\frac{1}{12}M(3a^2 + 3b^2 + 4h^2)$
<b>11.6.</b> Circular plate of radius $a$	$\frac{1}{2}Ma^2$ $\frac{1}{4}Ma^2$
(a) about axis perpendicular to plate through center (b) about axis coinciding with a diameter	

<b>11.7.</b>	Hollow circular plate or ring with outer radius $a$ and inner radius $b$	
(a)	about axis perpendicular to plane of plate through center	$\frac{1}{2}M(a^2 + b^2)$
(b)	about axis coinciding with a diameter	$\frac{1}{4}M(a^2 + b^2)$
<b>11.8.</b>	Thin circular ring of radius $a$	
(a)	about axis perpendicular to plane of ring through center	$Ma^2$
(b)	about axis coinciding with diameter	$\frac{1}{2}Ma^2$
<b>11.9.</b>	Sphere of radius $a$	
(a)	about axis coinciding with a diameter	$\frac{2}{5}Ma^2$
(b)	about axis tangent to the surface	$\frac{7}{5}Ma^2$
<b>11.10.</b>	Hollow sphere of outer radius $a$ and inner radius $b$	
(a)	about axis coinciding with a diameter	$\frac{2}{5}M(a^5 - b^5)/(a^3 - b^3)$
(b)	about axis tangent to the surface	$\frac{2}{5}M(a^5 - b^5)/(a^3 - b^3) + Ma^2$
<b>11.11.</b>	Hollow spherical shell of radius $a$	
(a)	about axis coinciding with a diameter	$\frac{2}{3}Ma^2$
(b)	about axis tangent to the surface	$\frac{5}{3}Ma^2$
<b>11.12.</b>	Ellipsoid with semi-axes $a, b, c$	
(a)	about axis coinciding with semi-axis $c$	$\frac{1}{5}M(a^2 + b^2)$
(b)	about axis tangent to surface, parallel to semi-axis $c$ and at distance $a$ from center	$\frac{1}{5}M(6a^2 + b^2)$
<b>11.13.</b>	Circular cone of radius $a$ and height $h$	
(a)	about axis of cone	$\frac{3}{10}Ma^2$
(b)	about axis through vertex and perpendicular to axis	$\frac{3}{20}M(a^2 + 4h^2)$
(c)	about axis through center of mass and perpendicular to axis	$\frac{3}{80}M(4a^2 + h^2)$
<b>11.14.</b>	Torus with outer radius $a$ and inner radius $b$	
(a)	about axis through center of mass and perpendicular to the plane of torus	$\frac{1}{4}M(7a^2 - 6ab + 3b^2)$
(b)	about axis through center of mass and in the plane of torus	$\frac{1}{4}M(9a^2 - 10ab + 5b^2)$

## Section III: Elementary Transcendental Functions

# 12 TRIGONOMETRIC FUNCTIONS

### Definition of Trigonometric Functions for a Right Triangle

Triangle  $ABC$  has a right angle ( $90^\circ$ ) at  $C$  and sides of length  $a, b, c$ . The trigonometric functions of angle  $A$  are defined as follows:

$$12.1. \text{ sine of } A = \sin A = \frac{a}{c} = \frac{\text{opposite}}{\text{hypotenuse}}$$

$$12.2. \text{ cosine of } A = \cos A = \frac{b}{c} = \frac{\text{adjacent}}{\text{hypotenuse}}$$

$$12.3. \text{ tangent of } A = \tan A = \frac{a}{b} = \frac{\text{opposite}}{\text{adjacent}}$$

$$12.4. \text{ cotangent of } A = \cot A = \frac{b}{a} = \frac{\text{adjacent}}{\text{opposite}}$$

$$12.5. \text{ secant of } A = \sec A = \frac{c}{b} = \frac{\text{hypotenuse}}{\text{adjacent}}$$

$$12.6. \text{ cosecant of } A = \csc A = \frac{c}{a} = \frac{\text{hypotenuse}}{\text{opposite}}$$




Fig. 12-1

### Extensions to Angles Which May be Greater Than $90^\circ$

Consider an  $xy$  coordinate system (see Figs. 12-2 and 12-3). A point  $P$  in the  $xy$  plane has coordinates  $(x, y)$  where  $x$  is considered as positive along  $OX$  and negative along  $OX'$  while  $y$  is positive along  $OY$  and negative along  $OY'$ . The distance from origin  $O$  to point  $P$  is positive and denoted by  $r = \sqrt{x^2 + y^2}$ . The angle  $A$  described *counterclockwise* from  $OX$  is considered *positive*. If it is described *clockwise* from  $OX$  it is considered *negative*. We call  $X'OX$  and  $Y'OY$  the *x* and *y* axis, respectively.

The various quadrants are denoted by I, II, III, and IV called the first, second, third, and fourth quadrants, respectively. In Fig. 12-2, for example, angle  $A$  is in the second quadrant while in Fig. 12-3 angle  $A$  is in the third quadrant.




Fig. 12-2




Fig. 12-3

For an angle  $A$  in any quadrant, the trigonometric functions of  $A$  are defined as follows.

$$12.7. \sin A = y/r$$

$$12.8. \cos A = x/r$$

$$12.9. \tan A = y/x$$

$$12.10. \cot A = x/y$$

$$12.11. \sec A = r/x$$

$$12.12. \csc A = r/y$$

### Relationship Between Degrees and Radians

A *radian* is that angle  $\theta$  subtended at center  $O$  of a circle by an arc  $MN$  equal to the radius  $r$ .

Since  $2\pi$  radians =  $360^\circ$  we have

$$12.13. 1 \text{ radian} = 180^\circ/\pi = 57.29577\ 95130\ 8232 \dots^\circ$$

$$12.14. 1^\circ = \pi/180 \text{ radians} = 0.01745\ 32925\ 19943\ 29576\ 92 \dots \text{ radians}$$




Fig. 12-4

### Relationships Among Trigonometric Functions

$$12.15. \tan A = \frac{\sin A}{\cos A}$$

$$12.19. \sin^2 A + \cos^2 A = 1$$

$$12.16. \cot A = \frac{1}{\tan A} = \frac{\cos A}{\sin A}$$

$$12.20. \sec^2 A - \tan^2 A = 1$$

$$12.17. \sec A = \frac{1}{\cos A}$$

$$12.21. \csc^2 A - \cot^2 A = 1$$

$$12.18. \csc A = \frac{1}{\sin A}$$

### Signs and Variations of Trigonometric Functions

Quadrant	$\sin A$	$\cos A$	$\tan A$	$\cot A$	$\sec A$	$\csc A$
I	+	+	+	+	+	+
	0 to 1	1 to 0	0 to $\infty$	$\infty$ to 0	1 to $\infty$	$\infty$ to 1
II	+	-	-	-	-	+
	1 to 0	0 to -1	$-\infty$ to 0	0 to $-\infty$	$-\infty$ to -1	1 to $\infty$
III	-	-	+	+	-	-
	0 to -1	-1 to 0	0 to $\infty$	$\infty$ to 0	-1 to $-\infty$	$-\infty$ to -1
IV	-	+	-	-	+	-
	-1 to 0	0 to 1	$-\infty$ to 0	0 to $-\infty$	$\infty$ to 1	-1 to $-\infty$

**Exact Values for Trigonometric Functions of Various Angles**

Angle $A$ in degrees	Angle $A$ in radians	$\sin A$	$\cos A$	$\tan A$	$\cot A$	$\sec A$	$\csc A$
0°	0	0	1	0	∞	1	∞
15°	$\pi/12$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$2-\sqrt{3}$	$2+\sqrt{3}$	$\sqrt{6}-\sqrt{2}$	$\sqrt{6}+\sqrt{2}$
30°	$\pi/6$	$\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	2
45°	$\pi/4$	$\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	1	1	$\sqrt{2}$	$\sqrt{2}$
60°	$\pi/3$	$\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	2	$\frac{2}{3}\sqrt{3}$
75°	$5\pi/12$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$2+\sqrt{3}$	$2-\sqrt{3}$	$\sqrt{6}+\sqrt{2}$	$\sqrt{6}-\sqrt{2}$
90°	$\pi/2$	1	0	±∞	0	±∞	1
105°	$7\pi/12$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-(2+\sqrt{3})$	$-(2-\sqrt{3})$	$-(\sqrt{6}+\sqrt{2})$	$\sqrt{6}-\sqrt{2}$
120°	$2\pi/3$	$\frac{1}{2}\sqrt{3}$	$-\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	-2	$\frac{2}{3}\sqrt{3}$
135°	$3\pi/4$	$\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	-1	-1	$-\sqrt{2}$	$\sqrt{2}$
150°	$5\pi/6$	$\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	2
165°	$11\pi/12$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-(2-\sqrt{3})$	$-(2+\sqrt{3})$	$-(\sqrt{6}-\sqrt{2})$	$\sqrt{6}+\sqrt{2}$
180°	$\pi$	0	-1	0	±∞	-1	±∞
195°	$13\pi/12$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$2-\sqrt{3}$	$2+\sqrt{3}$	$-(\sqrt{6}-\sqrt{2})$	$-(\sqrt{6}+\sqrt{2})$
210°	$7\pi/6$	$-\frac{1}{2}$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	$\sqrt{3}$	$-\frac{2}{3}\sqrt{3}$	-2
225°	$5\pi/4$	$-\frac{1}{2}\sqrt{2}$	$-\frac{1}{2}\sqrt{2}$	1	1	$-\sqrt{2}$	$-\sqrt{2}$
240°	$4\pi/3$	$-\frac{1}{2}\sqrt{3}$	$-\frac{1}{2}$	$\sqrt{3}$	$\frac{1}{3}\sqrt{3}$	-2	$-\frac{2}{3}\sqrt{3}$
255°	$17\pi/12$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$2+\sqrt{3}$	$2-\sqrt{3}$	$-(\sqrt{6}+\sqrt{2})$	$-(\sqrt{6}-\sqrt{2})$
270°	$3\pi/2$	-1	0	±∞	0	±∞	-1
285°	$19\pi/12$	$-\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$-(2+\sqrt{3})$	$-(2-\sqrt{3})$	$\sqrt{6}+\sqrt{2}$	$-(\sqrt{6}-\sqrt{2})$
300°	$5\pi/3$	$-\frac{1}{2}\sqrt{3}$	$\frac{1}{2}$	$-\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	2	$-\frac{2}{3}\sqrt{3}$
315°	$7\pi/4$	$-\frac{1}{2}\sqrt{2}$	$\frac{1}{2}\sqrt{2}$	-1	-1	$\sqrt{2}$	$-\sqrt{2}$
330°	$11\pi/6$	$-\frac{1}{2}$	$\frac{1}{2}\sqrt{3}$	$-\frac{1}{3}\sqrt{3}$	$-\sqrt{3}$	$\frac{2}{3}\sqrt{3}$	-2
345°	$23\pi/12$	$-\frac{1}{4}(\sqrt{6}-\sqrt{2})$	$\frac{1}{4}(\sqrt{6}+\sqrt{2})$	$-(2-\sqrt{3})$	$-(2+\sqrt{3})$	$\sqrt{6}-\sqrt{2}$	$-(\sqrt{6}+\sqrt{2})$
360°	$2\pi$	0	1	0	±∞	1	±∞

For other angles see Tables 2, 3, and 4.

### Graphs of Trigonometric Functions

In each graph  $x$  is in radians.

**12.22.**  $y = \sin x$




Fig. 12-5

**12.23.**  $y = \cos x$




Fig. 12-6

**12.24.**  $y = \tan x$




Fig. 12-7

**12.25.**  $y = \cot x$




Fig. 12-8

**12.26.**  $y = \sec x$




Fig. 12-9

**12.27.**  $y = \csc x$




Fig. 12-10

### Functions of Negative Angles

**12.28.**  $\sin(-A) = -\sin A$

**12.29.**  $\cos(-A) = \cos A$

**12.30.**  $\tan(-A) = -\tan A$

**12.31.**  $\csc(-A) = -\csc A$

**12.32.**  $\sec(-A) = \sec A$

**12.33.**  $\cot(-A) = -\cot A$

**Addition Formulas**

**12.34.**  $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$

**12.35.**  $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$

**12.36.**  $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$

**12.37.**  $\cot(A \pm B) = \frac{\cot A \cot B \mp 1}{\cot B \pm \cot A}$

**Functions of Angles in All Quadrants in Terms of Those in Quadrant I**

	$-A$	$90^\circ \pm A$ $\frac{\pi}{2} \pm A$	$180^\circ \pm A$ $\pi \pm A$	$270^\circ \pm A$ $\frac{3\pi}{2} \pm A$	$k(360^\circ) \pm A$ $2k\pi \pm A$ $k = \text{integer}$
sin	$-\sin A$	$\cos A$	$\sin A$	$-\cos A$	$\pm \sin A$
cos	$\cos A$	$\mp \sin A$	$-\cos A$	$\mp \sin A$	$\cos A$
tan	$-\tan A$	$\mp \cot A$	$\pm \tan A$	$\mp \cot A$	$\pm \tan A$
csc	$-\csc A$	$\sec A$	$\mp \csc A$	$-\sec A$	$\pm \csc A$
sec	$\sec A$	$\mp \csc A$	$-\sec A$	$\pm \csc A$	$\sec A$
cot	$-\cot A$	$\mp \tan A$	$\pm \cot A$	$\mp \tan A$	$\pm \cot A$

**Relationships Among Functions of Angles in Quadrant I**

	$\sin A = u$	$\cos A = u$	$\tan A = u$	$\cot A = u$	$\sec A = u$	$\csc A = u$
sin A	$u$	$\sqrt{1-u^2}$	$u/\sqrt{1+u^2}$	$1/\sqrt{1+u^2}$	$\sqrt{u^2-1}/u$	$1/u$
cos A	$\sqrt{1-u^2}$	$u$	$1/\sqrt{1+u^2}$	$u/\sqrt{1+u^2}$	$1/u$	$\sqrt{u^2-1}/u$
tan A	$u/\sqrt{1-u^2}$	$\sqrt{1-u^2}/u$	$u$	$1/u$	$\sqrt{u^2-1}$	$1/\sqrt{u^2-1}$
cot A	$\sqrt{1-u^2}/u$	$u/\sqrt{1-u^2}$	$1/u$	$u$	$1/\sqrt{u^2-1}$	$\sqrt{u^2-1}$
sec A	$1/\sqrt{1-u^2}$	$1/u$	$\sqrt{1+u^2}$	$\sqrt{1+u^2}/u$	$u$	$u/\sqrt{u^2-1}$
csc A	$1/u$	$1/\sqrt{1-u^2}$	$\sqrt{1+u^2}/u$	$\sqrt{1+u^2}$	$u/\sqrt{u^2-1}$	$u$

For extensions to other quadrants use appropriate signs as given in the preceding table.

### Double Angle Formulas

---

**12.38.**  $\sin 2A = 2 \sin A \cos A$

**12.39.**  $\cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = 2 \cos^2 A - 1$

**12.40.**  $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$

### Half Angle Formulas

---

**12.41.**  $\sin \frac{A}{2} = \pm \sqrt{\frac{1 - \cos A}{2}}$  [+ if  $A/2$  is in quadrant I or II  
- if  $A/2$  is in quadrant III or IV]

**12.42.**  $\cos \frac{A}{2} = \pm \sqrt{\frac{1 + \cos A}{2}}$  [+ if  $A/2$  is in quadrant I or IV  
- if  $A/2$  is in quadrant II or III]

**12.43.**  $\tan \frac{A}{2} = \pm \sqrt{\frac{1 - \cos A}{1 + \cos A}}$  [+ if  $A/2$  is in quadrant I or III  
- if  $A/2$  is in quadrant II or IV]

$$= \frac{\sin A}{1 + \cos A} = \frac{1 - \cos A}{\sin A} = \csc A - \cot A$$

### Multiple Angle Formulas

---

**12.44.**  $\sin 3A = 3 \sin A - 4 \sin^3 A$

**12.45.**  $\cos 3A = 4 \cos^3 A - 3 \cos A$

**12.46.**  $\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$

**12.47.**  $\sin 4A = 4 \sin A \cos A - 8 \sin^3 A \cos A$

**12.48.**  $\cos 4A = 8 \cos^4 A - 8 \cos^2 A + 1$

**12.49.**  $\tan 4A = \frac{4 \tan A - 4 \tan^3 A}{1 - 6 \tan^2 A + \tan^4 A}$

**12.50.**  $\sin 5A = 5 \sin A - 20 \sin^3 A + 16 \sin^5 A$

**12.51.**  $\cos 5A = 16 \cos^5 A - 20 \cos^3 A + 5 \cos A$

**12.52.**  $\tan 5A = \frac{\tan^5 A - 10 \tan^3 A + 5 \tan A}{1 - 10 \tan^2 A + 5 \tan^4 A}$

See also formulas 12.68 and 12.69.

### Powers of Trigonometric Functions

---

**12.53.**  $\sin^2 A = \frac{1}{2} - \frac{1}{2} \cos 2A$

**12.57.**  $\sin^4 A = \frac{3}{8} - \frac{1}{2} \cos 2A + \frac{1}{8} \cos 4A$

**12.54.**  $\cos^2 A = \frac{1}{2} + \frac{1}{2} \cos 2A$

**12.58.**  $\cos^4 A = \frac{3}{8} + \frac{1}{2} \cos 2A + \frac{1}{8} \cos 4A$

**12.55.**  $\sin^3 A = \frac{3}{4} \sin A - \frac{1}{4} \sin 3A$

**12.59.**  $\sin^5 A = \frac{5}{8} \sin A - \frac{5}{16} \sin 3A + \frac{1}{16} \sin 5A$

**12.56.**  $\cos^3 A = \frac{3}{4} \cos A + \frac{1}{4} \cos 3A$

**12.60.**  $\cos^5 A = \frac{5}{8} \cos A + \frac{5}{16} \cos 3A + \frac{1}{16} \cos 5A$

See also formulas 12.70 through 12.73.

### **Sum, Difference, and Product of Trigonometric Functions**

---

**12.61.**  $\sin A + \sin B = 2 \sin \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B)$

**12.62.**  $\sin A - \sin B = 2 \cos \frac{1}{2}(A+B) \sin \frac{1}{2}(A-B)$

**12.63.**  $\cos A + \cos B = 2 \cos \frac{1}{2}(A+B) \cos \frac{1}{2}(A-B)$

**12.64.**  $\cos A - \cos B = 2 \sin \frac{1}{2}(A+B) \sin \frac{1}{2}(B-A)$

**12.65.**  $\sin A \sin B = \frac{1}{2} \{ \cos(A-B) - \cos(A+B) \}$

**12.66.**  $\cos A \cos B = \frac{1}{2} \{ \cos(A-B) + \cos(A+B) \}$

**12.67.**  $\sin A \cos B = \frac{1}{2} \{ \sin(A-B) + \sin(A+B) \}$

### **General Formulas**

---

**12.68.**  $\sin nA = \sin A \left\{ (2 \cos A)^{n-1} - \binom{n-2}{1} (2 \cos A)^{n-3} + \binom{n-3}{2} (2 \cos A)^{n-5} - \dots \right\}$

**12.69.**  $\cos nA = \frac{1}{2} \left\{ (2 \cos A)^n - \frac{n}{1} (2 \cos A)^{n-2} + \frac{n}{2} \binom{n-3}{1} (2 \cos A)^{n-4} - \frac{n}{3} \binom{n-4}{2} (2 \cos A)^{n-6} + \dots \right\}$

**12.70.**  $\sin^{2n-1} A = \frac{(-1)^{n-1}}{2^{2n-2}} \left\{ \sin(2n-1)A - \binom{2n-1}{1} \sin(2n-3)A + \dots (-1)^{n-1} \binom{2n-1}{n-1} \sin A \right\}$

**12.71.**  $\cos^{2n-1} A = \frac{1}{2^{2n-2}} \left\{ \cos(2n-1)A + \binom{2n-1}{1} \cos(2n-3)A + \dots + \binom{2n-1}{n-1} \cos A \right\}$

**12.72.**  $\sin^{2n} A = \frac{1}{2^{2n}} \binom{2n}{n} + \frac{(-1)^n}{2^{2n-1}} \left\{ \cos 2nA - \binom{2n}{1} \cos(2n-2)A + \dots (-1)^{n-1} \binom{2n}{n-1} \cos 2A \right\}$

**12.73.**  $\cos^{2n} A = \frac{1}{2^{2n}} \binom{2n}{n} + \frac{1}{2^{2n-1}} \left\{ \cos 2nA + \binom{2n}{1} \cos(2n-2)A + \dots + \binom{2n}{n-1} \cos 2A \right\}$

### **Inverse Trigonometric Functions**

---

If  $x = \sin y$ , then  $y = \sin^{-1} x$ , i.e. the angle whose sine is  $x$  or inverse sine of  $x$  is a many-valued function of  $x$  which is a collection of single-valued functions called *branches*. Similarly, the other inverse trigonometric functions are multiple-valued.

For many purposes a particular branch is required. This is called the *principal branch* and the values for this branch are called *principal values*.

### Principal Values for Inverse Trigonometric Functions

Principal values for $x \geq 0$	Principal values for $x < 0$
$0 \leq \sin^{-1} x \leq \pi/2$	$-\pi/2 \leq \sin^{-1} x < 0$
$0 \leq \cos^{-1} x \leq \pi/2$	$\pi/2 < \cos^{-1} x \leq \pi$
$0 \leq \tan^{-1} x < \pi/2$	$-\pi/2 < \tan^{-1} x < 0$
$0 < \cot^{-1} x \leq \pi/2$	$\pi/2 < \cot^{-1} x < \pi$
$0 \leq \sec^{-1} x < \pi/2$	$\pi/2 < \sec^{-1} x \leq \pi$
$0 < \csc^{-1} x \leq \pi/2$	$-\pi/2 \leq \csc^{-1} x < 0$

### Relations Between Inverse Trigonometric Functions

In all cases it is assumed that principal values are used.

$$12.74. \quad \sin^{-1} x + \cos^{-1} x = \pi/2$$

$$12.80. \quad \sin^{-1}(-x) = -\sin^{-1} x$$

$$12.75. \quad \tan^{-1} x + \cot^{-1} x = \pi/2$$

$$12.81. \quad \cos^{-1}(-x) = \pi - \cos^{-1} x$$

$$12.76. \quad \sec^{-1} x + \csc^{-1} x = \pi/2$$

$$12.82. \quad \tan^{-1}(-x) = -\tan^{-1} x$$

$$12.77. \quad \csc^{-1} x = \sin^{-1}(1/x)$$

$$12.83. \quad \cot^{-1}(-x) = \pi - \cot^{-1} x$$

$$12.78. \quad \sec^{-1} x = \cos^{-1}(1/x)$$

$$12.84. \quad \sec^{-1}(-x) = \pi - \sec^{-1} x$$

$$12.79. \quad \cot^{-1} x = \tan^{-1}(1/x)$$

$$12.85. \quad \csc^{-1}(-x) = -\csc^{-1} x$$

### Graphs of Inverse Trigonometric Functions

In each graph  $y$  is in radians. Solid portions of curves correspond to principal values.

$$12.86. \quad y = \sin^{-1} x$$




Fig. 12-11

$$12.87. \quad y = \cos^{-1} x$$




Fig. 12-12

$$12.88. \quad y = \tan^{-1} x$$




Fig. 12-13

12.89.  $y = \cot^{-1} x$




Fig. 12-14

12.90.  $y = \sec^{-1} x$




Fig. 12-15

12.91.  $y = \csc^{-1} x$




Fig. 12-16

### Relationships Between Sides and Angles of a Plane Triangle

The following results hold for any plane triangle  $ABC$  with sides  $a, b, c$  and angles  $A, B, C$ .

#### 12.92. Law of Sines:

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

#### 12.93. Law of Cosines:

$$c^2 = a^2 + b^2 - 2ab \cos C$$

with similar relations involving the other sides and angles.

#### 12.94. Law of Tangents:

$$\frac{a+b}{a-b} = \frac{\tan \frac{1}{2}(A+B)}{\tan \frac{1}{2}(A-B)}$$

with similar relations involving the other sides and angles.




Fig. 12-17

12.95.  $\sin A = \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)}$

where  $s = \frac{1}{2}(a+b+c)$  is the semiperimeter of the triangle. Similar relations involving angles  $B$  and  $C$  can be obtained.

See also formula 7.5.

### Relationships Between Sides and Angles of a Spherical Triangle

Spherical triangle  $ABC$  is on the surface of a sphere as shown in Fig. 12-18. Sides  $a, b, c$  (which are arcs of great circles) are measured by their angles subtended at center  $O$  of the sphere.  $A, B, C$  are the angles opposite sides  $a, b, c$ , respectively. Then the following results hold.

#### 12.96. Law of Sines:

$$\frac{\sin a}{\sin A} = \frac{\sin b}{\sin B} = \frac{\sin c}{\sin C}$$

#### 12.97. Law of Cosines:

$$\cos a = \cos b \cos c + \sin b \sin c \cos A$$

$$\cos A = -\cos B \cos C + \sin B \sin C \cos a$$

with similar results involving other sides and angles.




Fig. 12-18

**12.98. Law of Tangents:**

$$\frac{\tan \frac{1}{2}(A+B)}{\tan \frac{1}{2}(A-B)} = \frac{\tan \frac{1}{2}(a+b)}{\tan \frac{1}{2}(a-b)}$$

with similar results involving other sides and angles.

$$12.99. \cos \frac{A}{2} = \sqrt{\frac{\sin s \sin(s-c)}{\sin b \sin c}}$$

where  $s = \frac{1}{2}(a+b+c)$ . Similar results hold for other sides and angles.

$$12.100. \cos \frac{a}{2} = \sqrt{\frac{\cos(S-B)\cos(S-C)}{\sin B \sin C}}$$

where  $S = \frac{1}{2}(A+B+C)$ . Similar results hold for other sides and angles.

See also formula 7.44.

**Napier's Rules for Right Angled Spherical Triangles**

Except for right angle  $C$ , there are five parts of spherical triangle  $ABC$  which, if arranged in the order as given in Fig. 12-19, would be  $a, b, A, c, B$ .




Fig. 12-19




Fig. 12-20

Suppose these quantities are arranged in a circle as in Fig. 12-20 where we attach the prefix "co" (indicating complement) to hypotenuse  $c$  and angles  $A$  and  $B$ .

Any one of the parts of this circle is called a *middle part*, the two neighboring parts are called *adjacent parts*, and the two remaining parts are called *opposite parts*. Then Napier's rules are

**12.101.** The sine of any middle part equals the product of the tangents of the adjacent parts.

**12.102.** The sine of any middle part equals the product of the cosines of the opposite parts.

**EXAMPLE:** Since  $\text{co}-A = 90^\circ - A$ ,  $\text{co}-B = 90^\circ - B$ , we have

$$\begin{array}{lll} \sin a = \tan b (\text{co}-B) & \text{or} & \sin a = \tan b \cot B \\ \sin (\text{co}-A) = \cos a \cos (\text{co}-B) & \text{or} & \cos A = \cos a \sin B \end{array}$$

These can of course be obtained also from the results of 12.97.

# 13 EXPONENTIAL and LOGARITHMIC FUNCTIONS

## Laws of Exponents

In the following  $p, q$  are real numbers,  $a, b$  are positive numbers, and  $m, n$  are positive integers.

$$13.1. \quad a^p \cdot a^q = a^{p+q}$$

$$13.2. \quad a^p/a^q = a^{p-q}$$

$$13.3. \quad (a^p)^q = a^{pq}$$

$$13.4. \quad a^0 = 1, \quad a \neq 0$$

$$13.5. \quad a^{-p} = 1/a^p$$

$$13.6. \quad (ab)^p = a^p b^p$$

$$13.7. \quad \sqrt[n]{a} = a^{1/n}$$

$$13.8. \quad \sqrt[n]{a^m} = a^{m/n}$$

$$13.9. \quad \sqrt[n]{a/b} = \sqrt[n]{a}/\sqrt[n]{b}$$

In  $a^p$ ,  $p$  is called the *exponent*,  $a$  is the *base*, and  $a^p$  is called the *pth power of a*. The function  $y = a^x$  is called an *exponential function*.

## Logarithms and Antilogarithms

If  $a^p = N$  where  $a \neq 0$  or 1, then  $p = \log_a N$  is called the *logarithm* of  $N$  to the base  $a$ . The number  $N = a^p$  is called the *antilogarithm* of  $p$  to the base  $a$ , written  $\text{antilog}_a p$ .

**Example:** Since  $3^2 = 9$  we have  $\log_3 9 = 2$ .  $\text{antilog}_3 2 = 9$ .

The function  $y = \log_a x$  is called a *logarithmic function*.

## Laws of Logarithms

$$13.10. \quad \log_a MN = \log_a M + \log_a N$$

$$13.11. \quad \log_a \frac{M}{N} = \log_a M - \log_a N$$

$$13.12. \quad \log_a M^p = p \log_a M$$

## Common Logarithms and Antilogarithms

Common logarithms and antilogarithms (also called *Briggsian*) are those in which the base  $a = 10$ . The common logarithm of  $N$  is denoted by  $\log_{10} N$  or briefly  $\log N$ . For numerical values of common logarithms, see Table 1.

## Natural Logarithms and Antilogarithms

Natural logarithms and antilogarithms (also called *Napierian*) are those in which the base  $a = e = 2.71828 \dots$  [see page 3]. The natural logarithm of  $N$  is denoted by  $\log_e N$  or  $\ln N$ . For numerical values of natural logarithms see Table 7. For values of natural antilogarithms (i.e., a table giving  $e^x$  for values of  $x$ ) see Table 8.

### Change of Base of Logarithms

---

The relationship between logarithms of a number  $N$  to different bases  $a$  and  $b$  is given by

$$13.13. \quad \log_a N = \frac{\log_b N}{\log_b a}$$

In particular,

$$13.14. \quad \log_e N = \ln N = 2.30258 \ 50929 \ 94 \dots \log_{10} N$$

$$13.15. \quad \log_{10} N = \log N = 0.43429 \ 44819 \ 03 \dots \log_e N$$

### Relationship Between Exponential and Trigonometric Functions

---

$$13.16. \quad e^{i\theta} = \cos \theta + i \sin \theta, \quad e^{-i\theta} = \cos \theta - i \sin \theta$$

These are called *Euler's identities*. Here  $i$  is the imaginary unit [see page 10].

$$13.17. \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$13.18. \quad \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}$$

$$13.19. \quad \tan \theta = \frac{e^{i\theta} - e^{-i\theta}}{i(e^{i\theta} + e^{-i\theta})} = -i \left( \frac{e^{i\theta} - e^{-i\theta}}{e^{i\theta} + e^{-i\theta}} \right)$$

$$13.20. \quad \cot \theta = i \left( \frac{e^{i\theta} + e^{-i\theta}}{e^{i\theta} - e^{-i\theta}} \right)$$

$$13.21. \quad \sec \theta = \frac{2}{e^{i\theta} + e^{-i\theta}}$$

$$13.22. \quad \csc \theta = \frac{2i}{e^{i\theta} - e^{-i\theta}}$$

### Periodicity of Exponential Functions

---

$$13.23. \quad e^{i(\theta + 2k\pi)} = e^{i\theta} \quad k = \text{integer}$$

From this it is seen that  $e^x$  has period  $2\pi i$ .

### Polar Form of Complex Numbers Expressed as an Exponential

---

The polar form (see 4.7) of a complex number  $z = x + iy$  can be written in terms of exponentials as follows:

$$13.24. \quad z = x + iy = r(\cos \theta + i \sin \theta) = re^{i\theta}$$

**Operations with Complex Numbers in Polar Form**

---

Formulas 4.8 to 4.11 are equivalent to the following:

$$13.25. \quad (r_1 e^{i\theta_1})(r_2 e^{i\theta_2}) = r_1 r_2 e^{i(\theta_1 + \theta_2)}$$

$$13.26. \quad \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}$$

$$13.27. \quad (re^{i\theta})^p = r^p e^{ip\theta} \quad (\text{De Moivre's theorem})$$

$$13.28. \quad (re^{i\theta})^{1/n} = [re^{i(\theta+2k\pi)}]^{1/n} = r^{1/n} e^{i(\theta+2k\pi)/n}$$

**Logarithm of a Complex Number**

---

$$13.29. \quad \ln(re^{i\theta}) = \ln r + i\theta + 2k\pi i \quad k = \text{integer}$$

# 14 HYPERBOLIC FUNCTIONS

## Definition of Hyperbolic Functions

---

$$14.1. \text{ Hyperbolic sine of } x = \sinh x = \frac{e^x - e^{-x}}{2}$$

$$14.2. \text{ Hyperbolic cosine of } x = \cosh x = \frac{e^x + e^{-x}}{2}$$

$$14.3. \text{ Hyperbolic tangent of } x = \tanh x = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

$$14.4. \text{ Hyperbolic cotangent of } x = \coth x = \frac{e^x + e^{-x}}{e^x - e^{-x}}$$

$$14.5. \text{ Hyperbolic secant of } x = \operatorname{sech} x = \frac{2}{e^x + e^{-x}}$$

$$14.6. \text{ Hyperbolic cosecant of } x = \operatorname{csch} x = \frac{2}{e^x - e^{-x}}$$

## Relationships Among Hyperbolic Functions

---

$$14.7. \tanh x = \frac{\sinh x}{\cosh x}$$

$$14.8. \coth x = \frac{1}{\tanh x} = \frac{\cosh x}{\sinh x}$$

$$14.9. \operatorname{sech} x = \frac{1}{\cosh x}$$

$$14.10. \operatorname{csch} x = \frac{1}{\sinh x}$$

$$14.11. \cosh^2 x - \sinh^2 x = 1$$

$$14.12. \operatorname{sech}^2 x + \tanh^2 x = 1$$

$$14.13. \coth^2 x - \operatorname{csch}^2 x = 1$$

## Functions of Negative Arguments

---

$$14.14. \sinh(-x) = -\sinh x$$

$$14.15. \cosh(-x) = \cosh x$$

$$14.16. \tanh(-x) = -\tanh x$$

$$14.17. \operatorname{csch}(-x) = -\operatorname{csch} x$$

$$14.18. \operatorname{sech}(-x) = \operatorname{sech} x$$

$$14.19. \coth(-x) = -\coth x$$

**Addition Formulas**

---

$$14.20. \sinh(x \pm y) = \sinh x \cosh y \pm \cosh x \sinh y$$

$$14.21. \cosh(x \pm y) = \cosh x \cosh y \pm \sinh x \sinh y$$

$$14.22. \tanh(x \pm y) = \frac{\tanh x \pm \tanh y}{1 \pm \tanh x \tanh y}$$

$$14.23. \coth(x \pm y) = \frac{\coth x \coth y \pm 1}{\coth y \pm \coth x}$$

**Double Angle Formulas**

---

$$14.24. \sinh 2x = 2 \sinh x \cosh x$$

$$14.25. \cosh 2x = \cosh^2 x + \sinh^2 x = 2 \cosh^2 x - 1 = 1 + 2 \sinh^2 x$$

$$14.26. \tanh 2x = \frac{2 \tanh x}{1 + \tanh^2 x}$$

**Half Angle Formulas**

---

$$14.27. \sinh \frac{x}{2} = \pm \sqrt{\frac{\cosh x - 1}{2}} \quad [+ \text{ if } x > 0, - \text{ if } x < 0]$$

$$14.28. \cosh \frac{x}{2} = \sqrt{\frac{\cosh x + 1}{2}}$$

$$14.29. \tanh \frac{x}{2} = \pm \sqrt{\frac{\cosh x - 1}{\cosh x + 1}} \quad [+ \text{ if } x > 0, - \text{ if } x < 0]$$

$$= \frac{\sinh x}{\cosh x + 1} = \frac{\cosh x - 1}{\sinh x}$$

**Multiple Angle Formulas**

---

$$14.30. \sinh 3x = 3 \sinh x + 4 \sinh^3 x$$

$$14.31. \cosh 3x = 4 \cosh^3 x - 3 \cosh x$$

$$14.32. \tanh 3x = \frac{3 \tanh x + \tanh^3 x}{1 + 3 \tanh^2 x}$$

$$14.33. \sinh 4x = 8 \sinh^3 x \cosh x + 4 \sinh x \cosh x$$

$$14.34. \cosh 4x = 8 \cosh^4 x - 8 \cosh^2 x + 1$$

$$14.35. \tanh 4x = \frac{4 \tanh x + 4 \tanh^3 x}{1 + 6 \tanh^2 x + \tanh^4 x}$$

### Powers of Hyperbolic Functions

---

**14.36.**  $\sinh^2 x = \frac{1}{2} \cosh 2x - \frac{1}{2}$

**14.37.**  $\cosh^2 x = \frac{1}{2} \cosh 2x + \frac{1}{2}$

**14.38.**  $\sinh^3 x = \frac{1}{4} \sinh 3x - \frac{3}{4} \sinh x$

**14.39.**  $\cosh^3 x = \frac{1}{4} \cosh 3x + \frac{3}{4} \cosh x$

**14.40.**  $\sinh^4 x = \frac{3}{8} - \frac{1}{2} \cosh 2x + \frac{1}{8} \cosh 4x$

**14.41.**  $\cosh^4 x = \frac{3}{8} + \frac{1}{2} \cosh 2x + \frac{1}{8} \cosh 4x$

### Sum, Difference, and Product of Hyperbolic Functions

---

**14.42.**  $\sinh x + \sinh y = 2 \sinh \frac{1}{2}(x+y) \cosh \frac{1}{2}(x-y)$

**14.43.**  $\sinh x - \sinh y = 2 \cosh \frac{1}{2}(x+y) \sinh \frac{1}{2}(x-y)$

**14.44.**  $\cosh x + \cosh y = 2 \cosh \frac{1}{2}(x+y) \cosh \frac{1}{2}(x-y)$

**14.45.**  $\cosh x - \cosh y = 2 \sinh \frac{1}{2}(x+y) \sinh \frac{1}{2}(x-y)$

**14.46.**  $\sinh x \sinh y = \frac{1}{2} \{\cosh(x+y) - \cosh(x-y)\}$

**14.47.**  $\cosh x \cosh y = \frac{1}{2} \{\cosh(x+y) + \cosh(x-y)\}$

**14.48.**  $\sinh x \cosh y = \frac{1}{2} \{\sinh(x+y) + \sinh(x-y)\}$

### Expression of Hyperbolic Functions in Terms of Others

---

In the following we assume  $x > 0$ . If  $x < 0$ , use the appropriate sign as indicated by formulas 14.14 to 14.19.

	$\sinh x = u$	$\cosh x = u$	$\tanh x = u$	$\coth x = u$	$\operatorname{sech} x = u$	$\operatorname{csch} x = u$
$\sinh x$	$u$	$\sqrt{u^2 - 1}$	$u/\sqrt{1-u^2}$	$1/\sqrt{u^2 - 1}$	$\sqrt{1-u^2}/u$	$1/u$
$\cosh x$	$\sqrt{1+u^2}$	$u$	$1/\sqrt{1-u^2}$	$u/\sqrt{u^2 - 1}$	$1/u$	$\sqrt{1+u^2}/u$
$\tanh x$	$u/\sqrt{1+u^2}$	$\sqrt{u^2 - 1}/u$	$u$	$1/u$	$\sqrt{1-u^2}$	$1/\sqrt{1+u^2}$
$\coth x$	$\sqrt{u^2 + 1}/u$	$u/\sqrt{u^2 - 1}$	$1/u$	$u$	$1/\sqrt{1-u^2}$	$\sqrt{1+u^2}$
$\operatorname{sech} x$	$1/\sqrt{1+u^2}$	$1/u$	$\sqrt{1-u^2}$	$\sqrt{u^2 - 1}/u$	$u$	$u/\sqrt{1+u^2}$
$\operatorname{csch} x$	$1/u$	$1/\sqrt{u^2 - 1}$	$\sqrt{1-u^2}/u$	$\sqrt{u^2 - 1}$	$u/\sqrt{1-u^2}$	$u$

**Graphs of Hyperbolic Functions**

14.49.  $y = \sinh x$




Fig. 14-1

14.50.  $y = \cosh x$




Fig. 14-2

14.51.  $y = \tanh x$




Fig. 14-3

14.52.  $y = \coth x$




Fig. 14-4

14.53.  $y = \operatorname{sech} x$




Fig. 14-5

14.54.  $y = \operatorname{csch} x$




Fig. 14-6

**Inverse Hyperbolic Functions**

If  $x = \sinh y$ , then  $y = \sinh^{-1} x$  is called the *inverse hyperbolic sine* of  $x$ . Similarly we define the other inverse hyperbolic functions. The inverse hyperbolic functions are multiple-valued and as in the case of inverse trigonometric functions [see page 49] we restrict ourselves to principal values for which they can be considered as single-valued.

The following list shows the principal values (unless otherwise indicated) of the inverse hyperbolic functions expressed in terms of logarithmic functions which are taken as real valued.

14.55.  $\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$        $-\infty < x < \infty$

14.56.  $\cosh^{-1} x = \ln(x + \sqrt{x^2 - 1})$        $x \geq 1$     ( $\cosh^{-1} x > 0$  is principal value)

14.57.  $\tanh^{-1} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$        $-1 < x < 1$

14.58.  $\coth^{-1} x = \frac{1}{2} \ln\left(\frac{x+1}{x-1}\right)$        $x > 1 \text{ or } x < -1$

14.59.  $\operatorname{sech}^{-1} x = \ln\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} - 1}\right)$        $0 < x \leq 1$     ( $\operatorname{sech}^{-1} x > 0$  is principal value)

14.60.  $\operatorname{csch}^{-1} x = \ln\left(\frac{1}{x} + \sqrt{\frac{1}{x^2} + 1}\right)$        $x \neq 0$

### Relations Between Inverse Hyperbolic Functions

**14.61.**  $\operatorname{csch}^{-1} x = \sinh^{-1}(1/x)$

**14.62.**  $\operatorname{sech}^{-1} x = \cosh^{-1}(1/x)$

**14.63.**  $\operatorname{coth}^{-1} x = \tanh^{-1}(1/x)$

**14.64.**  $\sinh^{-1}(-x) = -\sinh^{-1} x$

**14.65.**  $\tanh^{-1}(-x) = -\tanh^{-1} x$

**14.66.**  $\coth^{-1}(-x) = -\coth^{-1} x$

**14.67.**  $\operatorname{csch}^{-1}(-x) = -\operatorname{csch}^{-1} x$

### Graphs of Inverse Hyperbolic Functions

**14.68.**  $y = \sinh^{-1} x$




Fig. 14-7

**14.69.**  $y = \cosh^{-1} x$




Fig. 14-8

**14.70.**  $y = \tanh^{-1} x$




Fig. 14-9

**14.71.**  $y = \coth^{-1} x$




Fig. 14-10

**14.72.**  $y = \operatorname{sech}^{-1} x$




Fig. 14-11

**14.73.**  $y = \operatorname{csch}^{-1} x$




Fig. 14-12

**Relationship Between Hyperbolic and Trigonometric Functions**

14.74.  $\sin(ix) = i \sinh x$

14.75.  $\cos(ix) = \cosh x$

14.76.  $\tan(ix) = i \tanh x$

14.77.  $\csc(ix) = -i \operatorname{csch} x$

14.78.  $\sec(ix) = \operatorname{sech} x$

14.79.  $\cot(ix) = -i \operatorname{coth} x$

14.80.  $\sinh(ix) = i \sin x$

14.81.  $\cosh(ix) = \cos x$

14.82.  $\tanh(ix) = i \tan x$

14.83.  $\operatorname{csch}(ix) = -i \csc x$

14.84.  $\operatorname{sech}(ix) = \sec x$

14.85.  $\operatorname{coth}(ix) = -i \cot x$

**Periodicity of Hyperbolic Functions**

In the following  $k$  is any integer.

14.86.  $\sinh(x + 2k\pi i) = \sinh x$

14.87.  $\cosh(x + 2k\pi i) = \cosh x$

14.88.  $\tanh(x + k\pi i) = \tanh x$

14.89.  $\operatorname{csch}(x + 2k\pi i) = \operatorname{csch} x$

14.90.  $\operatorname{sech}(x + 2k\pi i) = \operatorname{sech} x$

14.91.  $\operatorname{coth}(x + k\pi i) = \operatorname{coth} x$

**Relationship Between Inverse Hyperbolic and Inverse Trigonometric Functions**

14.92.  $\sin^{-1}(ix) = i \sin^{-1} x$

14.93.  $\sinh^{-1}(ix) = i \sin^{-1} x$

14.94.  $\cos^{-1} x = \pm i \cosh^{-1} x$

14.95.  $\cosh^{-1} x = \pm i \cos^{-1} x$

14.96.  $\tan^{-1}(ix) = i \tanh^{-1} x$

14.97.  $\tanh^{-1}(ix) = i \tan^{-1} x$

14.98.  $\cot^{-1}(ix) = i \coth^{-1} x$

14.99.  $\coth^{-1}(ix) = -i \cot^{-1} x$

14.100.  $\sec^{-1} x = \pm i \operatorname{sech}^{-1} x$

14.101.  $\operatorname{sech}^{-1} x = \pm i \sec^{-1} x$

14.102.  $\csc^{-1}(ix) = -i \operatorname{csch}^{-1} x$

14.103.  $\operatorname{csch}^{-1}(ix) = -i \csc^{-1} x$

## Section IV: Calculus

# 15 DERIVATIVES

### Definition of a Derivative

---

Suppose  $y = f(x)$ . The derivative of  $y$  or  $f(x)$  is defined as

$$15.1. \frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}$$

where  $h = \Delta x$ . The derivative is also denoted by  $y'$ ,  $df/dx$  or  $f'(x)$ . The process of taking a derivative is called *differentiation*.

### General Rules of Differentiation

---

In the following,  $u, v, w$  are functions of  $x$ ;  $a, b, c, n$  are constants (restricted if indicated);  $e = 2.71828 \dots$  is the natural base of logarithms;  $\ln u$  is the natural logarithm of  $u$  (i.e., the logarithm to the base  $e$ ) where it is assumed that  $u > 0$  and all angles are in radians.

$$15.2. \frac{d}{dx}(c) = 0$$

$$15.3. \frac{d}{dx}(cx) = c$$

$$15.4. \frac{d}{dx}(cx^n) = ncx^{n-1}$$

$$15.5. \frac{d}{dx}(u \pm v \pm w \pm \dots) = \frac{du}{dx} \pm \frac{dv}{dx} \pm \frac{dw}{dx} \pm \dots$$

$$15.6. \frac{d}{dx}(cu) = c \frac{du}{dx}$$

$$15.7. \frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx}$$

$$15.8. \frac{d}{dx}(uvw) = uv \frac{dw}{dx} + uw \frac{dv}{dx} + vw \frac{du}{dx}$$

$$15.9. \frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v(du/dx) - u(dv/dx)}{v^2}$$

$$15.10. \frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$15.11. \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad (\text{Chain rule})$$

$$15.12. \frac{du}{dx} = \frac{1}{dx/du}$$

$$15.13. \frac{dy}{dx} = \frac{dy/du}{dx/du}$$

### Derivatives of Trigonometric and Inverse Trigonometric Functions

---

$$15.14. \frac{d}{dx} \sin u = \cos u \frac{du}{dx}$$

$$15.15. \frac{d}{dx} \cos u = -\sin u \frac{du}{dx}$$

$$15.16. \frac{d}{dx} \tan u = \sec^2 u \frac{du}{dx}$$

$$15.17. \frac{d}{dx} \cot u = -\csc^2 u \frac{du}{dx}$$

$$15.18. \frac{d}{dx} \sec u = \sec u \tan u \frac{du}{dx}$$

$$15.19. \frac{d}{dx} \csc u = -\csc u \cot u \frac{du}{dx}$$

$$15.20. \frac{d}{dx} \sin^{-1} u = \frac{1}{\sqrt{1-u^2}} \frac{du}{dx} \quad \left[ -\frac{\pi}{2} < \sin^{-1} u < \frac{\pi}{2} \right]$$

$$15.21. \frac{d}{dx} \cos^{-1} u = \frac{-1}{\sqrt{1-u^2}} \frac{du}{dx} \quad [0 < \cos^{-1} u < \pi]$$

$$15.22. \frac{d}{dx} \tan^{-1} u = \frac{1}{1+u^2} \frac{du}{dx} \quad \left[ -\frac{\pi}{2} < \tan^{-1} u < \frac{\pi}{2} \right]$$

$$15.23. \frac{d}{dx} \cot^{-1} u = \frac{-1}{1+u^2} \frac{du}{dx} \quad [0 < \cot^{-1} u < \pi]$$

$$15.24. \frac{d}{dx} \sec^{-1} u = \frac{1}{|u|\sqrt{u^2-1}} \frac{du}{dx} = \frac{\pm 1}{u\sqrt{u^2-1}} \frac{du}{dx} \quad \left[ \begin{array}{l} + \text{ if } 0 < \sec^{-1} u < \pi/2 \\ - \text{ if } \pi/2 < \sec^{-1} u < \pi \end{array} \right]$$

$$15.25. \frac{d}{dx} \csc^{-1} u = \frac{-1}{|u|\sqrt{u^2-1}} \frac{du}{dx} = \frac{\mp 1}{u\sqrt{u^2-1}} \frac{du}{dx} \quad \left[ \begin{array}{l} - \text{ if } 0 < \csc^{-1} u < \pi/2 \\ + \text{ if } -\pi/2 < \csc^{-1} u < 0 \end{array} \right]$$

### Derivatives of Exponential and Logarithmic Functions

---

$$15.26. \frac{d}{dx} \log_a u = \frac{\log_a e}{u} \frac{du}{dx} \quad a \neq 0, 1$$

$$15.27. \frac{d}{dx} \ln u = \frac{d}{dx} \log_e u = \frac{1}{u} \frac{du}{dx}$$

$$15.28. \frac{d}{dx} a^u = a^u \ln a \frac{du}{dx}$$

15.29.  $\frac{d}{dx} e^u = e^u \frac{du}{dx}$

15.30.  $\frac{d}{dx} u^v = \frac{d}{dx} e^{v \ln u} = e^{v \ln u} \frac{d}{dx} [v \ln u] = vu^{v-1} \frac{du}{dx} + u^v \ln u \frac{dv}{dx}$

### Derivatives of Hyperbolic and Inverse Hyperbolic Functions

---

15.31.  $\frac{d}{dx} \sinh u = \cosh u \frac{du}{dx}$

15.32.  $\frac{d}{dx} \cosh u = \sinh u \frac{du}{dx}$

15.33.  $\frac{d}{dx} \tanh u = \operatorname{sech}^2 u \frac{du}{dx}$

15.34.  $\frac{d}{dx} \coth u = -\operatorname{csch}^2 u \frac{du}{dx}$

15.35.  $\frac{d}{dx} \operatorname{sech} u = -\operatorname{sech} u \tanh u \frac{du}{dx}$

15.36.  $\frac{d}{dx} \operatorname{csch} u = -\operatorname{csch} u \coth u \frac{du}{dx}$

15.37.  $\frac{d}{dx} \sinh^{-1} u = \frac{1}{\sqrt{u^2 + 1}} \frac{du}{dx}$

15.38.  $\frac{d}{dx} \cosh^{-1} u = \frac{\pm 1}{\sqrt{u^2 - 1}} \frac{du}{dx}$

$[+ \text{ if } \cosh^{-1} u > 0, u > 1]$   
 $[- \text{ if } \cosh^{-1} u < 0, u > 1]$

15.39.  $\frac{d}{dx} \tanh^{-1} u = \frac{1}{1-u^2} \frac{du}{dx}$

$[-1 < u < 1]$

15.40.  $\frac{d}{dx} \coth^{-1} u = \frac{1}{1-u^2} \frac{du}{dx}$

$[u > 1 \text{ or } u < -1]$

15.41.  $\frac{d}{dx} \operatorname{sech}^{-1} u = \frac{\mp 1}{u \sqrt{1-u^2}} \frac{du}{dx}$

$[- \text{ if } \operatorname{sech}^{-1} u > 0, 0 < u < 1]$   
 $[+ \text{ if } \operatorname{sech}^{-1} u < 0, 0 < u < 1]$

15.42.  $\frac{d}{dx} \operatorname{csch}^{-1} u = \frac{-1}{|u| \sqrt{1+u^2}} \frac{du}{dx} = \frac{\mp 1}{u \sqrt{1+u^2}} \frac{du}{dx}$

$[- \text{ if } u > 0, + \text{ if } u < 0]$

### Higher Derivatives

---

The second, third, and higher derivatives are defined as follows.

15.43. Second derivative  $= \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d^2 y}{dx^2} = f''(x) = y''$

15.44. Third derivative  $= \frac{d}{dx} \left( \frac{d^2 y}{dx^2} \right) = \frac{d^3 y}{dx^3} = f'''(x) = y'''$

15.45.  $n$ th derivative  $= \frac{d}{dx} \left( \frac{d^{n-1} y}{dx^{n-1}} \right) = \frac{d^n y}{dx^n} = f^{(n)}(x) = y^{(n)}$

### Leibniz's Rule for Higher Derivatives of Products

---

Let  $D^p$  stand for the operator  $\frac{d^p}{dx^p}$  so that  $D^p u = \frac{d^p u}{dx^p}$  = the  $p$ th derivative of  $u$ . Then

$$15.46. \quad D^n(uv) = uD^n v + \binom{n}{1}(Du)(D^{n-1}v) + \binom{n}{2}(D^2u)(D^{n-2}v) + \cdots + vD^n u$$

where  $\binom{n}{1}, \binom{n}{2}, \dots$  are the binomial coefficients (see 3.5).

As special cases we have

$$15.47. \quad \frac{d^2}{dx^2}(uv) = u \frac{d^2v}{dx^2} + 2 \frac{du}{dx} \frac{dv}{dx} + v \frac{d^2u}{dx^2}$$

$$15.48. \quad \frac{d^3}{dx^3}(uv) = u \frac{d^3v}{dx^3} + 3 \frac{du}{dx} \frac{d^2v}{dx^2} + 3 \frac{d^2u}{dx^2} \frac{dv}{dx} + v \frac{d^3u}{dx^3}$$

### Differentials

---

Let  $y = f(x)$  and  $\Delta y = f(x + \Delta x) - f(x)$ . Then

$$15.49. \quad \frac{\Delta y}{\Delta x} = \frac{f(x + \Delta x) - f(x)}{\Delta x} = f'(x) + \epsilon = \frac{dy}{dx} + \epsilon$$

where  $\epsilon \rightarrow 0$  as  $\Delta x \rightarrow 0$ . Thus,

$$15.50. \quad \Delta y = f'(x)\Delta x + \epsilon\Delta x$$

If we call  $\Delta x = dx$  the differential of  $x$ , then we define the differential of  $y$  to be

$$15.51. \quad dy = f'(x)dx$$

### Rules for Differentials

---

The rules for differentials are exactly analogous to those for derivatives. As examples we observe that

$$15.52. \quad d(u \pm v \pm w \pm \cdots) = du \pm dv \pm dw \pm \cdots$$

$$15.53. \quad d(uv) = u dv + v du$$

$$15.54. \quad d\left(\frac{u}{v}\right) = \frac{v du - u dv}{v^2}$$

$$15.55. \quad d(u^n) = nu^{n-1} du$$

$$15.56. \quad d(\sin u) = \cos u du$$

$$15.57. \quad d(\cos u) = -\sin u du$$

## Partial Derivatives

---

Let  $z = f(x, y)$  be a function of the two variables  $x$  and  $y$ . Then we define the *partial derivative* of  $z$  or  $f(x, y)$  with respect to  $x$ , keeping  $y$  constant, to be

$$15.58. \quad \frac{\partial f}{\partial x} = \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x, y) - f(x, y)}{\Delta x}$$

This partial derivative is also denoted by  $\partial z / \partial x$ ,  $f_x$ , or  $z_x$ .

Similarly the partial derivative of  $z = f(x, y)$  with respect to  $y$ , keeping  $x$  constant, is defined to be

$$15.59. \quad \frac{\partial f}{\partial y} = \lim_{\Delta y \rightarrow 0} \frac{f(x, y + \Delta y) - f(x, y)}{\Delta y}$$

This partial derivative is also denoted by  $\partial z / \partial y$ ,  $f_y$ , or  $z_y$ .

Partial derivatives of higher order can be defined as follows:

$$15.60. \quad \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right), \quad \frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial y} \right)$$

$$15.61. \quad \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right), \quad \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right)$$

The results in 15.61 will be equal if the function and its partial derivatives are continuous; that is, in such cases, the order of differentiation makes no difference.

Extensions to functions of more than two variables are exactly analogous.

## Multivariable Differentials

---

The differential of  $z = f(x, y)$  is defined as

$$15.62. \quad dz = df = \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

where  $dx = \Delta x$  and  $dy = \Delta y$ . Note that  $dz$  is a function of four variables, namely  $x, y, dx, dy$ , and is linear in the variables  $dx$  and  $dy$ .

Extensions to functions of more than two variables are exactly analogous.

**EXAMPLE:** Let  $z = x^2 + 5xy + 2y^3$ . Then

$$z_x = 2x + 5y \quad \text{and} \quad z_y = 5x + 6y^2$$

and hence

$$dz = (2x + 5y) dx + (5x + 6y^2) dy$$

Suppose we want to find  $dz$  for  $dx = 2$ ,  $dy = 3$  and at the point  $P(4, 1)$ , i.e., when  $x = 4$  and  $y = 1$ . Substitution yields

$$dz = (8 + 5)2 + (20 + 6)3 = 26 + 78 = 104$$

# 16 INDEFINITE INTEGRALS

## Definition of an Indefinite Integral

If  $\frac{dy}{dx} = f(x)$ , then  $y$  is the function whose derivative is  $f(x)$  and is called the *anti-derivative* of  $f(x)$  or the *indefinite integral* of  $f(x)$ , denoted by  $\int f(x) dx$ . Similarly if  $y = \int f(u) du$ , then  $\frac{dy}{du} = f(u)$ . Since the derivative of a constant is zero, all indefinite integrals differ by an arbitrary constant.

For the definition of a definite integral, see 18.1. The process of finding an integral is called *integration*.

## General Rules of Integration

In the following,  $u, v, w$  are functions of  $x$ ;  $a, b, p, q, n$  any constants, restricted if indicated;  $e = 2.71828 \dots$  is the natural base of logarithms;  $\ln u$  denotes the natural logarithm of  $u$  where it is assumed that  $u > 0$  (in general, to extend formulas to cases where  $u < 0$  as well, replace  $\ln u$  by  $\ln |u|$ ); all angles are in radians; all constants of integration are omitted but implied.

$$16.1. \quad \int a \, dx = ax$$

$$16.2. \quad \int af(x) \, dx = a \int f(x) \, dx$$

$$16.3. \quad \int (u \pm v \pm w \pm \dots) \, dx = \int u \, dx \pm \int v \, dx \pm \int w \, dx \pm \dots$$

$$16.4. \quad \int u \, dv = uv - \int v \, du \quad (\text{Integration by parts})$$

For generalized integration by parts, see 16.48.

$$16.5. \quad \int f(ax) \, dx = \frac{1}{a} \int f(u) \, du$$

$$16.6. \quad \int F\{f(x)\} \, dx = \int F(u) \frac{dx}{du} \, du = \int \frac{F(u)}{f'(x)} \, du \quad \text{where } u = f(x)$$

$$16.7. \quad \int u^n \, du = \frac{u^{n+1}}{n+1}, \quad n \neq -1 \quad (\text{For } n = -1, \text{ see 16.8})$$

$$16.8. \quad \int \frac{du}{u} = \ln u \quad \text{if } u > 0 \text{ or } \ln(-u) \text{ if } u < 0 \\ = \ln |u|$$

$$16.9. \quad \int e^u \, du = e^u$$

$$16.10. \quad \int a^u \, du = \int e^{u \ln a} \, du = \frac{e^{u \ln a}}{\ln a} = \frac{a^u}{\ln a}, \quad a > 0, a \neq 1$$

**16.11.**  $\int \sin u du = -\cos u$

**16.12.**  $\int \cos u du = \sin u$

**16.13.**  $\int \tan u du = \ln \sec u = -\ln \cos u$

**16.14.**  $\int \cot u du = \ln \sin u$

**16.15.**  $\int \sec u du = \ln(\sec u + \tan u) = \ln \tan\left(\frac{u}{2} + \frac{\pi}{4}\right)$

**16.16.**  $\int \csc u du = \ln(\csc u - \cot u) = \ln \tan\frac{u}{2}$

**16.17.**  $\int \sec^2 u du = \tan u$

**16.18.**  $\int \csc^2 u du = -\cot u$

**16.19.**  $\int \tan^2 u du = \tan u - u$

**16.20.**  $\int \cot^2 u du = -\cot u - u$

**16.21.**  $\int \sin^2 u du = \frac{u}{2} - \frac{\sin 2u}{4} = \frac{1}{2}(u - \sin u \cos u)$

**16.22.**  $\int \cos^2 u du = \frac{u}{2} + \frac{\sin 2u}{4} = \frac{1}{2}(u + \sin u \cos u)$

**16.23.**  $\int \sec u \tan u du = \sec u$

**16.24.**  $\int \csc u \cot u du = -\csc u$

**16.25.**  $\int \sinh u du = \cosh u$

**16.26.**  $\int \cosh u du = \sinh u$

**16.27.**  $\int \tanh u du = \ln \cosh u$

**16.28.**  $\int \coth u du = \ln \sinh u$

**16.29.**  $\int \operatorname{sech} u du = \sin^{-1}(\tanh u) \text{ or } 2 \tan^{-1} e^u$

**16.30.**  $\int \operatorname{csch} u du = \ln \tanh\frac{u}{2} \text{ or } -\coth^{-1} e^u$

**16.31.**  $\int \operatorname{sech}^2 u du = \tanh u$

**16.32.**  $\int \operatorname{csch}^2 u du = -\coth u$

**16.33.**  $\int \tanh^2 u du = u - \tanh u$

**16.34.**  $\int \coth^2 u du = u - \coth u$

**16.35.**  $\int \sinh^2 u du = \frac{\sinh 2u}{4} - \frac{u}{2} = \frac{1}{2}(\sinh u \cosh u - u)$

**16.36.**  $\int \cosh^2 u du = \frac{\sinh 2u}{4} + \frac{u}{2} = \frac{1}{2}(\sinh u \cosh u + u)$

**16.37.**  $\int \operatorname{sech} u \tanh u du = -\operatorname{sech} u$

**16.38.**  $\int \operatorname{csch} u \coth u du = -\operatorname{csch} u$

**16.39.**  $\int \frac{du}{u^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{u}{a}$

**16.40.**  $\int \frac{du}{u^2 - a^2} = \frac{1}{2a} \ln \left( \frac{u-a}{u+a} \right) = -\frac{1}{a} \coth^{-1} \frac{u}{a} \quad u^2 > a^2$

**16.41.**  $\int \frac{du}{a^2 - u^2} = \frac{1}{2a} \ln \left( \frac{a+u}{a-u} \right) = \frac{1}{a} \tanh^{-1} \frac{u}{a} \quad u^2 < a^2$

**16.42.**  $\int \frac{du}{\sqrt{a^2 - u^2}} = \sin^{-1} \frac{u}{a}$

**16.43.**  $\int \frac{du}{\sqrt{u^2 + a^2}} = \ln(u + \sqrt{u^2 + a^2}) \quad \text{or} \quad \sinh^{-1} \frac{u}{a}$

**16.44.**  $\int \frac{du}{\sqrt{u^2 - a^2}} = \ln(u + \sqrt{u^2 - a^2})$

**16.45.**  $\int \frac{du}{u \sqrt{u^2 - a^2}} = \frac{1}{a} \sec^{-1} \left| \frac{u}{a} \right|$

**16.46.**  $\int \frac{du}{u \sqrt{u^2 + a^2}} = -\frac{1}{a} \ln \left( \frac{a + \sqrt{u^2 + a^2}}{u} \right)$

**16.47.**  $\int \frac{du}{u \sqrt{a^2 - u^2}} = -\frac{1}{a} \ln \left( \frac{a + \sqrt{a^2 - u^2}}{u} \right)$

**16.48.**  $\int f^{(n)} g dx = f^{(n-1)} g - f^{(n-2)} g' + f^{(n-3)} g'' - \cdots (-1)^n \int f g^{(n)} dx$

This is called *generalized integration by parts*.

### Important Transformations

---

Often in practice an integral can be simplified by using an appropriate transformation or substitution together with Formula 16.6. The following list gives some transformations and their effects.

$$16.49. \int F(ax + b) dx = \frac{1}{a} \int F(u) du \quad \text{where } u = ax + b$$

$$16.50. \int F(\sqrt{ax + b}) dx = \frac{2}{a} \int u F(u) du \quad \text{where } u = \sqrt{ax + b}$$

$$16.51. \int F(\sqrt[n]{ax + b}) dx = \frac{n}{a} \int u^{n-1} F(u) du \quad \text{where } u = \sqrt[n]{ax + b}$$

$$16.52. \int F(\sqrt{a^2 - x^2}) dx = a \int F(a \cos u) \cos u du \quad \text{where } x = a \sin u$$

$$16.53. \int F(\sqrt{x^2 + a^2}) dx = a \int F(a \sec u) \sec^2 u du \quad \text{where } x = a \tan u$$

$$16.54. \int F(\sqrt{x^2 - a^2}) dx = a \int F(a \tan u) \sec u \tan u du \quad \text{where } x = a \sec u$$

$$16.55. \int F(e^{ax}) dx = \frac{1}{a} \int \frac{F(u)}{u} du \quad \text{where } u = e^{ax}$$

$$16.56. \int F(\ln x) dx = \int F(u) e^u du \quad \text{where } u = \ln x$$

$$16.57. \int F\left(\sin^{-1} \frac{x}{a}\right) dx = a \int F(u) \cos u du \quad \text{where } u = \sin^{-1} \frac{x}{a}$$

Similar results apply for other inverse trigonometric functions.

$$16.58. \int F(\sin x, \cos x) dx = 2 \int F\left(\frac{2u}{1+u^2}, \frac{1-u^2}{1+u^2}\right) \frac{du}{1+u^2} \quad \text{where } u = \tan \frac{x}{2}$$

# 17

## TABLES of SPECIAL INDEFINITE INTEGRALS

Here we provide tables of special indefinite integrals. As stated in the remarks on page 67, here  $a, b, p, q, n$  are constants, restricted if indicated;  $e = 2.71828 \dots$  is the natural base of logarithms;  $\ln u$  denotes the natural logarithm of  $u$ , where it is assumed that  $u > 0$  (in general, to extend formulas to cases where  $u < 0$  as well, replace  $\ln u$  by  $\ln |u|$ ); all angles are in radians; and all constants of integration are omitted but implied. It is assumed in all cases that division by zero is excluded.

Our integrals are divided into types which involve the following algebraic expressions and functions:

(1) $ax + b$	(13) $\sqrt{ax^2 + bx + c}$	(25) $e^{ax}$
(2) $\sqrt{ax+b}$	(14) $x^3 + a^3$	(26) $\ln x$
(3) $ax + b$ and $px + q$	(15) $x^4 \pm a^4$	(27) $\sinh ax$
(4) $\sqrt{ax+b}$ and $px+q$	(16) $x^n \pm a^n$	(28) $\cosh ax$
(5) $\sqrt{ax+b}$ and $\sqrt{px+q}$	(17) $\sin ax$	(29) $\sinh ax$ and $\cosh ax$
(6) $x^2 + a^2$	(18) $\cos ax$	(30) $\tanh ax$
(7) $x^2 - a^2$ , with $x^2 > a^2$	(19) $\sin ax$ and $\cos ax$	(31) $\coth ax$
(8) $a^2 - x^2$ , with $x^2 < a^2$	(20) $\tan ax$	(32) $\operatorname{sech} ax$
(9) $\sqrt{x^2 + a^2}$	(21) $\cot ax$	(33) $\operatorname{csch} ax$
(10) $\sqrt{x^2 - a^2}$	(22) $\sec ax$	(34) inverse hyperbolic functions
(11) $\sqrt{a^2 - x^2}$	(23) $\csc ax$	
(12) $ax^2 + bx + c$	(24) inverse trigonometric functions	

Some integrals contain the Bernoulli numbers  $B_n$  and the Euler numbers  $E_n$  defined in Chapter 23.

### (1) Integrals Involving $ax + b$

---

$$17.1.1. \int \frac{dx}{ax+b} = \frac{1}{a} \ln(ax+b)$$

$$17.1.2. \int \frac{x dx}{ax+b} = \frac{x}{a} - \frac{b}{a^2} \ln(ax+b)$$

$$17.1.3. \int \frac{x^2 dx}{ax+b} = \frac{(ax+b)^2}{2a^3} - \frac{2b(ax+b)}{a^3} + \frac{b^2}{a^3} \ln(ax+b)$$

$$17.1.4. \int \frac{dx}{x(ax+b)} = \frac{1}{b} \ln\left(\frac{x}{ax+b}\right)$$

$$17.1.5. \int \frac{dx}{x^2(ax+b)} = -\frac{1}{bx} + \frac{a}{b^2} \ln\left(\frac{ax+b}{x}\right)$$

$$17.1.6. \int \frac{dx}{(ax+b)^2} = \frac{-1}{a(ax+b)}$$

$$17.1.7. \int \frac{x dx}{(ax+b)^2} = \frac{b}{a^2(ax+b)} + \frac{1}{a^2} \ln(ax+b)$$

$$17.1.8. \int \frac{x^2 dx}{(ax+b)^2} = \frac{ax+b}{a^3} - \frac{b^2}{a^3(ax+b)} - \frac{2b}{a^3} \ln(ax+b)$$

$$17.1.9. \int \frac{dx}{x(ax+b)^2} = \frac{1}{b(ax+b)} + \frac{1}{b^2} \ln\left(\frac{x}{ax+b}\right)$$

$$17.1.10. \int \frac{dx}{x^2(ax+b)^2} = \frac{-a}{b^2(ax+b)} - \frac{1}{b^2x} + \frac{2a}{b^3} \ln\left(\frac{ax+b}{x}\right)$$

$$17.1.11. \int \frac{dx}{(ax+b)^3} = \frac{-1}{2(ax+b)^2}$$

$$17.1.12. \int \frac{x dx}{(ax+b)^3} = \frac{-1}{a^2(ax+b)} + \frac{b}{2a^2(ax+b)^2}$$

$$17.1.13. \int \frac{x^2 dx}{(ax+b)^3} = \frac{2b}{a^3(ax+b)} - \frac{b^2}{2a^3(ax+b)^2} + \frac{1}{a^3} \ln(ax+b)$$

$$17.1.14. \int (ax+b)^n dx = \frac{(ax+b)^{n+1}}{(n+1)a}. \text{ If } n = -1, \text{ see 17.1.1.}$$

$$17.1.15. \int x(ax+b)^n dx = \frac{(ax+b)^{n+2}}{(n+2)a^2} - \frac{b(ax+b)^{n+1}}{(n+1)a^2}, \quad n \neq -1, -2$$

If  $n = -1, -2$ , see 17.1.2 and 17.1.7.

$$17.1.16. \int x^2(ax+b)^n dx = \frac{(ax+b)^{n+3}}{(n+3)a^3} - \frac{2b(ax+b)^{n+2}}{(n+2)a^3} + \frac{b^2(ax+b)^{n+1}}{(n+1)a^3}$$

If  $n = -1, -2, -3$ , see 17.1.3, 17.1.8, and 17.1.13.

$$17.1.17. \int x^m(ax+b)^n dx = \begin{cases} \frac{x^{m+1}(ax+b)^n}{m+n+1} + \frac{nb}{m+n+1} \int x^m(ax+b)^{n-1} dx \\ \frac{x^m(ax+b)^{n+1}}{(m+n+1)a} - \frac{mb}{(m+n+1)a} \int x^{m-1}(ax+b)^n dx \\ \frac{-x^{m+1}(ax+b)^{n+1}}{(n+1)b} + \frac{m+n+2}{(n+1)b} \int x^m(ax+b)^{n+1} dx \end{cases}$$

## (2) Integrals Involving $\sqrt{ax+b}$

---

$$17.2.1. \int \frac{dx}{\sqrt{ax+b}} = \frac{2\sqrt{ax+b}}{a}$$

$$17.2.2. \int \frac{x dx}{\sqrt{ax+b}} = \frac{2(ax-2b)}{3a^2} \sqrt{ax+b}$$

$$17.2.3. \int \frac{x^2 dx}{\sqrt{ax+b}} = \frac{2(3a^2x^2 - 4abx + 8b^2)}{15a^3} \sqrt{ax+b}$$

$$17.2.4. \int \frac{dx}{x\sqrt{ax+b}} = \begin{cases} \frac{1}{\sqrt{b}} \ln\left(\frac{\sqrt{ax+b} - \sqrt{b}}{\sqrt{ax+b} + \sqrt{b}}\right) \\ \frac{2}{\sqrt{-b}} \tan^{-1} \sqrt{\frac{ax+b}{-b}} \end{cases}$$

$$17.2.5. \int \frac{dx}{x^2\sqrt{ax+b}} = -\frac{\sqrt{ax+b}}{bx} - \frac{a}{2b} \int \frac{dx}{x\sqrt{ax+b}} \quad (\text{see 17.2.12.})$$

$$17.2.6. \int \sqrt{ax+b} dx = \frac{2\sqrt{(ax+b)^3}}{3a}$$

$$17.2.7. \int x\sqrt{ax+b} dx = \frac{2(3ax-2b)}{15a^2} \sqrt{(ax+b)^3}$$

$$17.2.8. \quad \int x^2 \sqrt{ax+b} dx = \frac{2(15a^2x^2 - 12abx + 8b^2)}{105a^3} \sqrt{(ax+b)^3}$$

$$17.2.9. \quad \int \frac{\sqrt{ax+b}}{x} dx = 2\sqrt{ax+b} + b \int \frac{dx}{x\sqrt{ax+b}} \quad (\text{See 17.2.12.})$$

$$17.2.10. \quad \int \frac{\sqrt{ax+b}}{x^2} dx = -\frac{\sqrt{ax+b}}{x} + \frac{a}{2} \int \frac{dx}{x\sqrt{ax+b}} \quad (\text{See 17.2.12.})$$

$$17.2.11. \quad \int \frac{x^m}{\sqrt{ax+b}} dx = \frac{2x^m \sqrt{ax+b}}{(2m+1)a} - \frac{2mb}{(2m+1)a} \int \frac{x^{m-1}}{\sqrt{ax+b}} dx$$

$$17.2.12. \quad \int \frac{dx}{x^m \sqrt{ax+b}} = -\frac{\sqrt{ax+b}}{(m-1)bx^{m-1}} - \frac{(2m-3)a}{(2m-2)b} \int \frac{dx}{x^{m-1} \sqrt{ax+b}}$$

$$17.2.13. \quad \int x^m \sqrt{ax+b} dx = \frac{2x^m}{(2m+3)a} (ax+b)^{3/2} - \frac{2mb}{(2m+3)a} \int x^{m-1} \sqrt{ax+b} dx$$

$$17.2.14. \quad \int \frac{\sqrt{ax+b}}{x^m} dx = -\frac{\sqrt{ax+b}}{(m-1)x^{m-1}} + \frac{a}{2(m-1)} \int \frac{dx}{x^{m-1} \sqrt{ax+b}}$$

$$17.2.15. \quad \int \frac{\sqrt{ax+b}}{x^m} dx = \frac{-(ax+b)^{3/2}}{(m-1)bx^{m-1}} - \frac{(2m-5)a}{(2m-2)b} \int \frac{\sqrt{ax+b}}{x^{m-1}} dx$$

$$17.2.16. \quad \int (ax+b)^{m/2} dx = \frac{2(ax+b)^{(m+2)/2}}{a^2(m+2)}$$

$$17.2.17. \quad \int x(ax+b)^{m/2} dx = \frac{2(ax+b)^{(m+4)/2}}{a^2(m+4)} - \frac{2b(ax+b)^{(m+2)/2}}{a^2(m+2)}$$

$$17.2.18. \quad \int x^2(ax+b)^{m/2} dx = \frac{2(ax+b)^{(m+6)/2}}{a^3(m+6)} - \frac{4b(ax+b)^{(m+4)/2}}{a^3(m+4)} + \frac{2b^2(ax+b)^{(m+2)/2}}{a^3(m+2)}$$

$$17.2.19. \quad \int \frac{(ax+b)^{m/2}}{x} dx = \frac{2(ax+b)^{m/2}}{m} + b \int \frac{(ax+b)^{(m-2)/2}}{x} dx$$

$$17.2.20. \quad \int \frac{(ax+b)^{m/2}}{x^2} dx = -\frac{(ax+b)^{(m+2)/2}}{bx} + \frac{ma}{2b} \int \frac{(ax+b)^{m/2}}{x} dx$$

$$17.2.21. \quad \int \frac{dx}{x(ax+b)^{m/2}} = \frac{2}{(m-2)b(ax+b)^{(m-2)/2}} + \frac{1}{b} \int \frac{dx}{x(ax+b)^{(m-2)/2}}$$

### (3) Integrals Involving $ax+b$ and $px+q$

---

$$17.3.1. \quad \int \frac{dx}{(ax+b)(px+q)} = \frac{1}{bp-aq} \ln \left( \frac{px+q}{ax+b} \right)$$

$$17.3.2. \quad \int \frac{x dx}{(ax+b)(px+q)} = \frac{1}{bp-aq} \left\{ \frac{b}{a} \ln(ax+b) - \frac{q}{p} \ln(px+q) \right\}$$

- 17.3.3.  $\int \frac{dx}{(ax+b)^2(px+q)} = \frac{1}{bp-aq} \left\{ \frac{1}{ax+b} + \frac{p}{bp-aq} \ln \left( \frac{px+q}{ax+b} \right) \right\}$
- 17.3.4.  $\int \frac{x dx}{(ax+b)^2(px+q)} = \frac{1}{bp-aq} \left\{ \frac{q}{bp-aq} \ln \left( \frac{ax+b}{px+q} \right) - \frac{b}{a(ax+b)} \right\}$
- 17.3.5.  $\int \frac{x^2 dx}{(ax+b)^2(px+q)} = \frac{b^2}{(bp-aq)a^2(ax+b)} + \frac{1}{(bp-aq)^2} \left\{ \frac{q^2}{p} \ln(px+q) + \frac{b(bp-2aq)}{a^2} \ln(ax+b) \right\}$
- 17.3.6.  $\int \frac{dx}{(ax+b)^m(px+q)^n} = \frac{-1}{(n-1)(bp-aq)} \left\{ \frac{1}{(ax+b)^{m-1}(px+q)^{n-1}} \right.$   

$$\left. + a(m+n-2) \int \frac{dx}{(ax+b)^m(px+q)^{n-1}} \right\}$$
- 17.3.7.  $\int \frac{ax+b}{px+q} dx = \frac{ax}{p} + \frac{bp-aq}{p^2} \ln(px+q)$   

$$\begin{aligned} & \quad \left. \frac{-1}{(n-1)(bp-aq)} \left\{ \frac{(ax+b)^{m+1}}{(px+q)^{n-1}} + (n-m-2)a \int \frac{(ax+b)^m}{(px+q)^{n-1}} dx \right\} \right. \\ 17.3.8. \quad \int \frac{(ax+b)^m}{(px+q)^n} dx = & \left. \frac{-1}{(n-m-1)p} \left\{ \frac{(ax+b)^m}{(px+q)^{n-1}} + m(bp-aq) \int \frac{(ax+b)^{m-1}}{(px+q)^n} dx \right\} \right. \\ & \left. \frac{-1}{(n-1)p} \left\{ \frac{(ax+b)^m}{(px+q)^{n-1}} - ma \int \frac{(ax+b)^{m-1}}{(px+q)^{n-1}} dx \right\} \right. \end{aligned}$$

#### (4) Integrals Involving $\sqrt{ax+b}$ and $px+q$

---

- 17.4.1.  $\int \frac{px+q}{\sqrt{ax+b}} dx = \frac{2(apx+3aq-2bp)}{3a^2} \sqrt{ax+b}$
- 17.4.2.  $\int \frac{dx}{(px+q)\sqrt{ax+b}} = \begin{cases} \frac{1}{\sqrt{bp-aq}\sqrt{p}} \ln \left( \frac{\sqrt{p(ax+b)} - \sqrt{bp-aq}}{\sqrt{p(ax+b)} + \sqrt{bp-aq}} \right) \\ \frac{2}{\sqrt{aq-bp}\sqrt{p}} \tan^{-1} \sqrt{\frac{p(ax+b)}{aq-bp}} \end{cases}$
- 17.4.3.  $\int \frac{\sqrt{ax+b}}{px+q} dx = \begin{cases} \frac{2\sqrt{ax+b}}{p} + \frac{\sqrt{bp-aq}}{p\sqrt{p}} \ln \left( \frac{\sqrt{p(ax+b)} - \sqrt{bp-aq}}{\sqrt{p(ax+b)} + \sqrt{bp-aq}} \right) \\ \frac{2\sqrt{ax+b}}{p} - \frac{2\sqrt{aq-bp}}{p\sqrt{p}} \tan^{-1} \sqrt{\frac{p(ax+b)}{aq-bp}} \end{cases}$
- 17.4.4.  $\int (px+q)^n \sqrt{ax+b} dx = \frac{2(px+q)^{n+1} \sqrt{ax+b}}{(2n+3)p} + \frac{bp-aq}{(2n+3)p} \int \frac{(px+q)^n}{\sqrt{ax+b}}$
- 17.4.5.  $\int \frac{dx}{(px+q)^n \sqrt{ax+b}} = \frac{\sqrt{ax+b}}{(n-1)(aq-bp)(px+q)^{n-1}} + \frac{(2n-3)a}{2(n-1)(aq-bp)} \int \frac{dx}{(px+q)^{n-1} \sqrt{ax+b}}$
- 17.4.6.  $\int \frac{(px+q)^n}{\sqrt{ax+b}} dx = \frac{2(px+q)^n \sqrt{ax+b}}{(2n+1)a} + \frac{2n(aq-bp)}{(2n+1)a} \int \frac{(px+q)^{n-1} dx}{\sqrt{ax+b}}$
- 17.4.7.  $\int \frac{\sqrt{ax+b}}{(px+q)^n} dx = \frac{-\sqrt{ax+b}}{(n-1)p(px+q)^{n-1}} + \frac{a}{2(n-1)p} \int \frac{dx}{(px+q)^{n-1} \sqrt{ax+b}}$

**(5) Integrals Involving  $\sqrt{ax+b}$  and  $\sqrt{px+q}$** 

---

$$17.5.1. \int \frac{dx}{\sqrt{(ax+b)(px+q)}} = \begin{cases} \frac{2}{\sqrt{ap}} \ln(\sqrt{a(px+q)} + \sqrt{p(ax+b)}) \\ \frac{2}{\sqrt{-ap}} \tan^{-1} \sqrt{\frac{-p(ax+b)}{a(px+q)}} \end{cases}$$

$$17.5.2. \int \frac{x dx}{\sqrt{(ax+b)(px+q)}} = \frac{\sqrt{(ax+b)(px+q)}}{ap} - \frac{bp+aq}{2ap} \int \frac{dx}{\sqrt{(ax+b)(px+q)}}$$

$$17.5.3. \int \sqrt{(ax+b)(px+q)} dx = \frac{2apx + bp + aq}{4ap} \sqrt{(ax+b)(px+q)} - \frac{(bp-aq)^2}{8ap} \int \frac{dx}{\sqrt{(ax+b)(px+q)}}$$

$$17.5.4. \int \sqrt{\frac{px+q}{ax+b}} dx = \frac{\sqrt{(ax+b)(px+q)}}{a} + \frac{aq-bp}{2a} \int \frac{dx}{\sqrt{(ax+b)(px+q)}}$$

$$17.5.5. \int \frac{dx}{(px+q)\sqrt{(ax+b)(px+q)}} = \frac{2\sqrt{ax+b}}{(aq-bp)\sqrt{px+q}}$$

**(6) Integrals Involving  $x^2 + a^2$** 

---

$$17.6.1. \int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a}$$

$$17.6.2. \int \frac{x dx}{x^2 + a^2} = \frac{1}{2} \ln(x^2 + a^2)$$

$$17.6.3. \int \frac{x^2 dx}{x^2 + a^2} = x - a \tan^{-1} \frac{x}{a}$$

$$17.6.4. \int \frac{x^3 dx}{x^2 + a^2} = \frac{x^2}{2} - \frac{a^2}{2} \ln(x^2 + a^2)$$

$$17.6.5. \int \frac{dx}{x(x^2 + a^2)} = \frac{1}{2a^2} \ln \left( \frac{x^2}{x^2 + a^2} \right)$$

$$17.6.6. \int \frac{dx}{x^2(x^2 + a^2)} = -\frac{1}{a^2 x} - \frac{1}{a^3} \tan^{-1} \frac{x}{a}$$

$$17.6.7. \int \frac{dx}{x^3(x^2 + a^2)} = -\frac{1}{2a^2 x^2} - \frac{1}{2a^4} \ln \left( \frac{x^2}{x^2 + a^2} \right)$$

$$17.6.8. \int \frac{dx}{(x^2 + a^2)^2} = \frac{x}{2a^2(x^2 + a^2)} + \frac{1}{2a^3} \tan^{-1} \frac{x}{a}$$

$$17.6.9. \int \frac{x dx}{(x^2 + a^2)^2} = \frac{-1}{2(x^2 + a^2)}$$

$$17.6.10. \int \frac{x^2 dx}{(x^2 + a^2)^2} = \frac{-x}{2(x^2 + a^2)} + \frac{1}{2a} \tan^{-1} \frac{x}{a}$$

$$17.6.11. \int \frac{x^3 dx}{(x^2 + a^2)^2} = \frac{a^2}{2(x^2 + a^2)} + \frac{1}{2} \ln(x^2 + a^2)$$

$$17.6.12. \int \frac{dx}{x(x^2 + a^2)^2} = \frac{1}{2a^2(x^2 + a^2)} + \frac{1}{2a^4} \ln\left(\frac{x^2}{x^2 + a^2}\right)$$

$$17.6.13. \int \frac{dx}{x^2(x^2 + a^2)^2} = -\frac{1}{a^4 x} - \frac{x}{2a^4(x^2 + a^2)} - \frac{3}{2a^5} \tan^{-1} \frac{x}{a}$$

$$17.6.14. \int \frac{dx}{x^3(x^2 + a^2)^2} = -\frac{1}{2a^4 x^2} - \frac{1}{2a^4(x^2 + a^2)} - \frac{1}{a^6} \ln\left(\frac{x^2}{x^2 + a^2}\right)$$

$$17.6.15. \int \frac{dx}{(x^2 + a^2)^n} = \frac{x}{2(n-1)a^2(x^2 + a^2)^{n-1}} + \frac{2n-3}{(2n-2)a^2} \int \frac{dx}{(x^2 + a^2)^{n-1}}$$

$$17.6.16. \int \frac{x dx}{(x^2 + a^2)^n} = \frac{-1}{2(n-1)(x^2 + a^2)^{n-1}}$$

$$17.6.17. \int \frac{dx}{x(x^2 + a^2)^n} = \frac{1}{2(n-1)a^2(x^2 + a^2)^{n-1}} + \frac{1}{a^2} \int \frac{dx}{x(x^2 + a^2)^{n-1}}$$

$$17.6.18. \int \frac{x^m dx}{(x^2 + a^2)^n} = \int \frac{x^{m-2} dx}{(x^2 + a^2)^{n-1}} - a^2 \int \frac{x^{m-2} dx}{(x^2 + a^2)^n}$$

$$17.6.19. \int \frac{dx}{x^m(x^2 + a^2)^n} = \frac{1}{a^2} \int \frac{dx}{x^m(x^2 + a^2)^{n-1}} - \frac{1}{a^2} \int \frac{dx}{x^{m-2}(x^2 + a^2)^n}$$

### (7) Integrals Involving $x^2 - a^2$ , $x^2 > a^2$

---

$$17.7.1. \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \ln\left(\frac{x-a}{x+a}\right) \quad \text{or} \quad -\frac{1}{a} \coth^{-1} \frac{x}{a}$$

$$17.7.2. \int \frac{x dx}{x^2 - a^2} = \frac{1}{2} \ln(x^2 - a^2)$$

$$17.7.3. \int \frac{x^2 dx}{x^2 - a^2} = x + \frac{a}{2} \ln\left(\frac{x-a}{x+a}\right)$$

$$17.7.4. \int \frac{x^3 dx}{x^2 - a^2} = \frac{x^2}{2} + \frac{a^2}{2} \ln(x^2 - a^2)$$

$$17.7.5. \int \frac{dx}{x(x^2 - a^2)} = \frac{1}{2a^2} \ln\left(\frac{x^2 - a^2}{x^2}\right)$$

$$17.7.6. \int \frac{dx}{x^2(x^2 - a^2)} = \frac{1}{a^2 x} + \frac{1}{2a^3} \ln\left(\frac{x-a}{x+a}\right)$$

$$17.7.7. \int \frac{dx}{x^3(x^2 - a^2)} = \frac{1}{2a^2 x^2} - \frac{1}{2a^4} \ln\left(\frac{x^2}{x^2 - a^2}\right)$$

$$17.7.8. \int \frac{dx}{(x^2 - a^2)^2} = \frac{-x}{2a^2(x^2 - a^2)} - \frac{1}{4a^3} \ln\left(\frac{x-a}{x+a}\right)$$

$$17.7.9. \quad \int \frac{x dx}{(x^2 - a^2)^2} = \frac{-1}{2(x^2 - a^2)}$$

$$17.7.10. \quad \int \frac{x^2 dx}{(x^2 - a^2)^2} = \frac{-x}{2(x^2 - a^2)} + \frac{1}{4a} \ln\left(\frac{x-a}{x+a}\right)$$

$$17.7.11. \quad \int \frac{x^3 dx}{(x^2 - a^2)^2} = \frac{-a^2}{2(x^2 - a^2)} + \frac{1}{2} \ln(x^2 - a^2)$$

$$17.7.12. \quad \int \frac{dx}{x(x^2 - a^2)^2} = \frac{-1}{2a^2(x^2 - a^2)} + \frac{1}{2a^4} \ln\left(\frac{x^2}{x^2 - a^2}\right)$$

$$17.7.13. \quad \int \frac{dx}{x^2(x^2 - a^2)^2} = -\frac{1}{a^4 x} - \frac{x}{2a^4(x^2 - a^2)} - \frac{3}{4a^5} \ln\left(\frac{x-a}{x+a}\right)$$

$$17.7.14. \quad \int \frac{dx}{x^3(x^2 - a^2)^2} = -\frac{1}{2a^4 x^2} - \frac{1}{2a^4(x^2 - a^2)} + \frac{1}{a^6} \ln\left(\frac{x^2}{x^2 - a^2}\right)$$

$$17.7.15. \quad \int \frac{dx}{(x^2 - a^2)^n} = \frac{-x}{2(n-1)a^2(x^2 - a^2)^{n-1}} - \frac{2n-3}{(2n-2)a^2} \int \frac{dx}{(x^2 - a^2)^{n-1}}$$

$$17.7.16. \quad \int \frac{x dx}{(x^2 - a^2)^n} = \frac{-1}{2(n-1)(x^2 - a^2)^{n-1}}$$

$$17.7.17. \quad \int \frac{dx}{x(x^2 - a^2)^n} = \frac{-1}{2(n-1)a^2(x^2 - a^2)^{n-1}} - \frac{1}{a^2} \int \frac{dx}{x(x^2 - a^2)^{n-1}}$$

$$17.7.18. \quad \int \frac{x^m dx}{(x^2 - a^2)^n} = \int \frac{x^{m-2} dx}{(x^2 - a^2)^{n-1}} + a^2 \int \frac{x^{m-2} dx}{(x^2 - a^2)^n}$$

$$17.7.19. \quad \int \frac{dx}{x^m(x^2 - a^2)^n} = \frac{1}{a^2} \int \frac{dx}{x^{m-2}(x^2 - a^2)^n} - \frac{1}{a^2} \int \frac{dx}{x^m(x^2 - a^2)^{n-1}}$$

### (8) Integrals Involving $x^2 - a^2, x^2 < a^2$

$$17.8.1. \quad \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln\left(\frac{a+x}{a-x}\right) \quad \text{or} \quad \frac{1}{a} \tanh^{-1} \frac{x}{a}$$

$$17.8.2. \quad \int \frac{x dx}{a^2 - x^2} = -\frac{1}{2} \ln(a^2 - x^2)$$

$$17.8.3. \quad \int \frac{x^2 dx}{a^2 - x^2} = -x + \frac{a}{2} \ln\left(\frac{a+x}{a-x}\right)$$

$$17.8.4. \quad \int \frac{x^3 dx}{a^2 - x^2} = -\frac{x^2}{2} - \frac{a^2}{2} \ln(a^2 - x^2)$$

$$17.8.5. \quad \int \frac{dx}{x(a^2 - x^2)} = \frac{1}{2a^2} \ln\left(\frac{x^2}{a^2 - x^2}\right)$$

$$17.8.6. \quad \int \frac{dx}{x^2(a^2 - x^2)} = -\frac{1}{a^2 x} + \frac{1}{2a^3} \ln\left(\frac{a+x}{a-x}\right)$$

$$17.8.7. \int \frac{dx}{x^3(a^2 - x^2)} = -\frac{1}{2a^2x^2} + \frac{1}{2a^4} \ln\left(\frac{x^2}{a^2 - x^2}\right)$$

$$17.8.8. \int \frac{dx}{(a^2 - x^2)^2} = \frac{x}{2a^2(a^2 - x^2)} + \frac{1}{4a^3} \ln\left(\frac{a+x}{a-x}\right)$$

$$17.8.9. \int \frac{x dx}{(a^2 - x^2)^2} = \frac{1}{2(a^2 - x^2)}$$

$$17.8.10. \int \frac{x^2 dx}{(a^2 - x^2)^2} = \frac{x}{2(a^2 - x^2)} - \frac{1}{4a} \ln\left(\frac{a+x}{a-x}\right)$$

$$17.8.11. \int \frac{x^3 dx}{(a^2 - x^2)^2} = \frac{a^2}{2(a^2 - x^2)} + \frac{1}{2} \ln(a^2 - x^2)$$

$$17.8.12. \int \frac{dx}{x(a^2 - x^2)^2} = \frac{1}{2a^2(a^2 - x^2)} + \frac{1}{2a^4} \ln\left(\frac{x^2}{a^2 - x^2}\right)$$

$$17.8.13. \int \frac{dx}{x^2(a^2 - x^2)^2} = \frac{-1}{a^4 x} + \frac{x}{2a^4(a^2 - x^2)} + \frac{3}{4a^5} \ln\left(\frac{a+x}{a-x}\right)$$

$$17.8.14. \int \frac{dx}{x^3(a^2 - x^2)^2} = \frac{-1}{2a^4 x^2} + \frac{1}{2a^4(a^2 - x^2)} + \frac{1}{a^6} \ln\left(\frac{x^2}{a^2 - x^2}\right)$$

$$17.8.15. \int \frac{dx}{(a^2 - x^2)^n} = \frac{x}{2(n-1)a^2(a^2 - x^2)^{n-1}} + \frac{2n-3}{(2n-2)a^2} \int \frac{dx}{(a^2 - x^2)^{n-1}}$$

$$17.8.16. \int \frac{x dx}{(a^2 - x^2)^n} = \frac{1}{2(n-1)(a^2 - x^2)^{n-1}}$$

$$17.8.17. \int \frac{dx}{x(a^2 - x^2)^n} = \frac{1}{2(n-1)a^2(a^2 - x^2)^{n-1}} + \frac{1}{a^2} \int \frac{dx}{x(a^2 - x^2)^{n-1}}$$

$$17.8.18. \int \frac{x^m dx}{(a^2 - x^2)^n} = a^2 \int \frac{x^{m-2} dx}{(a^2 - x^2)^n} - \int \frac{x^{m-2} dx}{(a^2 - x^2)^{n-1}}$$

$$17.8.19. \int \frac{dx}{x^m(a^2 - x^2)^n} = \frac{1}{a^2} \int \frac{dx}{x^m(a^2 - x^2)^{n-1}} + \frac{1}{a^2} \int \frac{dx}{x^{m-2}(a^2 - x^2)^n}$$

## (9) Integrals Involving $\sqrt{x^2 + a^2}$

---

$$17.9.1. \int \frac{dx}{\sqrt{x^2 + a^2}} = \ln(x + \sqrt{x^2 + a^2}) \quad \text{or} \quad \sinh^{-1} \frac{x}{a}$$

$$17.9.2. \int \frac{x dx}{\sqrt{x^2 + a^2}} = \sqrt{x^2 + a^2}$$

$$17.9.3. \int \frac{x^2 dx}{\sqrt{x^2 + a^2}} = \frac{x\sqrt{x^2 + a^2}}{2} - \frac{a^2}{2} \ln(x + \sqrt{x^2 + a^2})$$

$$17.9.4. \int \frac{x^3 dx}{\sqrt{x^2 + a^2}} = \frac{(x^2 + a^2)^{3/2}}{3} - a^2 \sqrt{x^2 + a^2}$$

$$17.9.5. \quad \int \frac{dx}{x\sqrt{x^2+a^2}} = -\frac{1}{a} \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.6. \quad \int \frac{dx}{x^2\sqrt{x^2+a^2}} = -\frac{\sqrt{x^2+a^2}}{a^2 x}$$

$$17.9.7. \quad \int \frac{dx}{x^3\sqrt{x^2+a^2}} = -\frac{\sqrt{x^2+a^2}}{2a^2 x^2} + \frac{1}{2a^3} \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.8. \quad \int \sqrt{x^2+a^2} dx = \frac{x\sqrt{x^2+a^2}}{2} + \frac{a^2}{2} \ln(x+\sqrt{x^2+a^2})$$

$$17.9.9. \quad \int x\sqrt{x^2+a^2} dx = \frac{(x^2+a^2)^{3/2}}{3}$$

$$17.9.10. \quad \int x^2\sqrt{x^2+a^2} dx = \frac{x(x^2+a^2)^{3/2}}{4} - \frac{a^2 x \sqrt{x^2+a^2}}{8} - \frac{a^4}{8} \ln(x+\sqrt{x^2+a^2})$$

$$17.9.11. \quad \int x^3\sqrt{x^2+a^2} dx = \frac{(x^2+a^2)^{5/2}}{5} - \frac{a^2(x^2+a^2)^{3/2}}{3}$$

$$17.9.12. \quad \int \frac{\sqrt{x^2+a^2}}{x} dx = \sqrt{x^2+a^2} - a \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.13. \quad \int \frac{\sqrt{x^2+a^2}}{x^2} dx = -\frac{\sqrt{x^2+a^2}}{x} + \ln(x+\sqrt{x^2+a^2})$$

$$17.9.14. \quad \int \frac{\sqrt{x^2+a^2}}{x^3} dx = -\frac{\sqrt{x^2+a^2}}{2x^2} - \frac{1}{2a} \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.15. \quad \int \frac{dx}{(x^2+a^2)^{3/2}} = \frac{x}{a^2\sqrt{x^2+a^2}}$$

$$17.9.16. \quad \int \frac{x dx}{(x^2+a^2)^{3/2}} = \frac{-1}{\sqrt{x^2+a^2}}$$

$$17.9.17. \quad \int \frac{x^2 dx}{(x^2+a^2)^{3/2}} = \frac{-x}{\sqrt{x^2+a^2}} + \ln(x+\sqrt{x^2+a^2})$$

$$17.9.18. \quad \int \frac{x^3 dx}{(x^2+a^2)^{3/2}} = \sqrt{x^2+a^2} + \frac{a^2}{\sqrt{x^2+a^2}}$$

$$17.9.19. \quad \int \frac{dx}{x(x^2+a^2)^{3/2}} = \frac{1}{a^2\sqrt{x^2+a^2}} - \frac{1}{a^3} \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.20. \quad \int \frac{dx}{x^2(x^2+a^2)^{3/2}} = -\frac{\sqrt{x^2+a^2}}{a^4 x} - \frac{x}{a^4\sqrt{x^2+a^2}}$$

$$17.9.21. \quad \int \frac{dx}{x^3(x^2+a^2)^{3/2}} = \frac{-1}{2a^2 x^2 \sqrt{x^2+a^2}} - \frac{3}{2a^4 \sqrt{x^2+a^2}} + \frac{3}{2a^5} \ln \left( \frac{a+\sqrt{x^2+a^2}}{x} \right)$$

$$17.9.22. \quad \int (x^2+a^2)^{3/2} dx = \frac{x(x^2+a^2)^{3/2}}{4} + \frac{3a^2 x \sqrt{x^2+a^2}}{8} + \frac{3}{8} a^4 \ln(x+\sqrt{x^2+a^2})$$

$$17.9.23. \quad \int x(x^2+a^2)^{3/2} dx = \frac{(x^2+a^2)^{5/2}}{5}$$

$$17.9.24. \int x^2(x^2 + a^2)^{3/2} dx = \frac{x(x^2 + a^2)^{5/2}}{6} - \frac{a^2 x(x^2 + a^2)^{3/2}}{24} - \frac{a^4 x \sqrt{x^2 + a^2}}{16} - \frac{a^6}{16} \ln(x + \sqrt{x^2 + a^2})$$

$$17.9.25. \int x^3(x^2 + a^2)^{3/2} dx = \frac{(x^2 + a^2)^{7/2}}{7} - \frac{a^2(x^2 + a^2)^{5/2}}{5}$$

$$17.9.26. \int \frac{(x^2 + a^2)^{3/2}}{x} dx = \frac{(x^2 + a^2)^{3/2}}{3} + a^2 \sqrt{x^2 + a^2} - a^3 \ln\left(\frac{a + \sqrt{x^2 + a^2}}{x}\right)$$

$$17.9.27. \int \frac{(x^2 + a^2)^{3/2}}{x^2} dx = -\frac{(x^2 + a^2)^{3/2}}{x} + \frac{3x\sqrt{x^2 + a^2}}{2} + \frac{3}{2}a^2 \ln(x + \sqrt{x^2 + a^2})$$

$$17.9.28. \int \frac{(x^2 + a^2)^{3/2}}{x^3} dx = -\frac{(x^2 + a^2)^{3/2}}{2x^2} + \frac{3}{2}\sqrt{x^2 + a^2} - \frac{3}{2}a \ln\left(\frac{a + \sqrt{x^2 + a^2}}{x}\right)$$

### (10) Integrals Involving $\sqrt{x^2 - a^2}$

---

$$17.10.1. \int \frac{dx}{\sqrt{x^2 - a^2}} = \ln(x + \sqrt{x^2 - a^2}), \int \frac{x dx}{\sqrt{x^2 - a^2}} = \sqrt{x^2 - a^2}$$

$$17.10.2. \int \frac{x^2 dx}{\sqrt{x^2 - a^2}} = \frac{x\sqrt{x^2 - a^2}}{2} + \frac{a^2}{2} \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.3. \int \frac{x^3 dx}{\sqrt{x^2 - a^2}} = \frac{(x^2 - a^2)^{3/2}}{3} + a^2 \sqrt{x^2 - a^2}$$

$$17.10.4. \int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \sec^{-1}\left|\frac{x}{a}\right|$$

$$17.10.5. \int \frac{dx}{x^2\sqrt{x^2 - a^2}} = \frac{\sqrt{x^2 - a^2}}{a^2 x}$$

$$17.10.6. \int \frac{dx}{x^3\sqrt{x^2 - a^2}} = \frac{\sqrt{x^2 - a^2}}{2a^2 x^2} + \frac{1}{2a^3} \sec^{-1}\left|\frac{x}{a}\right|$$

$$17.10.7. \int \sqrt{x^2 - a^2} dx = \frac{x\sqrt{x^2 - a^2}}{2} - \frac{a^2}{2} \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.8. \int x\sqrt{x^2 - a^2} dx = \frac{(x^2 - a^2)^{3/2}}{3}$$

$$17.10.9. \int x^2\sqrt{x^2 - a^2} dx = \frac{x(x^2 - a^2)^{3/2}}{4} + \frac{a^2 x \sqrt{x^2 - a^2}}{8} - \frac{a^4}{8} \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.10. \int x^3\sqrt{x^2 - a^2} dx = \frac{(x^2 - a^2)^{5/2}}{5} + \frac{a^2(x^2 - a^2)^{3/2}}{3}$$

$$17.10.11. \int \frac{\sqrt{x^2 - a^2}}{x} dx = \sqrt{x^2 - a^2} - a \sec^{-1}\left|\frac{x}{a}\right|$$

$$17.10.12. \int \frac{\sqrt{x^2 - a^2}}{x^2} dx = -\frac{\sqrt{x^2 - a^2}}{x} + \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.13. \int \frac{\sqrt{x^2 - a^2}}{x^3} dx = -\frac{\sqrt{x^2 - a^2}}{2x^2} + \frac{1}{2a} \sec^{-1} \left| \frac{x}{a} \right|$$

$$17.10.14. \int \frac{dx}{(x^2 - a^2)^{3/2}} = -\frac{x}{a^2 \sqrt{x^2 - a^2}}$$

$$17.10.15. \int \frac{x dx}{(x^2 - a^2)^{3/2}} = \frac{-1}{\sqrt{x^2 - a^2}}$$

$$17.10.16. \int \frac{x^2 dx}{(x^2 - a^2)^{3/2}} = -\frac{x}{\sqrt{x^2 - a^2}} + \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.17. \int \frac{x^3 dx}{(x^2 - a^2)^{3/2}} = \sqrt{x^2 - a^2} - \frac{a^2}{\sqrt{x^2 - a^2}}$$

$$17.10.18. \int \frac{dx}{x(x^2 - a^2)^{3/2}} = \frac{-1}{a^2 \sqrt{x^2 - a^2}} - \frac{1}{a^3} \sec^{-1} \left| \frac{x}{a} \right|$$

$$17.10.19. \int \frac{dx}{x^2(x^2 - a^2)^{3/2}} = -\frac{\sqrt{x^2 - a^2}}{a^4 x} - \frac{x}{a^4 \sqrt{x^2 - a^2}}$$

$$17.10.20. \int \frac{dx}{x^3(x^2 - a^2)^{3/2}} = \frac{1}{2a^2 x^2 \sqrt{x^2 - a^2}} - \frac{3}{2a^4 \sqrt{x^2 - a^2}} - \frac{3}{2a^5} \sec^{-1} \left| \frac{x}{a} \right|$$

$$17.10.21. \int (x^2 - a^2)^{3/2} dx = \frac{x(x^2 - a^2)^{3/2}}{4} - \frac{3a^2 x \sqrt{x^2 - a^2}}{8} + \frac{3}{8} a^4 \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.22. \int x(x^2 - a^2)^{3/2} dx = \frac{(x^2 - a^2)^{5/2}}{5}$$

$$17.10.23. \int x^2(x^2 - a^2)^{3/2} dx = \frac{x(x^2 - a^2)^{5/2}}{6} + \frac{a^2 x(x^2 - a^2)^{3/2}}{24} - \frac{a^4 x \sqrt{x^2 - a^2}}{16} + \frac{a^6}{16} \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.24. \int x^3(x^2 - a^2)^{3/2} dx = \frac{(x^2 - a^2)^{7/2}}{7} + \frac{a^2(x^2 - a^2)^{5/2}}{5}$$

$$17.10.25. \int \frac{(x^2 - a^2)^{3/2}}{x} dx = \frac{(x^2 - a^2)^{3/2}}{3} - a^2 \sqrt{x^2 - a^2} + a^3 \sec^{-1} \left| \frac{x}{a} \right|$$

$$17.10.26. \int \frac{(x^2 - a^2)^{3/2}}{x^2} dx = -\frac{(x^2 - a^2)^{3/2}}{x} + \frac{3x \sqrt{x^2 - a^2}}{2} - \frac{3}{2} a^2 \ln(x + \sqrt{x^2 - a^2})$$

$$17.10.27. \int \frac{(x^2 - a^2)^{3/2}}{x^3} dx = -\frac{(x^2 - a^2)^{3/2}}{2x^2} + \frac{3\sqrt{x^2 - a^2}}{2} - \frac{3}{2} a \sec^{-1} \left| \frac{x}{a} \right|$$

### (11) Integrals Involving $\sqrt{a^2 - x^2}$

---

$$17.11.1. \int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a}$$

$$17.11.2. \int \frac{x dx}{\sqrt{a^2 - x^2}} = -\sqrt{a^2 - x^2}$$

$$17.11.3. \int \frac{x^2 dx}{\sqrt{a^2 - x^2}} = -\frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}$$

$$17.11.4. \int \frac{x^3 dx}{\sqrt{a^2 - x^2}} = \frac{(a^2 - x^2)^{3/2}}{3} - a^2 \sqrt{a^2 - x^2}$$

$$17.11.5. \int \frac{dx}{x\sqrt{a^2 - x^2}} = -\frac{1}{a} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.6. \int \frac{dx}{x^2 \sqrt{a^2 - x^2}} = -\frac{\sqrt{a^2 - x^2}}{a^2 x}$$

$$17.11.7. \int \frac{dx}{x^3 \sqrt{a^2 - x^2}} = -\frac{\sqrt{a^2 - x^2}}{2a^2 x^2} - \frac{1}{2a^3} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.8. \int \sqrt{a^2 - x^2} dx = \frac{x\sqrt{a^2 - x^2}}{2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a}$$

$$17.11.9. \int x\sqrt{a^2 - x^2} dx = -\frac{(a^2 - x^2)^{3/2}}{3}$$

$$17.11.10. \int x^2 \sqrt{a^2 - x^2} dx = -\frac{x(a^2 - x^2)^{3/2}}{4} + \frac{a^2 x \sqrt{a^2 - x^2}}{8} + \frac{a^4}{8} \sin^{-1} \frac{x}{a}$$

$$17.11.11. \int x^3 \sqrt{a^2 - x^2} dx = \frac{(a^2 - x^2)^{5/2}}{5} - \frac{a^2 (a^2 - x^2)^{3/2}}{3}$$

$$17.11.12. \int \frac{\sqrt{a^2 - x^2}}{x} dx = \sqrt{a^2 - x^2} - a \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.13. \int \frac{\sqrt{a^2 - x^2}}{x^2} dx = -\frac{\sqrt{a^2 - x^2}}{x} - \sin^{-1} \frac{x}{a}$$

$$17.11.14. \int \frac{\sqrt{a^2 - x^2}}{x^3} dx = -\frac{\sqrt{a^2 - x^2}}{2x^2} + \frac{1}{2a} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.15. \int \frac{dx}{(a^2 - x^2)^{3/2}} = \frac{x}{a^2 \sqrt{a^2 - x^2}}$$

$$17.11.16. \int \frac{x dx}{(a^2 - x^2)^{3/2}} = \frac{1}{\sqrt{a^2 - x^2}}$$

$$17.11.17. \int \frac{x^2 dx}{(a^2 - x^2)^{3/2}} = \frac{x}{\sqrt{a^2 - x^2}} - \sin^{-1} \frac{x}{a}$$

$$17.11.18. \int \frac{x^3 dx}{(a^2 - x^2)^{3/2}} = \sqrt{a^2 - x^2} + \frac{a^2}{\sqrt{a^2 - x^2}}$$

$$17.11.19. \int \frac{dx}{x(a^2 - x^2)^{3/2}} = \frac{1}{a^2 \sqrt{a^2 - x^2}} - \frac{1}{a^3} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.20. \int \frac{dx}{x^2(a^2 - x^2)^{3/2}} = -\frac{\sqrt{a^2 - x^2}}{a^4 x} + \frac{x}{a^4 \sqrt{a^2 - x^2}}$$

$$17.11.21. \int \frac{dx}{x^3(a^2 - x^2)^{3/2}} = \frac{-1}{2a^2 x^2 \sqrt{a^2 - x^2}} + \frac{3}{2a^4 \sqrt{a^2 - x^2}} - \frac{3}{2a^5} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.22. \int (a^2 - x^2)^{3/2} dx = \frac{x(a^2 - x^2)^{3/2}}{4} + \frac{3a^2 x \sqrt{a^2 - x^2}}{8} + \frac{3}{8} a^4 \sin^{-1} \frac{x}{a}$$

$$17.11.23. \int x(a^2 - x^2)^{3/2} dx = -\frac{(a^2 - x^2)^{5/2}}{5}$$

$$17.11.24. \int x^2(a^2 - x^2)^{3/2} dx = -\frac{x(a^2 - x^2)^{5/2}}{6} + \frac{a^2 x(a^2 - x^2)^{3/2}}{24} + \frac{a^4 x \sqrt{a^2 - x^2}}{16} + \frac{a^6}{16} \sin^{-1} \frac{x}{a}$$

$$17.11.25. \int x^3(a^2 - x^2)^{3/2} dx = \frac{(a^2 - x^2)^{7/2}}{7} - \frac{a^2(a^2 - x^2)^{5/2}}{5}$$

$$17.11.26. \int \frac{(a^2 - x^2)^{3/2}}{x} dx = \frac{(a^2 - x^2)^{3/2}}{3} + a^2 \sqrt{a^2 - x^2} - a^3 \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.11.27. \int \frac{(a^2 - x^2)^{3/2}}{x^2} dx = -\frac{(a^2 - x^2)^{3/2}}{x} - \frac{3x \sqrt{a^2 - x^2}}{2} - \frac{3}{2} a^2 \sin^{-1} \frac{x}{a}$$

$$17.11.28. \int \frac{(a^2 - x^2)^{3/2}}{x^3} dx = -\frac{(a^2 - x^2)^{3/2}}{2x^2} - \frac{3\sqrt{a^2 - x^2}}{2} + \frac{3}{2} a \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

## (12) Integrals Involving $ax^2 + bx + c$

---

$$17.12.1. \int \frac{dx}{ax^2 + bx + c} = \begin{cases} \frac{2}{\sqrt{4ac - b^2}} \tan^{-1} \frac{2ax + b}{\sqrt{4ac - b^2}} \\ \frac{1}{\sqrt{b^2 - 4ac}} \ln \left( \frac{2ax + b - \sqrt{b^2 - 4ac}}{2ax + b + \sqrt{b^2 - 4ac}} \right) \end{cases}$$

If  $b^2 = 4ac$ ,  $ax^2 + bx + c = a(x + b/2a)^2$  and the results 17.1.6 to 17.1.10 and 17.1.14 to 17.1.17 can be used.  
If  $b = 0$  use results on page 75. If  $a$  or  $c = 0$  use results on pages 71–72.

$$17.12.2. \int \frac{x dx}{ax^2 + bx + c} = \frac{1}{2a} \ln(ax^2 + bx + c) - \frac{b}{2a} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.3. \int \frac{x^2 dx}{ax^2 + bx + c} = \frac{x}{a} - \frac{b}{2a^2} \ln(ax^2 + bx + c) + \frac{b^2 - 2ac}{2a^2} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.4. \int \frac{x^m dx}{ax^2 + bx + c} = \frac{x^{m-1}}{(m-1)a} - \frac{c}{a} \int \frac{x^{m-2} dx}{ax^2 + bx + c} - \frac{b}{a} \int \frac{x^{m-1} dx}{ax^2 + bx + c}$$

$$17.12.5. \int \frac{dx}{x(ax^2 + bx + c)} = \frac{1}{2c} \ln \left( \frac{x^2}{ax^2 + bx + c} \right) - \frac{b}{2c} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.6. \int \frac{dx}{x^2(ax^2 + bx + c)} = \frac{b}{2c^2} \ln \left( \frac{ax^2 + bx + c}{x^2} \right) - \frac{1}{cx} + \frac{b^2 - 2ac}{2c^2} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.7. \int \frac{dx}{x^n(ax^2 + bx + c)} = -\frac{1}{(n-1)c x^{n-1}} - \frac{b}{c} \int \frac{dx}{x^{n-1}(ax^2 + bx + c)} - \frac{a}{c} \int \frac{dx}{x^{n-2}(ax^2 + bx + c)}$$

$$17.12.8. \int \frac{dx}{(ax^2 + bx + c)^2} = \frac{2ax + b}{(4ac - b^2)(ax^2 + bx + c)} + \frac{2a}{4ac - b^2} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.9. \int \frac{x dx}{(ax^2 + bx + c)^2} = -\frac{bx + 2c}{(4ac - b^2)(ax^2 + bx + c)} - \frac{b}{4ac - b^2} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.10. \int \frac{x^2 dx}{(ax^2 + bx + c)^2} = \frac{(b^2 - 2ac)x + bc}{a(4ac - b^2)(ax^2 + bx + c)} + \frac{2c}{4ac - b^2} \int \frac{dx}{ax^2 + bx + c}$$

$$17.12.11. \int \frac{x^m dx}{(ax^2 + bx + c)^n} = -\frac{x^{m-1}}{(2n-m-1)a(ax^2 + bx + c)^{n-1}} + \frac{(m-1)c}{(2n-m-1)a} \int \frac{x^{m-2} dx}{(ax^2 + bx + c)^n} \\ - \frac{(n-m)b}{(2n-m-1)a} \int \frac{x^{m-1} dx}{(ax^2 + bx + c)^n}$$

$$17.12.12. \int \frac{x^{2n-1} dx}{(ax^2 + bx + c)^n} = \frac{1}{a} \int \frac{x^{2n-3} dx}{(ax^2 + bx + c)^{n-1}} - \frac{c}{a} \int \frac{x^{2n-3} dx}{(ax^2 + bx + c)^n} - \frac{b}{a} \int \frac{x^{2n-2} dx}{(ax^2 + bx + c)^n}$$

$$17.12.13. \int \frac{dx}{x(ax^2 + bx + c)^2} = \frac{1}{2c(ax^2 + bx + c)} - \frac{b}{2c} \int \frac{dx}{(ax^2 + bx + c)^2} + \frac{1}{c} \int \frac{dx}{x(ax^2 + bx + c)}$$

$$17.12.14. \int \frac{dx}{x^2(ax^2 + bx + c)^2} = -\frac{1}{cx(ax^2 + bx + c)} - \frac{3a}{c} \int \frac{dx}{(ax^2 + bx + c)^2} - \frac{2b}{c} \int \frac{dx}{x(ax^2 + bx + c)^2}$$

$$17.12.15. \int \frac{dx}{x^m(ax^2 + bx + c)^n} = -\frac{1}{(m-1)cx^{m-1}(ax^2 + bx + c)^{n-1}} - \frac{(m+2n-3)a}{(m-1)c} \int \frac{dx}{x^{m-2}(ax^2 + bx + c)^n} \\ - \frac{(m+n-2)b}{(m-1)c} \int \frac{dx}{x^{m-1}(ax^2 + bx + c)^n}$$

### (13) Integrals Involving $\sqrt{ax^2 + bx + c}$

In the following results if  $b^2 = 4ac$ ,  $\sqrt{ax^2 + bx + c} = \sqrt{a}(x + b/2a)$  and the results 17.1 can be used. If  $b = 0$  use the results 17.9. If  $a = 0$  or  $c = 0$  use the results 17.2 and 17.5.

$$17.13.1. \int \frac{dx}{\sqrt{ax^2 + bx + c}} = \begin{cases} \frac{1}{\sqrt{a}} \ln(2\sqrt{a}\sqrt{ax^2 + bx + c} + 2ax + b) \\ -\frac{1}{\sqrt{-a}} \sin^{-1}\left(\frac{2ax + b}{\sqrt{b^2 - 4ac}}\right) \text{ or } \frac{1}{\sqrt{a}} \sinh^{-1}\left(\frac{2ax + b}{\sqrt{4ac - b^2}}\right) \end{cases}$$

$$17.13.2. \int \frac{x dx}{\sqrt{ax^2 + bx + c}} = \frac{\sqrt{ax^2 + bx + c}}{a} - \frac{b}{2a} \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

$$17.13.3. \int \frac{x^2 dx}{\sqrt{ax^2 + bx + c}} = \frac{2ax - 3b}{4a^2} \sqrt{ax^2 + bx + c} + \frac{3b^2 - 4ac}{8a^2} \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

$$17.13.4. \int \frac{dx}{x\sqrt{ax^2 + bx + c}} = \begin{cases} -\frac{1}{\sqrt{c}} \ln\left(\frac{2\sqrt{c}\sqrt{ax^2 + bx + c} + bx + 2c}{x}\right) \\ \frac{1}{\sqrt{-c}} \sin^{-1}\left(\frac{bx + 2c}{|x|\sqrt{b^2 - 4ac}}\right) \text{ or } -\frac{1}{\sqrt{c}} \sinh^{-1}\left(\frac{bx + 2c}{|x|\sqrt{4ac - b^2}}\right) \end{cases}$$

$$17.13.5. \int \frac{dx}{x^2\sqrt{ax^2 + bx + c}} = -\frac{\sqrt{ax^2 + bx + c}}{cx} - \frac{b}{2c} \int \frac{dx}{x\sqrt{ax^2 + bx + c}}$$

$$17.13.6. \int \sqrt{ax^2 + bx + c} dx = \frac{(2ax + b)\sqrt{ax^2 + bx + c}}{4a} + \frac{4ac - b^2}{8a} \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

$$17.13.7. \int x\sqrt{ax^2+bx+c} dx = \frac{(ax^2+bx+c)^{3/2}}{3a} - \frac{b(2ax+b)}{8a^2}\sqrt{ax^2+bx+c} - \frac{b(4ac-b^2)}{16a^2} \int \frac{dx}{\sqrt{ax^2+bx+c}}$$

$$17.13.8. \int x^2\sqrt{ax^2+bx+c} dx = \frac{6ax-5b}{24a^2}(ax^2+bx+c)^{3/2} + \frac{5b^2-4ac}{16a^2} \int \sqrt{ax^2+bx+c} dx$$

$$17.13.9. \int \frac{\sqrt{ax^2+bx+c}}{x} dx = \sqrt{ax^2+bx+c} + \frac{b}{2} \int \frac{dx}{\sqrt{ax^2+bx+c}} + c \int \frac{dx}{x\sqrt{ax^2+bx+c}}$$

$$17.13.10. \int \frac{\sqrt{ax^2+bx+c}}{x^2} dx = -\frac{\sqrt{ax^2+bx+c}}{x} + a \int \frac{dx}{\sqrt{ax^2+bx+c}} + \frac{b}{2} \int \frac{dx}{x\sqrt{ax^2+bx+c}}$$

$$17.13.11. \int \frac{dx}{(ax^2+bx+c)^{3/2}} = \frac{2(2ax+b)}{(4ac-b^2)\sqrt{ax^2+bx+c}}$$

$$17.13.12. \int \frac{x dx}{(ax^2+bx+c)^{3/2}} = \frac{2(bx+2c)}{(b^2-4ac)\sqrt{ax^2+bx+c}}$$

$$17.13.13. \int \frac{x^2 dx}{(ax^2+bx+c)^{3/2}} = \frac{(2b^2-4ac)x+2bc}{a(4ac-b^2)\sqrt{ax^2+bx+c}} + \frac{1}{a} \int \frac{dx}{\sqrt{ax^2+bx+c}}$$

$$17.13.14. \int \frac{dx}{x(ax^2+bx+c)^{3/2}} = \frac{1}{c\sqrt{ax^2+bx+c}} + \frac{1}{c} \int \frac{dx}{x\sqrt{ax^2+bx+c}} - \frac{b}{2c} \int \frac{dx}{(ax^2+bx+c)^{3/2}}$$

$$17.13.15. \int \frac{dx}{x^2(ax^2+bx+c)^{3/2}} = -\frac{ax^2+2bx+c}{c^2x\sqrt{ax^2+bx+c}} + \frac{b^2-2ac}{2c^2} \int \frac{dx}{(ax^2+bx+c)^{3/2}} - \frac{3b}{2c^2} \int \frac{dx}{x\sqrt{ax^2+bx+c}}$$

$$17.13.16. \int (ax^2+bx+c)^{n+1/2} dx = \frac{(2ax+b)(ax^2+bx+c)^{n+1/2}}{4a(n+1)} + \frac{(2n+1)(4ac-b^2)}{8a(n+1)} \int (ax^2+bx+c)^{n-1/2} dx$$

$$17.13.17. \int x(ax^2+bx+c)^{n+1/2} dx = \frac{(ax^2+bx+c)^{n+3/2}}{a(2n+3)} - \frac{b}{2a} \int (ax^2+bx+c)^{n+1/2} dx$$

$$17.13.18. \int \frac{dx}{(ax^2+bx+c)^{n+1/2}} = \frac{2(2ax+b)}{(2n-1)(4ac-b^2)(ax^2+bx+c)^{n-1/2}} + \frac{8a(n-1)}{(2n-1)(4ac-b^2)} \int \frac{dx}{(ax^2+bx+c)^{n-1/2}}$$

$$17.13.19. \int \frac{dx}{x(ax^2+bx+c)^{n+1/2}} = \frac{1}{(2n-1)c(ax^2+bx+c)^{n-1/2}} + \frac{1}{c} \int \frac{dx}{x(ax^2+bx+c)^{n-1/2}} - \frac{b}{2c} \int \frac{dx}{(ax^2+bx+c)^{n+1/2}}$$

**(14) Integrals Involving  $x^3 + a^3$** 

---

Note that for formulas involving  $x^3 - a^3$  replace  $a$  with  $-a$ .

$$17.14.1. \int \frac{dx}{x^3 + a^3} = \frac{1}{6a^2} \ln \left( \frac{(x+a)^2}{x^2 - ax + a^2} \right) + \frac{1}{a^2 \sqrt{3}} \tan^{-1} \frac{2x-a}{a\sqrt{3}}$$

$$17.14.2. \int \frac{x dx}{x^3 + a^3} = \frac{1}{6a} \ln \left( \frac{x^2 - ax + a^2}{(x+a)^2} \right) + \frac{1}{a\sqrt{3}} \tan^{-1} \frac{2x-a}{a\sqrt{3}}$$

$$17.14.3. \int \frac{x^2 dx}{x^3 + a^3} = \frac{1}{3} \ln(x^3 + a^3)$$

$$17.14.4. \int \frac{dx}{x(x^3 + a^3)} = \frac{1}{3a^3} \ln \left( \frac{x^3}{x^3 + a^3} \right)$$

$$17.14.5. \int \frac{dx}{x^2(x^3 + a^3)} = -\frac{1}{a^3 x} - \frac{1}{6a^4} \ln \left( \frac{x^2 - ax + a^2}{(x+a)^2} \right) - \frac{1}{a^4 \sqrt{3}} \tan^{-1} \frac{2x-a}{a\sqrt{3}}$$

$$17.14.6. \int \frac{dx}{(x^3 + a^3)^2} = \frac{x}{3a^3(x^3 + a^3)} + \frac{1}{9a^5} \ln \left( \frac{(x+a)^2}{x^2 - ax + a^2} \right) + \frac{2}{3a^5 \sqrt{3}} \tan^{-1} \frac{2x-a}{a\sqrt{3}}$$

$$17.14.7. \int \frac{x dx}{(x^3 + a^3)^2} = \frac{x^2}{3a^3(x^3 + a^3)} + \frac{1}{18a^4} \ln \left( \frac{x^2 - ax + a^2}{(x+a)^2} \right) + \frac{1}{3a^4 \sqrt{3}} \tan^{-1} \frac{2x-a}{a\sqrt{3}}$$

$$17.14.8. \int \frac{x^2 dx}{(x^3 + a^3)^2} = -\frac{1}{3(x^3 + a^3)}$$

$$17.14.9. \int \frac{dx}{x(x^3 + a^3)^2} = \frac{1}{3a^3(x^3 + a^3)} + \frac{1}{3a^6} \ln \left( \frac{x^3}{x^3 + a^3} \right)$$

$$17.14.10. \int \frac{dx}{x^2(x^3 + a^3)^2} = -\frac{1}{a^6 x} - \frac{x^2}{3a^6(x^3 + a^3)} - \frac{4}{3a^6} \int \frac{x dx}{x^3 + a^3} \quad (\text{See 17.14.2.})$$

$$17.14.11. \int \frac{x^m dx}{x^3 + a^3} = \frac{x^{m-2}}{m-2} - a^3 \int \frac{x^{m-3} dx}{x^3 + a^3}$$

$$17.14.12. \int \frac{dx}{x^n(x^3 + a^3)} = \frac{-1}{a^3(n-1)x^{n-1}} - \frac{1}{a^3} \int \frac{dx}{x^{n-3}(x^3 + a^3)}$$

**(15) Integrals Involving  $x^4 \pm a^4$** 

---

$$17.15.1. \int \frac{dx}{x^4 + a^4} = \frac{1}{4a^3 \sqrt{2}} \ln \left( \frac{x^2 + ax\sqrt{2} + a^2}{x^2 - ax\sqrt{2} + a^2} \right) - \frac{1}{2a^3 \sqrt{2}} \left[ \tan^{-1} \left( 1 - \frac{x\sqrt{2}}{a} \right) - \tan^{-1} \left( 1 + \frac{x\sqrt{2}}{a} \right) \right]$$

$$17.15.2. \int \frac{x dx}{x^4 + a^4} = \frac{1}{2a^2} \tan^{-1} \frac{x^2}{a^2}$$

$$17.15.3. \int \frac{x^2 dx}{x^4 + a^4} = \frac{1}{4a\sqrt{2}} \ln \left( \frac{x^2 - ax\sqrt{2} + a^2}{x^2 + ax\sqrt{2} + a^2} \right) - \frac{1}{2a\sqrt{2}} \left[ \tan^{-1} \left( 1 - \frac{x\sqrt{2}}{a} \right) - \tan^{-1} \left( 1 + \frac{x\sqrt{2}}{a} \right) \right]$$

$$17.15.4. \int \frac{x^3 dx}{x^4 + a^4} = \frac{1}{4} \ln(x^4 + a^4)$$

$$17.15.5. \int \frac{dx}{x(x^4 + a^4)} = \frac{1}{4a^4} \ln\left(\frac{x^4}{x^4 + a^4}\right)$$

$$17.15.6. \int \frac{dx}{x^2(x^4 + a^4)} = -\frac{1}{a^4 x} - \frac{1}{4a^5 \sqrt{2}} \ln\left(\frac{x^2 - ax\sqrt{2} + a^2}{x^2 + ax\sqrt{2} + a^2}\right) \\ + \frac{1}{2a^5 \sqrt{2}} \left[ \tan^{-1}\left(1 - \frac{x\sqrt{2}}{a}\right) - \tan^{-1}\left(1 + \frac{x\sqrt{2}}{a}\right) \right]$$

$$17.15.7. \int \frac{dx}{x^3(x^4 + a^4)} = -\frac{1}{2a^4 x^2} - \frac{1}{2a^6} \tan^{-1}\frac{x^2}{a^2}$$

$$17.15.8. \int \frac{dx}{x^4 - a^4} = \frac{1}{4a^3} \ln\left(\frac{x-a}{x+a}\right) - \frac{1}{2a^3} \tan^{-1}\frac{x}{a}$$

$$17.15.9. \int \frac{x dx}{x^4 - a^4} = \frac{1}{4a^2} \ln\left(\frac{x^2 - a^2}{x^2 + a^2}\right)$$

$$17.15.10. \int \frac{x^2 dx}{x^4 - a^4} = \frac{1}{4a} \ln\left(\frac{x-a}{x+a}\right) + \frac{1}{2a} \tan^{-1}\frac{x}{a}$$

$$17.15.11. \int \frac{x^3 dx}{x^4 - a^4} = \frac{1}{4} \ln(x^4 - a^4)$$

$$17.15.12. \int \frac{dx}{x(x^4 - a^4)} = \frac{1}{4a^4} \ln\left(\frac{x^4 - a^4}{x^4}\right)$$

$$17.15.13. \int \frac{dx}{x^3(x^4 - a^4)} = \frac{1}{a^4 x} + \frac{1}{4a^5} \ln\left(\frac{x-a}{x+a}\right) + \frac{1}{2a^5} \tan^{-1}\frac{x}{a}$$

$$17.15.14. \int \frac{dx}{x^3(x^4 - a^4)} = \frac{1}{2a^4 x^2} + \frac{1}{4a^6} \ln\left(\frac{x^2 - a^2}{x^2 + a^2}\right)$$

### (16) Integrals Involving $x^n \pm a^n$

---

$$17.16.1. \int \frac{dx}{x(x^n + a^n)} = \frac{1}{na^n} \ln\left(\frac{x^n}{x^n + a^n}\right)$$

$$17.16.2. \int \frac{x^{n-1} dx}{x^n + a^n} = \frac{1}{n} \ln(x^n + a^n)$$

$$17.16.3. \int \frac{x^m dx}{(x^n + a^n)^r} = \int \frac{x^{m-n} dx}{(x^n + a^n)^{r-1}} - a^n \int \frac{x^{m-n} dx}{(x^n + a^n)^r}$$

$$17.16.4. \int \frac{dx}{x^m (x^n + a^n)^r} = \frac{1}{a^n} \int \frac{dx}{x^m (x^n + a^n)^{r-1}} - \frac{1}{a^n} \int \frac{dx}{x^{m-n} (x^n + a^n)^r}$$

$$17.16.5. \int \frac{dx}{x\sqrt{x^n + a^n}} = \frac{1}{n\sqrt{a^n}} \ln\left(\frac{\sqrt{x^n + a^n} - \sqrt{a^n}}{\sqrt{x^n + a^n} + \sqrt{a^n}}\right)$$

$$17.16.6. \int \frac{dx}{x(x^n - a^n)} = \frac{1}{na^n} \ln \left( \frac{x^n - a^n}{x^n} \right)$$

$$17.16.7. \int \frac{x^{n-1} dx}{x^n - a^n} = \frac{1}{n} \ln(x^n - a^n)$$

$$17.16.8. \int \frac{x^m dx}{(x^n - a^n)^r} = a^n \int \frac{x^{m-n} dx}{(x^n - a^n)^r} + \int \frac{x^{m-n} dx}{(x^n - a^n)^{r-1}}$$

$$17.16.9. \int \frac{dx}{x^m (x^n - a^n)^r} = \frac{1}{a^n} \int \frac{dx}{x^{m-n} (x^n - a^n)^r} - \frac{1}{a^n} \int \frac{dx}{x^m (x^n - a^n)^{r-1}}$$

$$17.16.10. \int \frac{dx}{x\sqrt{x^n - a^n}} = \frac{2}{n\sqrt{a^n}} \cos^{-1} \sqrt{\frac{a^n}{x^n}}$$

$$17.16.11. \int \frac{x^{p-1} dx}{x^{2m} + a^{2m}} = \frac{1}{ma^{2m-p}} \sum_{k=1}^m \sin \frac{(2k-1)p\pi}{2m} \tan^{-1} \left( \frac{x + a \cos[(2k-1)\pi/2m]}{a \sin[(2k-1)\pi/2m]} \right) \\ - \frac{1}{2ma^{2m-p}} \sum_{k=1}^m \cos \frac{(2k-1)p\pi}{2m} \ln \left( x^2 + 2ax \cos \frac{(2k-1)\pi}{2m} + a^2 \right)$$

where  $0 < p \leq 2m$ .

$$17.16.12. \int \frac{x^{p-1} dx}{x^{2m} - a^{2m}} = \frac{1}{2ma^{2m-p}} \sum_{k=1}^{m-1} \cos \frac{kp\pi}{m} \ln \left( x^2 - 2ax \cos \frac{k\pi}{m} + a^2 \right) \\ - \frac{1}{ma^{2m-p}} \sum_{k=1}^{m-1} \sin \frac{kp\pi}{m} \tan^{-1} \left( \frac{x - a \cos(k\pi/m)}{a \sin(k\pi/m)} \right) \\ + \frac{1}{2ma^{2m-p}} \{ \ln(x-a) + (-1)^p \ln(x+a) \}$$

where  $0 < p \leq 2m$ .

$$17.16.13. \int \frac{x^{p-1} dx}{x^{2m+1} + a^{2m+1}} = \frac{2(-1)^{p-1}}{(2m+1)a^{2m-p+1}} \sum_{k=1}^m \sin \frac{2kp\pi}{2m+1} \tan^{-1} \left( \frac{x + a \cos[2k\pi/(2m+1)]}{a \sin[2k\pi/(2m+1)]} \right) \\ - \frac{(-1)^{p-1}}{(2m+1)a^{2m-p+1}} \sum_{k=1}^m \cos \frac{2kp\pi}{2m+1} \ln \left( x^2 + 2ax \cos \frac{2k\pi}{2m+1} + a^2 \right) \\ + \frac{(-1)^{p-1} \ln(x+a)}{(2m+1)a^{2m-p+1}}$$

where  $0 < p \leq 2m+1$ .

$$17.16.14. \int \frac{x^{p-1} dx}{x^{2m+1} - a^{2m+1}} = \frac{-2}{(2m+1)a^{2m-p+1}} \sum_{k=1}^m \sin \frac{2kp\pi}{2m+1} \tan^{-1} \left( \frac{x - a \cos[2k\pi/(2m+1)]}{a \sin[2k\pi/(2m+1)]} \right) \\ + \frac{1}{(2m+1)a^{2m-p+1}} \sum_{k=1}^m \cos \frac{2kp\pi}{2m+1} \ln \left( x^2 - 2ax \cos \frac{2k\pi}{2m+1} + a^2 \right) \\ + \frac{\ln(x-a)}{(2m+1)a^{2m-p+1}}$$

where  $0 < p \leq 2m+1$ .

**(17) Integrals Involving  $\sin ax$** 

$$17.17.1. \int \sin ax dx = -\frac{\cos ax}{a}$$

$$17.17.2. \int x \sin ax dx = \frac{\sin ax}{a^2} - \frac{x \cos ax}{a}$$

$$17.17.3. \int x^2 \sin ax dx = \frac{2x}{a^2} \sin ax + \left( \frac{2}{a^3} - \frac{x^2}{a} \right) \cos ax$$

$$17.17.4. \int x^3 \sin ax dx = \left( \frac{3x^2}{a^2} - \frac{6}{a^4} \right) \sin ax + \left( \frac{6x}{a^3} - \frac{x^3}{a} \right) \cos ax$$

$$17.17.5. \int \frac{\sin ax}{x} dx = ax - \frac{(ax)^3}{3 \cdot 3!} + \frac{(ax)^5}{5 \cdot 5!} - \dots$$

$$17.17.6. \int \frac{\sin ax}{x^2} dx = -\frac{\sin ax}{x} + a \int \frac{\cos ax}{x} dx \quad (\text{See 17.18.5.})$$

$$17.17.7. \int \frac{dx}{\sin ax} = \frac{1}{a} \ln(\csc ax - \cot ax) = \frac{1}{a} \ln \tan \frac{ax}{2}$$

$$17.17.8. \int \frac{x dx}{\sin ax} = \frac{1}{a^2} \left\{ ax + \frac{(ax)^3}{18} + \frac{7(ax)^5}{1800} + \dots + \frac{2(2^{2n-1}-1)B_n(ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.17.9. \int \sin^2 ax dx = \frac{x}{2} - \frac{\sin 2ax}{4a}$$

$$17.17.10. \int x \sin^2 ax dx = \frac{x^2}{4} - \frac{x \sin 2ax}{4a} - \frac{\cos 2ax}{8a^2}$$

$$17.17.11. \int \sin^3 ax dx = -\frac{\cos ax}{a} + \frac{\cos^3 ax}{3a}$$

$$17.17.12. \int \sin^4 ax dx = \frac{3x}{8} - \frac{\sin 2ax}{4a} + \frac{\sin 4ax}{32a}$$

$$17.17.13. \int \frac{dx}{\sin^2 ax} = -\frac{1}{a} \cot ax$$

$$17.17.14. \int \frac{dx}{\sin^3 ax} = -\frac{\cos ax}{2a \sin^2 ax} + \frac{1}{2a} \ln \tan \frac{ax}{2}$$

$$17.17.15. \int \sin px \sin qx dx = \frac{\sin(p-q)x}{2(p-q)} - \frac{\sin(p+q)x}{2(p+q)} \quad (\text{If } p = \pm q, \text{ see 17.17.9.})$$

$$17.17.16. \int \frac{dx}{1-\sin ax} = \frac{1}{a} \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right)$$

$$17.17.17. \int \frac{x dx}{1-\sin ax} = \frac{x}{a} \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) + \frac{2}{a^2} \ln \sin \left( \frac{\pi}{4} - \frac{ax}{2} \right)$$

$$17.17.18. \int \frac{dx}{1+\sin ax} = -\frac{1}{a} \tan \left( \frac{\pi}{4} - \frac{ax}{2} \right)$$

$$17.17.19. \int \frac{x dx}{1 + \sin ax} = -\frac{x}{a} \tan\left(\frac{\pi}{4} - \frac{ax}{2}\right) + \frac{2}{a^2} \ln \sin\left(\frac{\pi}{4} + \frac{ax}{2}\right)$$

$$17.17.20. \int \frac{dx}{(1 - \sin ax)^2} = \frac{1}{2a} \tan\left(\frac{\pi}{4} + \frac{ax}{2}\right) + \frac{1}{6a} \tan^3\left(\frac{\pi}{4} + \frac{ax}{2}\right)$$

$$17.17.21. \int \frac{dx}{(1 + \sin ax)^2} = -\frac{1}{2a} \tan\left(\frac{\pi}{4} - \frac{ax}{2}\right) - \frac{1}{6a} \tan^3\left(\frac{\pi}{4} - \frac{ax}{2}\right)$$

$$17.17.22. \int \frac{dx}{p + q \sin ax} = \begin{cases} \frac{2}{a\sqrt{p^2 - q^2}} \tan^{-1} \frac{p \tan \frac{1}{2}ax + q}{\sqrt{p^2 - q^2}} \\ \frac{1}{a\sqrt{q^2 - p^2}} \ln \left( \frac{p \tan \frac{1}{2}ax + q - \sqrt{q^2 - p^2}}{p \tan \frac{1}{2}ax + q + \sqrt{q^2 - p^2}} \right) \end{cases}$$

(If  $p = \pm q$ , see 17.17.16 and 17.17.18.)

$$17.17.23. \int \frac{dx}{(p + q \sin ax)^2} = \frac{q \cos ax}{a(p^2 - q^2)(p + q \sin ax)} + \frac{p}{p^2 - q^2} \int \frac{dx}{p + q \sin ax}$$

(If  $p = \pm q$ , see 17.17.20 and 17.17.21.)

$$17.17.24. \int \frac{dx}{p^2 + q^2 \sin^2 ax} = \frac{1}{ap\sqrt{p^2 + q^2}} \tan^{-1} \frac{\sqrt{p^2 + q^2} \tan ax}{p}$$

$$17.17.25. \int \frac{dx}{p^2 - q^2 \sin^2 ax} = \begin{cases} \frac{1}{ap\sqrt{p^2 - q^2}} \tan^{-1} \frac{\sqrt{p^2 - q^2} \tan ax}{p} \\ \frac{1}{2ap\sqrt{q^2 - p^2}} \ln \left( \frac{\sqrt{q^2 - p^2} \tan ax + p}{\sqrt{q^2 - p^2} \tan ax - p} \right) \end{cases}$$

$$17.17.26. \int x^m \sin ax dx = -\frac{x^m \cos ax}{a} + \frac{mx^{m-1} \sin ax}{a^2} - \frac{m(m-1)}{a^2} \int x^{m-2} \sin ax dx$$

$$17.17.27. \int \frac{\sin ax}{x^n} dx = -\frac{\sin ax}{(n-1)x^{n-1}} + \frac{a}{n-1} \int \frac{\cos ax}{x^{n-1}} dx \quad (\text{See 17.18.30.})$$

$$17.17.28. \int \sin^n ax dx = -\frac{\sin^{n-1} ax \cos ax}{an} + \frac{n-1}{n} \int \sin^{n-2} ax dx$$

$$17.17.29. \int \frac{dx}{\sin^n ax} = \frac{-\cos ax}{a(n-1)\sin^{n-1} ax} + \frac{n-2}{n-1} \int \frac{dx}{\sin^{n-2} ax}$$

$$17.17.30. \int \frac{x dx}{\sin^n ax} = \frac{-x \cos ax}{a(n-1)\sin^{n-1} ax} - \frac{1}{a^2(n-1)(n-2)\sin^{n-2} ax} + \frac{n-2}{n-1} \int \frac{x dx}{\sin^{n-2} ax}$$

## (18) Integrals Involving $\cos ax$

---

$$17.18.1. \int \cos ax dx = \frac{\sin ax}{a}$$

$$17.18.2. \int x \cos ax dx = \frac{\cos ax}{a^2} + \frac{x \sin ax}{a}$$

$$17.18.3. \int x^2 \cos ax dx = \frac{2x}{a^2} \cos ax + \left( \frac{x^2}{a} - \frac{2}{a^3} \right) \sin ax$$

$$17.18.4. \int x^3 \cos ax dx = \left( \frac{3x^2}{a^2} - \frac{6}{a^4} \right) \cos ax + \left( \frac{x^3}{a} - \frac{6x}{a^3} \right) \sin ax$$

$$17.18.5. \int \frac{\cos ax}{x} dx = \ln x - \frac{(ax)^2}{2 \cdot 2!} + \frac{(ax)^4}{4 \cdot 4!} - \frac{(ax)^6}{6 \cdot 6!} + \dots$$

$$17.18.6. \int \frac{\cos ax}{x^2} dx = -\frac{\cos ax}{x} - a \int \frac{\sin ax}{x} dx \quad (\text{See 17.17.5.})$$

$$17.18.7. \int \frac{dx}{\cos ax} = \frac{1}{a} \ln(\sec ax + \tan ax) = \frac{1}{a} \ln \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right)$$

$$17.18.8. \int \frac{xdx}{\cos ax} = \frac{1}{a^2} \left\{ \frac{(ax)^2}{2} + \frac{(ax)^4}{8} + \frac{5(ax)^6}{144} + \dots + \frac{E_n(ax)^{2n+2}}{(2n+2)(2n)!} + \dots \right\}$$

$$17.18.9. \int \cos^2 ax dx = \frac{x}{2} + \frac{\sin 2ax}{4a}$$

$$17.18.10. \int x \cos^2 ax dx = \frac{x^2}{4} + \frac{x \sin 2ax}{4a} + \frac{\cos 2ax}{8a^2}$$

$$17.18.11. \int \cos^3 ax dx = \frac{\sin ax}{a} - \frac{\sin^3 ax}{3a}$$

$$17.18.12. \int \cos^4 ax dx = \frac{3x}{8} + \frac{\sin 2ax}{4a} + \frac{\sin 4ax}{32a}$$

$$17.18.13. \int \frac{dx}{\cos^2 ax} = \frac{\tan ax}{a}$$

$$17.18.14. \int \frac{dx}{\cos^3 ax} = \frac{\sin ax}{2a \cos^2 ax} + \frac{1}{2a} \ln \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right)$$

$$17.18.15. \int \cos ax \cos px dx = \frac{\sin(a-p)x}{2(a-p)} + \frac{\sin(a+p)x}{2(a+p)} \quad (\text{If } a = \pm p, \text{ see 17.18.9.})$$

$$17.18.16. \int \frac{dx}{1-\cos ax} = -\frac{1}{a} \cot \frac{ax}{2}$$

$$17.18.17. \int \frac{xdx}{1-\cos ax} = -\frac{x}{a} \cot \frac{ax}{2} + \frac{2}{a^2} \ln \sin \frac{ax}{2}$$

$$17.18.18. \int \frac{dx}{1+\cos ax} = \frac{1}{a} \tan \frac{ax}{2}$$

$$17.18.19. \int \frac{xdx}{1+\cos ax} = \frac{x}{a} \tan \frac{ax}{2} + \frac{2}{a^2} \ln \cos \frac{ax}{2}$$

$$17.18.20. \int \frac{dx}{(1-\cos ax)^2} = -\frac{1}{2a} \cot \frac{ax}{2} - \frac{1}{6a} \cot^3 \frac{ax}{2}$$

$$17.18.21. \int \frac{dx}{(1+\cos ax)^2} = \frac{1}{2a} \tan \frac{ax}{2} + \frac{1}{6a} \tan^3 \frac{ax}{2}$$

$$17.18.22. \int \frac{dx}{p+q \cos ax} = \begin{cases} \frac{2}{a\sqrt{p^2-q^2}} \tan^{-1} \sqrt{(p-q)/(p+q)} \tan \frac{1}{2} ax \\ \frac{1}{a\sqrt{q^2-p^2}} \ln \left( \frac{\tan \frac{1}{2} ax + \sqrt{(q+p)/(q-p)}}{\tan \frac{1}{2} ax - \sqrt{(q+p)/(q-p)}} \right) \end{cases}$$

(If  $p = \pm q$ , see 17.18.16 and 17.18.18.)

17.18.23.  $\int \frac{dx}{(p+q\cos ax)^2} = \frac{q \sin ax}{a(q^2 - p^2)(p+q\cos ax)} - \frac{p}{q^2 - p^2} \int \frac{dx}{p+q\cos ax}$  (If  $p=\pm q$  see 17.18.19 and 17.18.20.)

17.18.24.  $\int \frac{dx}{p^2 + q^2 \cos^2 ax} = \frac{1}{ap\sqrt{p^2 + q^2}} \tan^{-1} \frac{p \tan ax}{\sqrt{p^2 + q^2}}$

17.18.25.  $\int \frac{dx}{p^2 - q^2 \cos^2 ax} = \begin{cases} \frac{1}{ap\sqrt{p^2 - q^2}} \tan^{-1} \frac{p \tan ax}{\sqrt{p^2 - q^2}} \\ \frac{1}{2ap\sqrt{q^2 - p^2}} \ln \left( \frac{p \tan ax - \sqrt{q^2 - p^2}}{p \tan ax + \sqrt{q^2 - p^2}} \right) \end{cases}$

17.18.26.  $\int x^m \cos ax dx = \frac{x^m \sin ax}{a} + \frac{mx^{m-1}}{a^2} \cos ax - \frac{m(m-1)}{a^2} \int x^{m-2} \cos ax dx$

17.18.27.  $\int \frac{\cos ax}{x^n} dx = -\frac{\cos ax}{(n-1)x^{n-1}} - \frac{a}{n-1} \int \frac{\sin ax}{x^{n-1}} dx$  (See 17.17.27.)

17.18.28.  $\int \cos^n ax dx = \frac{\sin ax \cos^{n-1} ax}{an} + \frac{n-1}{n} \int \cos^{n-2} ax dx$

17.18.29.  $\int \frac{dx}{\cos^n ax} = \frac{\sin ax}{a(n-1)\cos^{n-1} ax} + \frac{n-2}{b-1} \int \frac{dx}{\cos^{n-2} ax}$

17.18.30.  $\int \frac{x dx}{\cos^n ax} = \frac{x \sin ax}{a(n-1)\cos^{n-1} ax} - \frac{1}{a^2(n-1)(n-2)\cos^{n-2} ax} + \frac{n-2}{n-1} \int \frac{x dx}{\cos^{n-2} ax}$

### (19) Integrals Involving $\sin ax$ and $\cos ax$

17.19.1.  $\int \sin ax \cos ax dx = \frac{\sin^2 ax}{2a}$

17.19.2.  $\int \sin px \cos qx dx = -\frac{\cos(p-q)x}{2(p-q)} - \frac{\cos(p+q)x}{2(p+q)}$

17.19.3.  $\int \sin^n ax \cos ax dx = \frac{\sin^{n+1} ax}{(n+1)a}$  (If  $n=-1$ , see 17.21.1.)

17.19.4.  $\int \cos^n ax \sin ax dx = -\frac{\cos^{n+1} ax}{(n+1)a}$  (If  $n=-1$ , see 17.20.1.)

17.19.5.  $\int \sin^2 ax \cos^2 ax dx = \frac{x}{8} - \frac{\sin 4ax}{32a}$

17.19.6.  $\int \frac{dx}{\sin ax \cos ax} = \frac{1}{a} \ln \tan ax$

17.19.7.  $\int \frac{dx}{\sin^2 ax \cos ax} = \frac{1}{a} \ln \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) - \frac{1}{a \sin ax}$

17.19.8.  $\int \frac{dx}{\sin ax \cos^2 ax} = \frac{1}{a} \ln \tan \frac{ax}{2} + \frac{1}{a \cos ax}$

17.19.9.  $\int \frac{dx}{\sin^2 ax \cos^2 ax} = -\frac{2 \cot 2ax}{a}$

$$17.19.10. \int \frac{\sin^2 ax}{\cos ax} dx = -\frac{\sin ax}{a} + \frac{1}{a} \ln \tan \left( \frac{ax}{2} + \frac{\pi}{4} \right)$$

$$17.19.11. \int \frac{\cos^2 ax}{\sin ax} dx = \frac{\cos ax}{a} + \frac{1}{a} \ln \tan \frac{ax}{2}$$

$$17.19.12. \int \frac{dx}{\cos ax(1 \pm \sin ax)} = \mp \frac{1}{2a(1 \pm \sin ax)} + \frac{1}{2a} \ln \tan \left( \frac{ax}{2} + \frac{\pi}{4} \right)$$

$$17.19.13. \int \frac{dx}{\sin ax(1 \pm \cos ax)} = \pm \frac{1}{2a(1 \pm \cos ax)} + \frac{1}{2a} \ln \tan \frac{ax}{2}$$

$$17.19.14. \int \frac{dx}{\sin ax \pm \cos ax} = \frac{1}{a\sqrt{2}} \ln \tan \left( \frac{ax}{2} \pm \frac{\pi}{8} \right)$$

$$17.19.15. \int \frac{\sin ax dx}{\sin ax \pm \cos ax} = \frac{x}{2} \mp \frac{1}{2a} \ln(\sin ax \pm \cos ax)$$

$$17.19.16. \int \frac{\cos ax dx}{\sin ax \pm \cos ax} = \pm \frac{x}{2} + \frac{1}{2a} \ln(\sin ax \pm \cos ax)$$

$$17.19.17. \int \frac{\sin ax dx}{p+q \cos ax} = -\frac{1}{aq} \ln(p+q \cos ax)$$

$$17.19.18. \int \frac{\cos ax dx}{p+q \sin ax} = \frac{1}{aq} \ln(p+q \sin ax)$$

$$17.19.19. \int \frac{\sin ax dx}{(p+q \cos ax)^n} = \frac{1}{aq(n-1)(p+q \cos ax)^{n-1}}$$

$$17.19.20. \int \frac{\cos ax dx}{(p+q \sin ax)^n} = \frac{-1}{aq(n-1)(p+q \sin ax)^{n-1}}$$

$$17.19.21. \int \frac{dx}{p \sin ax + q \cos ax} = \frac{1}{a\sqrt{p^2+q^2}} \ln \tan \left( \frac{ax + \tan^{-1}(q/p)}{2} \right)$$

$$17.19.22. \int \frac{dx}{p \sin ax + q \cos ax + r} = \begin{cases} \frac{2}{a\sqrt{r^2-p^2-q^2}} \tan^{-1} \left( \frac{p+(r-q)\tan(ax/2)}{\sqrt{r^2-p^2-q^2}} \right) \\ \frac{1}{a\sqrt{p^2+q^2-r^2}} \ln \left( \frac{p-\sqrt{p^2+q^2-r^2}+(r-q)\tan(ax/2)}{p+\sqrt{p^2+q^2-r^2}+(r-q)\tan(ax/2)} \right) \end{cases}$$

(If  $r = q$  see 17.19.23. If  $r^2 = p^2 + q^2$  see 17.19.24.)

$$17.19.23. \int \frac{dx}{p \sin ax + q(1+\cos ax)} = \frac{1}{ap} \ln \left( q + p \tan \frac{ax}{2} \right)$$

$$17.19.24. \int \frac{dx}{p \sin ax + q \cos ax \pm \sqrt{p^2+q^2}} = \frac{-1}{a\sqrt{p^2+q^2}} \tan \left( \frac{\pi}{4} \mp \frac{ax + \tan^{-1}(q/p)}{2} \right)$$

$$17.19.25. \int \frac{dx}{p^2 \sin^2 ax + q^2 \cos^2 ax} = \frac{1}{apq} \tan^{-1} \left( \frac{p \tan ax}{q} \right)$$

$$17.19.26. \int \frac{dx}{p^2 \sin^2 ax - q^2 \cos^2 ax} = \frac{1}{2apq} \ln \left( \frac{p \tan ax - q}{p \tan ax + q} \right)$$

$$17.19.27. \int \sin^m ax \cos^n ax dx = \begin{cases} -\frac{\sin^{m-1} ax \cos^{n+1} ax}{a(m+n)} + \frac{m-1}{m+n} \int \sin^{m-2} ax \cos^n ax dx \\ \frac{\sin^{m+1} ax \cos^{n-1} ax}{a(m+n)} + \frac{n-1}{m+n} \int \sin^m ax \cos^{n-2} ax dx \end{cases}$$

$$17.19.28. \int \frac{\sin^m ax}{\cos^n ax} dx = \begin{cases} \frac{\sin^{m-1} ax}{a(n-1)\cos^{n-1} ax} - \frac{m-1}{n-1} \int \frac{\sin^{m-2} ax}{\cos^{n-2} ax} dx \\ \frac{\sin^{m+1} ax}{a(n-1)\cos^{n-1} ax} - \frac{m-n+2}{n-1} \int \frac{\sin^m ax}{\cos^{n-2} ax} dx \\ \frac{-\sin^{m-1} ax}{a(m-n)\cos^{n-1} ax} + \frac{m-1}{m-n} \int \frac{\sin^{m-2} ax}{\cos^n ax} dx \end{cases}$$

$$17.19.29. \int \frac{\cos^m ax}{\sin^n ax} dx = \begin{cases} \frac{-\cos^{m-1} ax}{a(n-1)\sin^{n-1} ax} - \frac{m-1}{n-1} \int \frac{\cos^{m-2} ax}{\sin^{n-2} ax} dx \\ \frac{-\cos^{m+1} ax}{a(n-1)\sin^{n-1} ax} - \frac{m-n+2}{n-1} \int \frac{\cos^m ax}{\sin^{n-2} ax} dx \\ \frac{\cos^{m-1} ax}{a(m-n)\sin^{n-1} ax} + \frac{m-1}{m-n} \int \frac{\cos^{m-2} ax}{\sin^n ax} dx \end{cases}$$

$$17.19.30. \int \frac{dx}{\sin^m ax \cos^n ax} = \begin{cases} \frac{1}{a(n-1)\sin^{m-1} ax \cos^{n-1} ax} + \frac{m+n-2}{n-1} \int \frac{dx}{\sin^m ax \cos^{n-2} ax} \\ \frac{-1}{a(m-1)\sin^{m-1} ax \cos^{n-1} ax} + \frac{m+n-2}{m-1} \int \frac{dx}{\sin^{m-2} ax \cos^n ax} \end{cases}$$

## (20) Integrals Involving $\tan ax$

---

$$17.20.1. \int \tan ax dx = -\frac{1}{a} \ln |\cos ax| = \frac{1}{a} \ln |\sec ax|$$

$$17.20.2. \int \tan^2 ax dx = \frac{\tan ax}{a} - x$$

$$17.20.3. \int \tan^3 ax dx = \frac{\tan^2 ax}{2a} + \frac{1}{a} \ln |\cos ax|$$

$$17.20.4. \int \tan^n ax \sec^2 ax dx = \frac{\tan^{n+1} ax}{(n+1)a}$$

$$17.20.5. \int \frac{\sec^2 ax}{\tan ax} dx = \frac{1}{a} \ln |\tan ax|$$

$$17.20.6. \int \frac{dx}{\tan ax} = \frac{1}{a} \ln |\sin ax|$$

$$17.20.7. \int x \tan ax dx = \frac{1}{a^2} \left\{ \frac{(ax)^3}{3} + \frac{(ax)^5}{15} + \frac{2(ax)^7}{105} + \dots + \frac{2^{2n}(2^{2n}-1)B_n(ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.20.8. \int \frac{\tan ax}{x} dx = ax + \frac{(ax)^3}{9} + \frac{2(ax)^5}{75} + \dots + \frac{2^{2n}(2^{2n}-1)B_n(ax)^{2n+1}}{(2n-1)(2n)!} + \dots$$

$$17.20.9. \int x \tan^2 ax dx = \frac{x \tan ax}{a} + \frac{1}{a^2} \ln |\cos ax| - \frac{x^2}{2}$$

$$17.20.10. \int \frac{dx}{p + q \tan ax} = \frac{px}{p^2 + q^2} + \frac{q}{a(p^2 + q^2)} \ln(q \sin ax + p \cos ax)$$

$$17.20.11. \int \tan^n ax dx = \frac{\tan^{n-1} ax}{(n-1)a} - \int \tan^{n-2} ax dx$$

### (21) Integrals Involving $\cot ax$

---

$$17.21.1. \int \cot ax dx = \frac{1}{a} \ln \sin ax$$

$$17.21.2. \int \cot^2 ax dx = -\frac{\cot ax}{a} - x$$

$$17.21.3. \int \cot^3 ax dx = -\frac{\cot^2 ax}{2a} - \frac{1}{a} \ln \sin ax$$

$$17.21.4. \int \cot^n ax \csc^2 ax dx = -\frac{\cot^{n+1} ax}{(n+1)a}$$

$$17.21.5. \int \frac{\csc^2 ax}{\cot ax} dx = -\frac{1}{a} \ln \cot ax$$

$$17.21.6. \int \frac{dx}{\cot ax} = -\frac{1}{a} \ln \cos ax$$

$$17.21.7. \int x \cot ax dx = \frac{1}{a^2} \left\{ ax - \frac{(ax)^3}{9} - \frac{(ax)^5}{225} - \dots - \frac{2^{2n} B_n (ax)^{2n+1}}{(2n+1)!} - \dots \right\}$$

$$17.21.8. \int \frac{\cot ax}{x} dx = -\frac{1}{ax} - \frac{ax}{3} - \frac{(ax)^3}{135} - \dots - \frac{2^{2n} B_n (ax)^{2n-1}}{(2n-1)(2n)!} - \dots$$

$$17.21.9. \int x \cot^2 ax dx = -\frac{x \cot ax}{a} + \frac{1}{a^2} \ln \sin ax - \frac{x^2}{2}$$

$$17.21.10. \int \frac{dx}{p + q \cot ax} = \frac{px}{p^2 + q^2} - \frac{q}{a(p^2 + q^2)} \ln(q \sin ax + p \cos ax)$$

$$17.21.11. \int \cot^n ax dx = -\frac{\cot^{n-1} ax}{(n-1)a} - \int \cot^{n-2} ax dx$$

### (22) Integrals Involving $\sec ax$

---

$$17.22.1. \int \sec ax dx = \frac{1}{a} \ln(\sec ax + \tan ax) = \frac{1}{a} \ln \tan \left( \frac{ax}{2} + \frac{\pi}{4} \right)$$

$$17.22.2. \int \sec^2 ax dx = \frac{\tan ax}{a}$$

$$17.22.3. \int \sec^3 ax dx = \frac{\sec ax \tan ax}{2a} + \frac{1}{2a} \ln(\sec ax + \tan ax)$$

$$17.22.4. \int \sec^n ax \tan ax dx = \frac{\sec^n ax}{na}$$

$$17.22.5. \int \frac{dx}{\sec ax} = \frac{\sin ax}{a}$$

$$17.22.6. \int x \sec ax dx = \frac{1}{a^2} \left\{ \frac{(ax)^2}{2} + \frac{(ax)^4}{8} + \frac{5(ax)^6}{144} + \dots + \frac{E_n(ax)^{2n+2}}{(2n+2)(2n)!} + \dots \right\}$$

$$17.22.7. \int \frac{\sec ax}{x} dx = \ln x + \frac{(ax)^2}{4} + \frac{5(ax)^4}{96} + \frac{61(ax)^6}{4320} + \dots + \frac{E_n(ax)^{2n}}{2n(2n)!} + \dots$$

$$17.22.8. \int x \sec^2 ax dx = \frac{x}{a} \tan ax + \frac{1}{a^2} \ln \cos ax$$

$$17.22.9. \int \frac{dx}{q + p \sec ax} = \frac{x}{q} - \frac{p}{q} \int \frac{dx}{p + q \cos ax}$$

$$17.22.10. \int \sec^n ax dx = \frac{\sec^{n-2} ax \tan ax}{a(n-1)} + \frac{n-2}{n-1} \int \sec^{n-2} ax dx$$

### (23) Integrals Involving $\csc ax$

$$17.23.1. \int \csc ax dx = \frac{1}{a} \ln(\csc ax - \cot ax) = \frac{1}{a} \ln \tan \frac{ax}{2}$$

$$17.23.2. \int \csc^2 ax dx = -\frac{\cot ax}{a}$$

$$17.23.3. \int \csc^3 ax dx = -\frac{\csc ax \cot ax}{2a} + \frac{1}{2a} \ln \tan \frac{ax}{2}$$

$$17.23.4. \int \csc^n ax \cot ax dx = -\frac{\csc^n ax}{na}$$

$$17.23.5. \int \frac{dx}{\csc ax} = -\frac{\cos ax}{a}$$

$$17.23.6. \int x \csc ax dx = \frac{1}{a^2} \left\{ ax + \frac{(ax)^3}{18} + \frac{7(ax)^5}{1800} + \dots + \frac{2(2^{2n-1}-1)B_n(ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.23.7. \int \frac{\csc ax}{x} dx = -\frac{1}{ax} + \frac{ax}{6} + \frac{7(ax)^3}{1080} + \dots + \frac{2(2^{2n-1}-1)B_n(ax)^{2n-1}}{(2n-1)(2n)!} + \dots$$

$$17.23.8. \int x \csc^2 ax dx = -\frac{x \cot ax}{a} + \frac{1}{a^2} \ln \sin ax$$

$$17.23.9. \int \frac{dx}{q + p \csc ax} = \frac{x}{q} - \frac{p}{q} \int \frac{dx}{p + q \sin ax} \quad (\text{See 17.17.22.})$$

$$17.23.10. \int \csc^n ax dx = -\frac{\csc^{n-2} ax \cot ax}{a(n-1)} + \frac{n-2}{n-1} \int \csc^{n-2} ax dx$$

**(24) Integrals Involving Inverse Trigonometric Functions**

$$17.24.1. \int \sin^{-1} \frac{x}{a} dx = x \sin^{-1} \frac{x}{a} + \sqrt{a^2 - x^2}$$

$$17.24.2. \int x \sin^{-1} \frac{x}{a} dx = \left( \frac{x^2}{2} - \frac{a^2}{4} \right) \sin^{-1} \frac{x}{a} + \frac{x\sqrt{a^2 - x^2}}{4}$$

$$17.24.3. \int x^2 \sin^{-1} \frac{x}{a} dx = \frac{x^3}{3} \sin^{-1} \frac{x}{a} + \frac{(x^2 + 2a^2)\sqrt{a^2 - x^2}}{9}$$

$$17.24.4. \int \frac{\sin^{-1}(x/a)}{x} dx = \frac{x}{a} + \frac{(x/a)^3}{2 \cdot 3 \cdot 3} + \frac{1 \cdot 3 \cdot 5(x/a)^5}{2 \cdot 4 \cdot 5 \cdot 5} + \frac{1 \cdot 3 \cdot 5 \cdot 7(x/a)^7}{2 \cdot 4 \cdot 6 \cdot 7 \cdot 7} + \dots$$

$$17.24.5. \int \frac{\sin^{-1}(x/a)}{x^2} dx = -\frac{\sin^{-1}(x/a)}{x} - \frac{1}{a} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.24.6. \int \left( \sin^{-1} \frac{x}{a} \right)^2 dx = x \left( \sin^{-1} \frac{x}{a} \right)^2 - 2x + 2\sqrt{a^2 - x^2} \sin^{-1} \frac{x}{a}$$

$$17.24.7. \int \cos^{-1} \frac{x}{a} dx = x \cos^{-1} \frac{x}{a} - \sqrt{a^2 - x^2}$$

$$17.24.8. \int x \cos^{-1} \frac{x}{a} dx = \left( \frac{x^2}{2} - \frac{a^2}{4} \right) \cos^{-1} \frac{x}{a} - \frac{x\sqrt{a^2 - x^2}}{4}$$

$$17.24.9. \int x^2 \cos^{-1} \frac{x}{a} dx = \frac{x^3}{3} \cos^{-1} \frac{x}{a} - \frac{(x^2 + 2a^2)\sqrt{a^2 - x^2}}{9}$$

$$17.24.10. \int \frac{\cos^{-1}(x/a)}{x} dx = \frac{\pi}{2} \ln x - \int \frac{\sin^{-1}(x/a)}{x} dx \quad (\text{See 17.24.4.})$$

$$17.24.11. \int \frac{\cos^{-1}(x/a)}{x^2} dx = -\frac{\cos^{-1}(x/a)}{x} + \frac{1}{a} \ln \left( \frac{a + \sqrt{a^2 - x^2}}{x} \right)$$

$$17.24.12. \int \left( \cos^{-1} \frac{x}{a} \right)^2 dx = x \left( \cos^{-1} \frac{x}{a} \right)^2 - 2x - 2\sqrt{a^2 - x^2} \cos^{-1} \frac{x}{a}$$

$$17.24.13. \int \tan^{-1} \frac{x}{a} dx = x \tan^{-1} \frac{x}{a} - \frac{a}{2} \ln(x^2 + a^2)$$

$$17.24.14. \int x \tan^{-1} \frac{x}{a} dx = \frac{1}{2} (x^2 + a^2) \tan^{-1} \frac{x}{a} - \frac{ax}{2}$$

$$17.24.15. \int x^2 \tan^{-1} \frac{x}{a} dx = \frac{x^3}{3} \tan^{-1} \frac{x}{a} - \frac{ax^2}{6} + \frac{a^3}{6} \ln(x^2 + a^2)$$

$$17.24.16. \int \frac{\tan^{-1}(x/a)}{x} dx = \frac{x}{a} - \frac{(x/a)^3}{3^2} + \frac{(x/a)^5}{5^2} - \frac{(x/a)^7}{7^2} + \dots$$

$$17.24.17. \int \frac{\tan^{-1}(x/a)}{x^2} dx = -\frac{1}{x} \tan^{-1} \frac{x}{a} - \frac{1}{2a} \ln \left( \frac{x^2 + a^2}{x^2} \right)$$

$$17.24.18. \int \cot^{-1} \frac{x}{a} dx = x \cot^{-1} \frac{x}{a} + \frac{a}{2} \ln(x^2 + a^2)$$

$$17.24.19. \int x \cot^{-1} \frac{x}{a} dx = \frac{1}{2}(x^2 + a^2) \cot^{-1} \frac{x}{a} + \frac{ax}{2}$$

$$17.24.20. \int x^2 \cot^{-1} \frac{x}{a} dx = \frac{x^3}{3} \cot^{-1} \frac{x}{a} + \frac{ax^2}{6} - \frac{a^3}{6} \ln(x^2 + a^2)$$

$$17.24.21. \int \frac{\cot^{-1}(x/a)}{x} dx = \frac{\pi}{2} \ln x - \int \frac{\tan^{-1}(x/a)}{x} dx \quad (\text{See 17.24.16.})$$

$$17.24.22. \int \frac{\cot^{-1}(x/a)}{x^2} dx = \frac{\cot^{-1}(x/a)}{x} + \frac{1}{2a} \ln \left( \frac{x^2 + a^2}{x^2} \right)$$

$$17.24.23. \int \sec^{-1} \frac{x}{a} dx = \begin{cases} x \sec^{-1} \frac{x}{a} - a \ln(x + \sqrt{x^2 - a^2}) & 0 < \sec^{-1} \frac{x}{a} < \frac{\pi}{2} \\ x \sec^{-1} \frac{x}{a} + a \ln(x + \sqrt{x^2 - a^2}) & \frac{\pi}{2} < \sec^{-1} \frac{x}{a} < \pi \end{cases}$$

$$17.24.24. \int x \sec^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^2}{2} \sec^{-1} \frac{x}{a} - \frac{a\sqrt{x^2 - a^2}}{2} & 0 < \sec^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^2}{2} \sec^{-1} \frac{x}{a} + \frac{a\sqrt{x^2 - a^2}}{2} & \frac{\pi}{2} < \sec^{-1} \frac{x}{a} < \pi \end{cases}$$

$$17.24.25. \int x^2 \sec^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^3}{3} \sec^{-1} \frac{x}{a} - \frac{ax\sqrt{x^2 - a^2}}{6} - \frac{a^3}{6} \ln(x + \sqrt{x^2 - a^2}) & 0 < \sec^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^3}{3} \sec^{-1} \frac{x}{a} + \frac{ax\sqrt{x^2 - a^2}}{6} + \frac{a^3}{6} \ln(x + \sqrt{x^2 - a^2}) & \frac{\pi}{2} < \sec^{-1} \frac{x}{a} < \pi \end{cases}$$

$$17.24.26. \int \frac{\sec^{-1}(x/a)}{x} dx = \frac{\pi}{2} \ln x + \frac{a}{x} + \frac{(a/x)^3}{2 \cdot 3 \cdot 3} + \frac{1 \cdot 3 \cdot 5 (a/x)^5}{2 \cdot 4 \cdot 5 \cdot 5} + \frac{1 \cdot 3 \cdot 5 (a/x)^7}{2 \cdot 4 \cdot 6 \cdot 7 \cdot 7} + \dots$$

$$17.24.27. \int \frac{\sec^{-1}(x/a)}{x^2} dx = \begin{cases} -\frac{\sec^{-1}(x/a)}{x} + \frac{\sqrt{x^2 - a^2}}{ax} & 0 < \sec^{-1} \frac{x}{a} < \frac{\pi}{2} \\ -\frac{\sec^{-1}(x/a)}{x} - \frac{\sqrt{x^2 - a^2}}{ax} & \frac{\pi}{2} < \sec^{-1} \frac{x}{a} < \pi \end{cases}$$

$$17.24.28. \int \csc^{-1} \frac{x}{a} dx = \begin{cases} x \csc^{-1} \frac{x}{a} + a \ln(x + \sqrt{x^2 - a^2}) & 0 < \csc^{-1} \frac{x}{a} < \frac{\pi}{2} \\ x \csc^{-1} \frac{x}{a} - a \ln(x + \sqrt{x^2 - a^2}) & -\frac{\pi}{2} < \csc^{-1} \frac{x}{a} < 0 \end{cases}$$

$$17.24.29. \int x \csc^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^2}{2} \csc^{-1} \frac{x}{a} + \frac{a\sqrt{x^2 - a^2}}{2} & 0 < \csc^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^2}{2} \csc^{-1} \frac{x}{a} - \frac{a\sqrt{x^2 - a^2}}{2} - \frac{\pi}{2} & -\frac{\pi}{2} < \csc^{-1} \frac{x}{a} < 0 \end{cases}$$

$$17.24.30. \int x^2 \csc^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^3}{3} \csc^{-1} \frac{x}{a} + \frac{ax\sqrt{x^2 - a^2}}{6} + \frac{a^3}{6} \ln(x + \sqrt{x^2 - a^2}) & 0 < \csc^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^3}{3} \csc^{-1} \frac{x}{a} - \frac{ax\sqrt{x^2 - a^2}}{6} - \frac{a^3}{6} \ln(x + \sqrt{x^2 - a^2}) & -\frac{\pi}{2} < \csc^{-1} \frac{x}{a} < 0 \end{cases}$$

$$17.24.31. \int \frac{\csc^{-1}(x/a)}{x} dx = -\left( \frac{a}{x} + \frac{(ax)^3}{2 \cdot 3 \cdot 3} + \frac{1 \cdot 3(ax)^5}{2 \cdot 4 \cdot 5 \cdot 5} + \frac{1 \cdot 3 \cdot 5(ax)^7}{2 \cdot 4 \cdot 6 \cdot 7 \cdot 7} + \dots \right)$$

$$17.24.32. \int \frac{\csc^{-1}(x/a)}{x^2} dx = \begin{cases} \frac{\csc^{-1}(x/a)}{x} - \frac{\sqrt{x^2 - a^2}}{ax} & 0 < \csc^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{\csc^{-1}(x/a)}{x} + \frac{\sqrt{x^2 - a^2}}{ax} & -\frac{\pi}{2} < \csc^{-1} \frac{x}{a} < 0 \end{cases}$$

$$17.24.33. \int x^m \sin^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \sin^{-1} \frac{x}{a} - \frac{1}{m+1} \int \frac{x^{m+1}}{\sqrt{a^2 - x^2}} dx$$

$$17.24.34. \int x^m \cos^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \cos^{-1} \frac{x}{a} + \frac{1}{m+1} \int \frac{x^{m+1}}{\sqrt{a^2 - x^2}} dx$$

$$17.24.35. \int x^m \tan^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \tan^{-1} \frac{x}{a} - \frac{a}{m+1} \int \frac{x^{m+1}}{x^2 + a^2} dx$$

$$17.24.36. \int x^m \cot^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \cot^{-1} \frac{x}{a} + \frac{a}{m+1} \int \frac{x^{m+1}}{x^2 + a^2} dx$$

$$17.24.37. \int x^m \sec^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^{m+1} \sec^{-1}(x/a)}{m+1} - \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{x^2 - a^2}} & 0 < \sec^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^{m+1} \sec^{-1}(x/a)}{m+1} + \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{x^2 - a^2}} & \frac{\pi}{2} < \sec^{-1} \frac{x}{a} < \pi \end{cases}$$

$$17.24.38. \int x^m \csc^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^{m+1} \sec^{-1}(x/a)}{m+1} + \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{x^2 - a^2}} & 0 < \csc^{-1} \frac{x}{a} < \frac{\pi}{2} \\ \frac{x^{m+1} \csc^{-1}(x/a)}{m+1} - \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{x^2 - a^2}} & -\frac{\pi}{2} < \csc^{-1} \frac{x}{a} < 0 \end{cases}$$

### (25) Integrals Involving $e^{ax}$

---

$$17.25.1. \int e^{ax} dx = \frac{e^{ax}}{a}$$

$$17.25.2. \int x e^{ax} dx = \frac{e^{ax}}{a} \left( x - \frac{1}{a} \right)$$

$$17.25.3. \int x^2 e^{ax} dx = \frac{e^{ax}}{a} \left( x^2 - \frac{2x}{a} + \frac{2}{a^2} \right)$$

$$\begin{aligned} 17.25.4. \int x^n e^{ax} dx &= \frac{x^n e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} dx \\ &= \frac{e^{ax}}{a} \left( x^n - \frac{nx^{n-1}}{a} + \frac{n(n-1)x^{n-2}}{a^2} - \dots - \frac{(-1)^n n!}{a^n} \right) \text{ if } n = \text{positive integer} \end{aligned}$$

$$17.25.5. \int \frac{e^{ax}}{x} dx = \ln x + \frac{ax}{1 \cdot 1!} + \frac{(ax)^2}{2 \cdot 2!} + \frac{(ax)^3}{3 \cdot 3!} + \dots$$

$$17.25.6. \int \frac{e^{ax}}{x^n} dx = \frac{-e^{ax}}{(n-1)x^{n-1}} + \frac{a}{n-1} \int \frac{e^{ax}}{x^{n-1}} dx$$

$$17.25.7. \int \frac{dx}{p+qe^{ax}} = \frac{x}{p} - \frac{1}{ap} \ln(p+qe^{ax})$$

$$17.25.8. \int \frac{dx}{(p+qe^{ax})^2} = \frac{x}{p^2} + \frac{1}{ap(p+qe^{ax})} - \frac{1}{ap^2} \ln(p+qe^{ax})$$

$$17.25.9. \int \frac{dx}{pe^{ax}+qe^{-ax}} = \begin{cases} \frac{1}{a\sqrt{pq}} \tan^{-1}\left(\sqrt{\frac{p}{q}}e^{ax}\right) \\ \frac{1}{2a\sqrt{-pq}} \ln\left(\frac{e^{ax}-\sqrt{-q/p}}{e^{ax}+\sqrt{-q/p}}\right) \end{cases}$$

$$17.25.10. \int e^{ax} \sin bx dx = \frac{e^{ax}(a \sin bx - b \cos bx)}{a^2 + b^2}$$

$$17.25.11. \int e^{ax} \cos bx dx = \frac{e^{ax}(a \cos bx + b \sin bx)}{a^2 + b^2}$$

$$17.25.12. \int xe^{ax} \sin bx dx = \frac{xe^{ax}(a \sin bx - b \cos bx)}{a^2 + b^2} - \frac{e^{ax}\{(a^2 - b^2)\sin bx - 2ab \cos bx\}}{(a^2 + b^2)^2}$$

$$17.25.13. \int xe^{ax} \cos bx dx = \frac{xe^{ax}(a \cos bx + b \sin bx)}{a^2 + b^2} - \frac{e^{ax}\{(a^2 - b^2)\cos bx + 2ab \sin bx\}}{(a^2 + b^2)^2}$$

$$17.25.14. \int e^{ax} \ln x dx = \frac{e^{ax} \ln x}{a} - \frac{1}{a} \int \frac{e^{ax}}{x} dx$$

$$17.25.15. \int e^{ax} \sin^n bx dx = \frac{e^{ax} \sin^{n-1} bx}{a^2 + n^2 b^2} (a \sin bx - nb \cos bx) + \frac{n(n-1)b^2}{a^2 + n^2 b^2} \int e^{ax} \sin^{n-2} bx dx$$

$$17.25.16. \int e^{ax} \cos^n bx dx = \frac{e^{ax} \cos^{n-1} bx}{a^2 + n^2 b^2} (a \cos bx + nb \sin bx) + \frac{n(n-1)b^2}{a^2 + n^2 b^2} \int e^{ax} \cos^{n-2} bx dx$$

## (26) Integrals Involving $\ln x$

---

$$17.26.1. \int \ln x dx = x \ln x - x$$

$$17.26.2. \int x \ln x dx = \frac{x^2}{2} \left( \ln x - \frac{1}{2} \right)$$

$$17.26.3. \int x^m \ln x dx = \frac{x^{m+1}}{m+1} \left( \ln x - \frac{1}{m+1} \right) \quad (\text{If } m = -1, \text{ see 17.26.4.})$$

$$17.26.4. \int \frac{\ln x}{x} dx = \frac{1}{2} \ln^2 x$$

$$17.26.5. \int \frac{\ln x}{x^2} dx = -\frac{\ln x}{x} - \frac{1}{x}$$

$$17.26.6. \int \ln^2 x dx = x \ln^2 x - 2x \ln x + 2x$$

$$17.26.7. \int \frac{\ln^n x dx}{x} = \frac{\ln^{n+1} x}{n+1} \quad (\text{If } n = -1, \text{ see 17.26.8.})$$

$$17.26.8. \int \frac{dx}{x \ln x} = \ln(\ln x)$$

$$17.26.9. \quad \int \frac{dx}{\ln x} = \ln(\ln x) + \ln x + \frac{\ln^2 x}{2 \cdot 2!} + \frac{\ln^3 x}{3 \cdot 3!} + \dots$$

$$17.26.10. \quad \int \frac{x^m dx}{\ln x} = \ln(\ln x) + (m+1)\ln x + \frac{(m+1)^2 \ln^2 x}{2 \cdot 2!} + \frac{(m+1)^3 \ln^3 x}{3 \cdot 3!} + \dots$$

$$17.26.11. \quad \int \ln^n x dx = x \ln^n x - n \int \ln^{n-1} x dx$$

$$17.26.12. \quad \int x^m \ln^n x dx = \frac{x^{m+1} \ln^n x}{m+1} - \frac{n}{m+1} \int x^m \ln^{n-1} x dx$$

If  $m = -1$ , see 17.26.7.

$$17.26.13. \quad \int \ln(x^2 + a^2) dx = x \ln(x^2 + a^2) - 2x + 2a \tan^{-1} \frac{x}{a}$$

$$17.26.14. \quad \int \ln(x^2 - a^2) dx = x \ln(x^2 - a^2) - 2x + a \ln \left( \frac{x+a}{x-a} \right)$$

$$17.26.15. \quad \int x^m \ln(x^2 \pm a^2) dx = \frac{x^{m+1} \ln(x^2 \pm a^2)}{m+1} - \frac{2}{m+1} \int \frac{x^{m+2}}{x^2 \pm a^2} dx$$

## (27) Integrals Involving $\sinh ax$

---

$$17.27.1. \quad \int \sinh ax dx = \frac{\cosh ax}{a}$$

$$17.27.2. \quad \int x \sinh ax dx = \frac{x \cosh ax}{a} - \frac{\sinh ax}{a^2}$$

$$17.27.3. \quad \int x^2 \sinh ax dx = \left( \frac{x^2}{a} + \frac{2}{a^3} \right) \cosh ax - \frac{2x}{a^2} \sinh ax$$

$$17.27.4. \quad \int \frac{\sinh ax}{x} dx = ax + \frac{(ax)^3}{3 \cdot 3!} + \frac{(ax)^5}{5 \cdot 5!} + \dots$$

$$17.27.5. \quad \int \frac{\sinh ax}{x^2} dx = -\frac{\sinh ax}{x} + a \int \frac{\cosh ax}{x} dx \quad (\text{See 17.28.4.})$$

$$17.27.6. \quad \int \frac{dx}{\sinh ax} = \frac{1}{a} \ln \tanh \frac{ax}{2}$$

$$17.27.7. \quad \int \frac{x dx}{\sinh ax} = \frac{1}{a^2} \left\{ ax - \frac{(ax)^3}{18} + \frac{7(ax)^5}{1800} - \dots + \frac{2(-1)^n (2^{2n}-1) B_n (ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.27.8. \quad \int \sinh^2 ax dx = \frac{\sinh ax \cosh ax}{2a} - \frac{x}{2}$$

$$17.27.9. \quad \int x \sinh^2 ax dx = \frac{x \sinh 2ax}{4a} - \frac{\cosh 2ax}{8a^2} - \frac{x^2}{4}$$

$$17.27.10. \quad \int \frac{dx}{\sinh^2 ax} = -\frac{\coth ax}{a}$$

$$17.27.11. \quad \int \sinh ax \sinh px dx = \frac{\sinh(a+p)x}{2(a+p)} - \frac{\sinh(a-p)x}{2(a-p)}$$

For  $a = \pm p$  see 17.27.8.

$$17.27.12. \quad \int x^m \sinh ax dx = \frac{x^m \cosh ax}{a} - \frac{m}{a} \int x^{m-1} \cosh ax dx \quad (\text{See 17.28.12.})$$

$$17.27.13. \quad \int \sinh^n ax dx = \frac{\sinh^{n-1} ax \cosh ax}{an} - \frac{n-1}{n} \int \sinh^{n-2} ax dx$$

$$17.27.14. \quad \int \frac{\sinh ax}{x^n} dx = \frac{-\sinh ax}{(n-1)x^{n-1}} + \frac{a}{n-1} \int \frac{\cosh ax}{x^{n-1}} dx \quad (\text{See 17.28.14.})$$

$$17.27.15. \quad \int \frac{dx}{\sinh^n ax} = \frac{-\cosh ax}{a(n-1)\sinh^{n-1} ax} - \frac{n-2}{n-1} \int \frac{dx}{\sinh^{n-2} ax}$$

$$17.27.16. \quad \int \frac{x dx}{\sinh^n ax} = \frac{-x \cosh ax}{a(n-1)\sinh^{n-1} ax} - \frac{1}{a^2(n-1)(n-2)\sinh^{n-2} ax} - \frac{n-2}{n-1} \int \frac{x dx}{\sinh^{n-2} ax}$$

## (28) Integrals Involving $\cosh ax$

---

$$17.28.1. \quad \int \cosh ax dx = \frac{\sinh ax}{a}$$

$$17.28.2. \quad \int x \cosh ax dx = \frac{x \sinh ax}{a} - \frac{\cosh ax}{a^2}$$

$$17.28.3. \quad \int x^2 \cosh ax dx = -\frac{2x \cosh ax}{a^2} + \left( \frac{x^2}{a} + \frac{2}{a^3} \right) \sinh ax$$

$$17.28.4. \quad \int \frac{\cosh ax}{x} dx = \ln x + \frac{(ax)^2}{2 \cdot 2!} + \frac{(ax)^4}{4 \cdot 4!} + \frac{(ax)^6}{6 \cdot 6!} + \dots$$

$$17.28.5. \quad \int \frac{\cosh ax}{x^2} dx = -\frac{\cosh ax}{x} + a \int \frac{\sinh ax}{x} dx \quad (\text{See 17.27.4.})$$

$$17.28.6. \quad \int \frac{dx}{\cosh ax} = \frac{2}{a} \tan^{-1} e^{ax}$$

$$17.28.7. \quad \int \frac{x dx}{\cosh ax} = \frac{1}{a^2} \left\{ \frac{(ax)^2}{2} - \frac{(ax)^4}{8} + \frac{5(ax)^6}{144} + \dots + \frac{(-1)^n E_n(ax)^{2n+2}}{(2n+2)(2n)!} + \dots \right\}$$

$$17.28.8. \quad \int \cosh^2 ax dx = \frac{x}{2} + \frac{\sinh ax \cosh ax}{2a}$$

$$17.28.9. \quad \int x \cosh^2 ax dx = \frac{x^2}{4} + \frac{x \sinh 2ax}{4a} - \frac{\cosh 2ax}{8a^2}$$

$$17.28.10. \quad \int \frac{dx}{\cosh^2 ax} = \frac{\tanh ax}{a}$$

$$17.28.11. \quad \int \cosh ax \cosh px dx = \frac{\sinh(a-p)x}{2(a-p)} + \frac{\sinh(a+p)x}{2(a+p)}$$

$$17.28.12. \quad \int x^m \cosh ax dx = \frac{x^m \sinh ax}{a} - \frac{m}{a} \int x^{m-1} \sinh ax dx \quad (\text{See 17.27.12.})$$

$$17.28.13. \quad \int \cosh^n ax dx = \frac{\cosh^{n-1} ax \sinh ax}{an} + \frac{n-1}{n} \int \cosh^{n-2} ax dx$$

$$17.28.14. \quad \int \frac{\cosh ax}{x^n} dx = \frac{-\cosh ax}{(n-1)x^{n-1}} + \frac{a}{n-1} \int \frac{\sinh ax}{x^{n-1}} dx \quad (\text{See 17.27.14.})$$

$$17.28.15. \quad \int \frac{dx}{\cosh^n ax} = \frac{\sinh ax}{a(n-1) \cosh^{n-1} ax} + \frac{n-2}{n-1} \int \frac{dx}{\cosh^{n-2} ax}$$

$$17.28.16. \quad \int \frac{x dx}{\cosh^n ax} = \frac{x \sinh ax}{a(n-1) \cosh^{n-1} ax} + \frac{1}{(n-1)(n-2)a^2 \cosh^{n-2} ax} + \frac{n-2}{n-1} \int \frac{x dx}{\cosh^{n-2} ax}$$

### (29) Integrals Involving $\sinh ax$ and $\cosh ax$

---

$$17.29.1. \quad \int \sinh ax \cosh ax dx = \frac{\sinh^2 ax}{2a}$$

$$17.29.2. \quad \int \sinh px \cosh qx dx = \frac{\cosh(p+q)x}{2(p+q)} + \frac{\cosh(p-q)x}{2(p-q)}$$

$$17.29.3. \quad \int \sinh^2 ax \cosh^2 ax dx = \frac{\sinh 4ax}{32a} - \frac{x}{8}$$

$$17.29.4. \quad \int \frac{dx}{\sinh ax \cosh ax} = \frac{1}{a} \ln \tanh ax$$

$$17.29.5. \quad \int \frac{dx}{\sinh^2 ax \cosh^2 ax} = -\frac{2 \coth 2ax}{a}$$

$$17.29.6. \quad \int \frac{\sinh^2 ax}{\cosh ax} dx = \frac{\sinh ax}{a} - \frac{1}{a} \tan^{-1} \sinh ax$$

$$17.29.7. \quad \int \frac{\cosh^2 ax}{\sinh ax} dx = \frac{\cosh ax}{a} + \frac{1}{a} \ln \tanh \frac{ax}{2}$$

**(30) Integrals Involving  $\tanh ax$** 

---

$$17.30.1. \int \tanh ax dx = \frac{1}{a} \ln \cosh ax$$

$$17.30.2. \int \tanh^2 ax dx = x - \frac{\tanh ax}{a}$$

$$17.30.3. \int \tanh^3 ax dx = \frac{1}{a} \ln \cosh ax - \frac{\tanh^2 ax}{2a}$$

$$17.30.4. \int x \tanh ax dx = \frac{1}{a^2} \left\{ \frac{(ax)^3}{3} - \frac{(ax)^5}{15} + \frac{2(ax)^7}{105} - \dots - \frac{(-1)^{n-1} 2^{2n} (2^{2n}-1) B_n (ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.30.5. \int x \tanh^2 ax dx = \frac{x^2}{2} - \frac{x \tanh ax}{a} + \frac{1}{a^2} \ln \cosh ax$$

$$17.30.6. \int \frac{\tanh ax}{x} dx = ax - \frac{(ax)^3}{9} + \frac{2(ax)^5}{75} - \dots - \frac{(-1)^{n-1} 2^{2n} (2^{2n}-1) B_n (ax)^{2n-1}}{(2n-1)(2n)!} + \dots$$

$$17.30.7. \int \frac{dx}{p+q \tanh ax} = \frac{px}{p^2-q^2} - \frac{q}{a(p^2-q^2)} \ln(q \sinh ax + p \cosh ax)$$

$$17.30.8. \int \tanh^n ax dx = \frac{-\tanh^{n-1} ax}{a(a-1)} + \int \tanh^{n-2} ax dx$$

**(31) Integrals Involving  $\coth ax$** 

---

$$17.31.1. \int \coth ax dx = \frac{1}{a} \ln \sinh ax$$

$$17.31.2. \int \coth^2 ax dx = x - \frac{\coth ax}{a}$$

$$17.31.3. \int \coth^3 ax dx = \frac{1}{a} \ln \sinh ax - \frac{\coth^2 ax}{2a}$$

$$17.31.4. \int x \coth ax dx = \frac{1}{a^2} \left\{ ax + \frac{(ax)^3}{9} - \frac{(ax)^5}{225} + \dots - \frac{(-1)^{n-1} 2^{2n} B_n (ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.31.5. \int x \coth^2 ax dx = \frac{x^2}{2} - \frac{x \coth ax}{a} + \frac{1}{a^2} \ln \sinh ax$$

$$17.31.6. \int \frac{\coth ax}{x} dx = -\frac{1}{ax} + \frac{ax}{3} - \frac{(ax)^3}{135} + \dots - \frac{(-1)^n 2^{2n} B_n (ax)^{2n-1}}{(2n-1)(2n)!} + \dots$$

$$17.31.7. \int \frac{dx}{p+q \coth ax} = \frac{px}{p^2-q^2} - \frac{q}{a(p^2-q^2)} \ln(p \sinh ax + q \cosh ax)$$

$$17.31.8. \int \coth^n ax dx = -\frac{\coth^{n-1} ax}{a(n-1)} + \int \coth^{n-2} ax dx$$

**(32) Integrals Involving  $\operatorname{sech} ax$** 

---

$$17.32.1. \int \operatorname{sech} ax dx = \frac{2}{a} \tan^{-1} e^{ax}$$

$$17.32.2. \int \operatorname{sech}^2 ax dx = \frac{\tanh ax}{a}$$

$$17.32.3. \int \operatorname{sech}^3 ax dx = \frac{\operatorname{sech} ax \tanh ax}{2a} + \frac{1}{2a} \tan^{-1} \sinh ax$$

$$17.32.4. \int x \operatorname{sech} ax dx = \frac{1}{a^2} \left\{ \frac{(ax)^2}{2} - \frac{(ax)^4}{8} + \frac{5(ax)^6}{144} + \dots \frac{(-1)^n E_n(ax)^{2n+2}}{(2n+2)(2n)!} + \dots \right\}$$

$$17.32.5. \int x \operatorname{sech}^2 ax dx = \frac{x \tanh ax}{a} - \frac{1}{a^2} \ln \cosh ax$$

$$17.32.6. \int \frac{\operatorname{sech} ax}{x} dx = \ln x - \frac{(ax)^2}{4} + \frac{5(ax)^4}{96} - \frac{61(ax)^6}{4320} + \dots \frac{(-1)^n E_n(ax)^{2n}}{2n(2n)!} + \dots$$

$$17.32.7. \int \operatorname{sech}^n ax dx = \frac{\operatorname{sech}^{n-2} ax \tanh ax}{a(n-1)} + \frac{n-2}{n-1} \int \operatorname{sech}^{n-2} ax dx$$

**(33) Integrals Involving  $\operatorname{csch} ax$** 

---

$$17.33.1. \int \operatorname{csch} ax dx = \frac{1}{a} \ln \tanh \frac{ax}{2}$$

$$17.33.2. \int \operatorname{csch}^2 ax dx = -\frac{\coth ax}{a}$$

$$17.33.3. \int \operatorname{csch}^3 ax dx = -\frac{\operatorname{csch} ax \coth ax}{2a} - \frac{1}{2a} \ln \tanh \frac{ax}{2}$$

$$17.33.4. \int x \operatorname{csch} ax dx = \frac{1}{a^2} \left\{ ax - \frac{(ax)^3}{18} + \frac{7(ax)^5}{1800} + \dots + \frac{2(-1)^n (2^{2n-1} - 1) B_n (ax)^{2n+1}}{(2n+1)!} + \dots \right\}$$

$$17.33.5. \int x \operatorname{csch}^2 ax dx = -\frac{x \coth ax}{a} + \frac{1}{a^2} \ln \sinh ax$$

$$17.33.6. \int \frac{\operatorname{csch} ax}{x} dx = -\frac{1}{ax} - \frac{ax}{6} + \frac{7(ax)^3}{1080} + \dots \frac{(-1)^n 2(2^{2n-1} - 1) B_n (ax)^{2n-1}}{(2n-1)(2n)!} + \dots$$

$$17.33.7. \int \operatorname{csch}^n ax dx = \frac{-\operatorname{csch}^{n-2} ax \coth ax}{a(n-1)} - \frac{n-2}{n-1} \int \operatorname{csch}^{n-2} ax dx$$

**(34) Integrals Involving Inverse Hyperbolic Functions**

$$17.34.1. \int \sinh^{-1} \frac{x}{a} dx = x \sinh^{-1} \frac{x}{a} - \sqrt{x^2 + a^2}$$

$$17.34.2. \int x \sinh^{-1} \frac{x}{a} dx = \left( \frac{x^2}{2} + \frac{a^2}{4} \right) \sinh^{-1} \frac{x}{a} - x \frac{\sqrt{x^2 + a^2}}{4}$$

$$17.34.3. \int \frac{\sinh^{-1}(x/a)}{x} dx = \begin{cases} \frac{x}{a} - \frac{(x/a)^3}{2 \cdot 3 \cdot 3} + \frac{1 \cdot 3 \cdot 5(x/a)^5}{2 \cdot 4 \cdot 5 \cdot 5} - \frac{1 \cdot 3 \cdot 5 \cdot 7(x/a)^7}{2 \cdot 4 \cdot 6 \cdot 7 \cdot 7} + \dots & |x| < a \\ \frac{\ln^2(2x/a)}{2} - \frac{(a/x)^2}{2 \cdot 2 \cdot 2} + \frac{1 \cdot 3(a/x)^4}{2 \cdot 4 \cdot 4 \cdot 4} - \frac{1 \cdot 3 \cdot 5(a/x)^6}{2 \cdot 4 \cdot 6 \cdot 6 \cdot 6} + \dots & x > a \\ -\frac{\ln^2(-2x/a)}{2} + \frac{(a/x)^2}{2 \cdot 2 \cdot 2} - \frac{1 \cdot 3(a/x)^4}{2 \cdot 4 \cdot 4 \cdot 4} + \frac{1 \cdot 3 \cdot 5(a/x)^6}{2 \cdot 4 \cdot 6 \cdot 6 \cdot 6} - \dots & x < -a \end{cases}$$

$$17.34.4. \int \cosh^{-1} \frac{x}{a} dx = \begin{cases} x \cosh^{-1}(x/a) - \sqrt{x^2 - a^2}, & \cosh^{-1}(x/a) > 0 \\ x \cosh^{-1}(x/a) + \sqrt{x^2 - a^2}, & \cosh^{-1}(x/a) < 0 \end{cases}$$

$$17.34.5. \int x \cosh^{-1} \frac{x}{a} dx = \begin{cases} \frac{1}{4}(2x^2 - a^2) \cosh^{-1}(x/a) - \frac{1}{4}x \sqrt{x^2 - a^2}, & \cosh^{-1}(x/a) > 0 \\ \frac{1}{4}(2x^2 - a^2) \cosh^{-1}(x/a) + \frac{1}{4}x \sqrt{x^2 - a^2}, & \cosh^{-1}(x/a) < 0 \end{cases}$$

$$17.34.6. \int \frac{\cosh^{-1}(x/a)}{x} dx = \pm \left[ \frac{1}{2} \ln^2(2x/a) + \frac{(a/x)^2}{2 \cdot 2 \cdot 2} + \frac{1 \cdot 3(a/x)^4}{2 \cdot 4 \cdot 4 \cdot 4} + \frac{1 \cdot 3 \cdot 5(a/x)^6}{2 \cdot 4 \cdot 6 \cdot 6 \cdot 6} + \dots \right] \\ + \text{if } \cosh^{-1}(x/a) > 0, - \text{if } \cosh^{-1}(x/a) < 0$$

$$17.34.7. \int \tanh^{-1} \frac{x}{a} dx = x \tanh^{-1} \frac{x}{a} + \frac{a}{2} \ln(a^2 - x^2)$$

$$17.34.8. \int x \tanh^{-1} \frac{x}{a} dx = \frac{ax}{2} + \frac{1}{2}(x^2 - a^2) \tanh^{-1} \frac{x}{a}$$

$$17.34.9. \int \frac{\tanh^{-1}(x/a)}{x} dx = \frac{x}{a} + \frac{(x/a)^3}{3^2} + \frac{(x/a)^5}{5^2} + \dots$$

$$17.34.10. \int \coth^{-1} \frac{x}{a} dx = x \coth^{-1} x + \frac{a}{2} \ln(x^2 - a^2)$$

$$17.34.11. \int x \coth^{-1} \frac{x}{a} dx = \frac{ax}{2} + \frac{1}{2}(x^2 - a^2) \coth^{-1} \frac{x}{a}$$

$$17.34.12. \int \frac{\coth^{-1}(x/a)}{x} dx = - \left( \frac{a}{x} + \frac{(a/x)^3}{3^2} + \frac{(a/x)^5}{5^2} + \dots \right)$$

$$17.34.13. \int \operatorname{sech}^{-1} \frac{x}{a} dx = \begin{cases} x \operatorname{sech}^{-1}(x/a) + a \sin^{-1}(x/a), & \operatorname{sech}^{-1}(x/a) > 0 \\ x \operatorname{sech}^{-1}(x/a) - a \sin^{-1}(x/a), & \operatorname{sech}^{-1}(x/a) < 0 \end{cases}$$

$$17.34.14. \int \operatorname{csch}^{-1} \frac{x}{a} dx = x \operatorname{csch}^{-1} \frac{x}{a} \pm a \sinh^{-1} \frac{x}{a} \quad (+\text{if } x > 0, -\text{if } x < 0)$$

$$17.34.15. \int x^m \sinh^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \sinh^{-1} \frac{x}{a} - \frac{1}{m+1} \int \frac{x^{m+1}}{\sqrt{x^2 + a^2}} dx$$

$$17.34.16. \int x^m \cosh^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^{m+1}}{m+1} \cosh^{-1} \frac{x}{a} - \frac{1}{m+1} \int \frac{x^{m+1}}{\sqrt{x^2 - a^2}} dx & \cosh^{-1}(x/a) > 0 \\ \frac{x^{m+1}}{m+1} \cosh^{-1} \frac{x}{a} + \frac{1}{m+1} \int \frac{x^{m+1}}{\sqrt{x^2 - a^2}} dx & \cosh^{-1}(x/a) < 0 \end{cases}$$

$$17.34.17. \int x^m \tanh^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \tanh^{-1} \frac{x}{a} - \frac{a}{m+1} \int \frac{x^{m+1}}{a^2 - x^2} dx$$

$$17.34.18. \int x^m \coth^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \coth^{-1} \frac{x}{a} - \frac{1}{m+1} \int \frac{x^{m+1}}{a^2 - x^2} dx$$

$$17.34.19. \int x^m \operatorname{sech}^{-1} \frac{x}{a} dx = \begin{cases} \frac{x^{m+1}}{m+1} \operatorname{sech}^{-1} \frac{x}{a} + \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{a^2 - x^2}} & \operatorname{sech}^{-1}(x/a) > 0 \\ \frac{x^{m+1}}{m+1} \operatorname{sech}^{-1} \frac{x}{a} - \frac{1}{m+1} \int \frac{x^m dx}{\sqrt{a^2 - x^2}} & \operatorname{sech}^{-1}(x/a) < 0 \end{cases}$$

$$17.34.20. \int x^m \operatorname{csch}^{-1} \frac{x}{a} dx = \frac{x^{m+1}}{m+1} \operatorname{csch}^{-1} \frac{x}{a} \pm \frac{a}{m+1} \int \frac{x^m dx}{\sqrt{x^2 + a^2}} \quad (+\text{if } x > 0, -\text{if } x < 0)$$

# 18 DEFINITE INTEGRALS

## Definition of a Definite Integral

Let  $f(x)$  be defined in an interval  $a \leq x \leq b$ . Divide the interval into  $n$  equal parts of length  $\Delta x = (b - a)/n$ . Then the definite integral of  $f(x)$  between  $x = a$  and  $x = b$  is defined as

$$18.1. \int_a^b f(x) dx = \lim_{n \rightarrow \infty} \{f(a)\Delta x + f(a + \Delta x)\Delta x + f(a + 2\Delta x)\Delta x + \dots + f(a + (n-1)\Delta x)\Delta x\}$$

The limit will certainly exist if  $f(x)$  is piecewise continuous.

If  $f(x) = \frac{d}{dx} g(x)$ , then by the fundamental theorem of the integral calculus the above definite integral can be evaluated by using the result

$$18.2. \int_a^b f(x) dx = \int_a^b \frac{d}{dx} g(x) dx = g(x) \Big|_a^b = g(b) - g(a)$$

If the interval is infinite or if  $f(x)$  has a singularity at some point in the interval, the definite integral is called an *improper integral* and can be defined by using appropriate limiting procedures. For example,

$$18.3. \int_a^{\infty} f(x) dx = \lim_{b \rightarrow \infty} \int_a^b f(x) dx$$

$$18.4. \int_{-\infty}^{\infty} f(x) dx = \lim_{\substack{a \rightarrow -\infty \\ b \rightarrow \infty}} \int_a^b f(x) dx$$

$$18.5. \int_a^b f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{a-\epsilon}^{b-\epsilon} f(x) dx \quad \text{if } b \text{ is a singular point.}$$

$$18.6. \int_a^b f(x) dx = \lim_{\epsilon \rightarrow 0} \int_{a+\epsilon}^b f(x) dx \quad \text{if } a \text{ is a singular point.}$$

## General Formulas Involving Definite Integrals

$$18.7. \int_a^b \{f(x) \pm g(x) \pm h(x) \pm \dots\} dx = \int_a^b f(x) dx \pm \int_a^b g(x) dx \pm \int_a^b h(x) dx \pm \dots$$

$$18.8. \int_a^b cf(x) dx = c \int_a^b f(x) dx \quad \text{where } c \text{ is any constant.}$$

$$18.9. \int_a^a f(x) dx = 0$$

$$18.10. \int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$18.11. \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

**18.12.**  $\int_a^b f(x) dx = (b - a) f(c)$  where  $c$  is between  $a$  and  $b$ .

This is called the *mean value theorem* for definite integrals and is valid if  $f(x)$  is continuous in  $a \leq x \leq b$ .

**18.13.**  $\int_a^b f(x) g(x) dx = f(c) \int_a^b g(x) dx$  where  $c$  is between  $a$  and  $b$

This is a generalization of 18.12 and is valid if  $f(x)$  and  $g(x)$  are continuous in  $a \leq x \leq b$  and  $g(x) \geq 0$ .

### Leibnitz's Rules for Differentiation of Integrals

---

**18.14.**  $\frac{d}{d\alpha} \int_{\phi_1(\alpha)}^{\phi_2(\alpha)} F(x, \alpha) dx = \int_{\phi_1(\alpha)}^{\phi_2(\alpha)} \frac{\partial F}{d\alpha} dx + F(\phi_2, \alpha) \frac{d\phi_2}{d\alpha} - F(\phi_1, \alpha) \frac{d\phi_1}{d\alpha}$

### Approximate Formulas for Definite Integrals

---

In the following the interval from  $x = a$  to  $x = b$  is subdivided into  $n$  equal parts by the points  $a = x_0, x_1, x_2, \dots, x_{n-1}, x_n = b$  and we let  $y_0 = f(x_0), y_1 = f(x_1), y_2 = f(x_2), \dots, y_n = f(x_n), h = (b - a)/n$ .

**Rectangular formula:**

**18.15.**  $\int_a^b f(x) dx \approx h(y_0 + y_1 + y_2 + \dots + y_{n-1})$

**Trapezoidal formula:**

**18.16.**  $\int_a^b f(x) dx \approx \frac{h}{2}(y_0 + 2y_1 + 2y_2 + \dots + 2y_{n-1} + y_n)$

**Simpson's formula (or parabolic formula) for  $n$  even:**

**18.17.**  $\int_a^b f(x) dx \approx \frac{h}{3}(y_0 + 4y_1 + 2y_2 + 4y_3 + \dots + 2y_{n-2} + 4y_{n-1} + y_n)$

### Definite Integrals Involving Rational or Irrational Expressions

---

**18.18.**  $\int_0^\infty \frac{dx}{x^2 + a^2} = \frac{\pi}{2a}$

**18.19.**  $\int_0^\infty \frac{x^{p-1} dx}{1+x} = \frac{\pi}{\sin p\pi}, \quad 0 < p < 1$

**18.20.**  $\int_0^\infty \frac{x^m dx}{x^n + a^n} = \frac{\pi a^{m+1-n}}{n \sin[(m+1)\pi/n]}, \quad 0 < m+1 < n$

**18.21.**  $\int_0^\infty \frac{x^m dx}{1+2x \cos \beta + x^2} = \frac{\pi}{\sin m\pi} \frac{\sin m\beta}{\sin \beta}$

**18.22.**  $\int_0^a \frac{dx}{\sqrt{a^2 - x^2}} = \frac{\pi}{2}$

**18.23.**  $\int_0^a \sqrt{a^2 - x^2} dx = \frac{\pi a^2}{4}$

$$18.24. \int_0^a x^m (a^n - x^n)^p dx = \frac{a^{m+1+n p} \Gamma[(m+1)/n] \Gamma(p+1)}{n \Gamma[(m+1)/n + p + 1]}$$

$$18.25. \int_0^\infty \frac{x^m dx}{(x^n + a^n)^r} = \frac{(-1)^{r-1} \pi a^{m+1-n r} \Gamma[(m+1)/n]}{n \sin[(m+1)\pi/n] (r-1)! \Gamma[(m+1)/n - r + 1]}, \quad 0 < m+1 < nr$$

### Definite Integrals Involving Trigonometric Functions

---

All letters are considered positive unless otherwise indicated.

$$18.26. \int_0^\pi \sin mx \sin nx dx = \begin{cases} 0 & m, n \text{ integers and } m \neq n \\ \pi/2 & m, n \text{ integers and } m = n \end{cases}$$

$$18.27. \int_0^\pi \cos mx \cos nx dx = \begin{cases} 0 & m, n \text{ integers and } m \neq n \\ \pi/2 & m, n \text{ integers and } m = n \end{cases}$$

$$18.28. \int_0^\pi \sin mx \cos nx dx = \begin{cases} 0 & m, n \text{ integers and } m+n \text{ even} \\ 2m/(m^2 - n^2) & m, n \text{ integers and } m+n \text{ odd} \end{cases}$$

$$18.29. \int_0^{\pi/2} \sin^2 x dx = \int_0^{\pi/2} \cos^2 x dx = \frac{\pi}{4}$$

$$18.30. \int_0^{\pi/2} \sin^{2m} x dx = \int_0^{\pi/2} \cos^{2m} x dx = \frac{1 \cdot 3 \cdot 5 \cdots 2m-1}{2 \cdot 4 \cdot 6 \cdots 2m} \frac{\pi}{2}, \quad m=1, 2, \dots$$

$$18.31. \int_0^{\pi/2} \sin^{2m+1} x dx = \int_0^{\pi/2} \cos^{2m+1} x dx = \frac{2 \cdot 4 \cdot 6 \cdots 2m}{1 \cdot 3 \cdot 5 \cdots 2m+1}, \quad m=1, 2, \dots$$

$$18.32. \int_0^{\pi/2} \sin^{2p-1} x \cos^{2q-1} x dx = \frac{\Gamma(p) \Gamma(q)}{2 \Gamma(p+q)}$$

$$18.33. \int_0^\infty \frac{\sin px}{x} dx = \begin{cases} \pi/2 & p > 0 \\ 0 & p = 0 \\ -\pi/2 & p < 0 \end{cases}$$

$$18.34. \int_0^\infty \frac{\sin px \cos qx}{x} dx = \begin{cases} 0 & p > q > 0 \\ \pi/2 & 0 < p < q \\ \pi/4 & p = q > 0 \end{cases}$$

$$18.35. \int_0^\infty \frac{\sin px \sin qx}{x^2} dx = \begin{cases} \pi p/2 & 0 < p \leq q \\ \pi q/2 & p \geq q > 0 \end{cases}$$

$$18.36. \int_0^\infty \frac{\sin^2 px}{x^2} dx = \frac{\pi p}{2}$$

$$18.37. \int_0^\infty \frac{1 - \cos px}{x^2} dx = \frac{\pi p}{2}$$

$$18.38. \int_0^\infty \frac{\cos px - \cos qx}{x} dx = \ln \frac{q}{p}$$

$$18.39. \int_0^\infty \frac{\cos px - \cos qx}{x^2} dx = \frac{\pi(q-p)}{2}$$

$$18.40. \int_0^\infty \frac{\cos mx}{x^2 + a^2} dx = \frac{\pi}{2a} e^{-ma}$$

$$18.41. \int_0^\infty \frac{x \sin mx}{x^2 + a^2} dx = \frac{\pi}{2} e^{-ma}$$

$$18.42. \int_0^\infty \frac{\sin mx}{x(x^2 + a^2)} dx = \frac{\pi}{2a^2} (1 - e^{-ma})$$

$$18.43. \int_0^{2\pi} \frac{dx}{a + b \sin x} = \frac{2\pi}{\sqrt{a^2 - b^2}}$$

$$18.44. \int_0^{2\pi} \frac{dx}{a + b \cos x} = \frac{2\pi}{\sqrt{a^2 - b^2}}$$

$$18.45. \int_0^{\pi/2} \frac{dx}{a + b \cos x} = \frac{\cos^{-1}(b/a)}{\sqrt{a^2 - b^2}}$$

$$18.46. \int_0^{2\pi} \frac{dx}{(a + b \sin x)^2} = \int_0^{2\pi} \frac{dx}{(a + b \cos x)^2} = \frac{2\pi a}{(a^2 - b^2)^{3/2}}$$

$$18.47. \int_0^{2\pi} \frac{dx}{1 - 2a \cos x + a^2} = \frac{2\pi}{1 - a^2}, \quad 0 < a < 1$$

$$18.48. \int_0^\pi \frac{x \sin x dx}{1 - 2a \cos x + a^2} = \begin{cases} (\pi/a) \ln(1+a), & |a| < 1 \\ \pi \ln(1+1/a), & |a| > 1 \end{cases}$$

$$18.49. \int_0^\pi \frac{\cos mx dx}{1 - 2a \cos x + a^2} = \frac{\pi a^m}{1 - a^2}, \quad a^2 < 1, \quad m = 0, 1, 2, \dots$$

$$18.50. \int_0^\infty \sin ax^2 dx = \int_0^\infty \cos ax^2 dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}}$$

$$18.51. \int_0^\infty \sin ax^n dx = \frac{1}{na^{1/n}} \Gamma(1/n) \sin \frac{\pi}{2n}, \quad n > 1$$

$$18.52. \int_0^\infty \cos ax^n dx = \frac{1}{na^{1/n}} \Gamma(1/n) \cos \frac{\pi}{2n}, \quad n > 1$$

$$18.53. \int_0^\infty \frac{\sin x}{\sqrt{x}} dx = \int_0^\infty \frac{\cos x}{\sqrt{x}} dx = \sqrt{\frac{\pi}{2}}$$

$$18.54. \int_0^\infty \frac{\sin x}{x^p} dx = \frac{\pi}{2\Gamma(p)\sin(p\pi/2)}, \quad 0 < p < 1$$

$$18.55. \int_0^\infty \frac{\cos x}{x^p} dx = \frac{\pi}{2\Gamma(p)\cos(p\pi/2)}, \quad 0 < p < 1$$

$$18.56. \int_0^\infty \sin ax^2 \cos 2bx dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}} \left( \cos \frac{b^2}{a} - \sin \frac{b^2}{a} \right)$$

$$18.57. \int_0^{\infty} \cos ax^2 \cos 2bx dx = \frac{1}{2} \sqrt{\frac{\pi}{2a}} \left( \cos \frac{b^2}{a} + \sin \frac{b^2}{a} \right)$$

$$18.58. \int_0^{\infty} \frac{\sin^3 x}{x^3} dx = \frac{3\pi}{8}$$

$$18.59. \int_0^{\infty} \frac{\sin^4 x}{x^4} dx = \frac{\pi}{3}$$

$$18.60. \int_0^{\infty} \frac{\tan x}{x} dx = \frac{\pi}{2}$$

$$18.61. \int_0^{\pi/2} \frac{dx}{1 + \tan^m x} = \frac{\pi}{4}$$

$$18.62. \int_0^{\pi/2} \frac{x}{\sin x} dx = 2 \left\{ \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \dots \right\}$$

$$18.63. \int_0^1 \frac{\tan^{-1} x}{x} dx = \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \frac{1}{7^2} + \dots$$

$$18.64. \int_0^1 \frac{\sin^{-1} x}{x} dx = \frac{\pi}{2} \ln 2$$

$$18.65. \int_0^1 \frac{1 - \cos x}{x} dx - \int_1^{\infty} \frac{\cos x}{x} dx = \gamma$$

$$18.66. \int_0^{\infty} \left( \frac{1}{1+x^2} - \cos x \right) \frac{dx}{x} = \gamma$$

$$18.67. \int_0^{\infty} \frac{\tan^{-1} px - \tan^{-1} qx}{x} dx = \frac{\pi}{2} \ln \frac{p}{q}$$

### Definite Integrals Involving Exponential Functions

---

Some integrals contain Euler's constant  $\gamma = 0.5772156 \dots$  (see 1.3, page 3).

$$18.68. \int_0^{\infty} e^{-ax} \cos bx dx = \frac{a}{a^2 + b^2}$$

$$18.69. \int_0^{\infty} e^{-ax} \sin bx dx = \frac{b}{a^2 + b^2}$$

$$18.70. \int_0^{\infty} \frac{e^{-ax} \sin bx}{x} dx = \tan^{-1} \frac{b}{a}$$

$$18.71. \int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} dx = \ln \frac{b}{a}$$

$$18.72. \int_0^{\infty} e^{-ax^2} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}}$$

$$18.73. \int_0^{\infty} e^{-ax^2} \cos bx dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-b^2/4a}$$

$$18.74. \int_0^\infty e^{-(ax^2+bx+c)} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{(b^2-4ac)/4a} \operatorname{erfc} \frac{b}{2\sqrt{a}}$$

where  $\operatorname{erfc}(p) = \frac{2}{\sqrt{\pi}} \int_p^\infty e^{-x^2} dx$

$$18.75. \int_{-\infty}^\infty e^{-(ax^2+bx+c)} dx = \sqrt{\frac{\pi}{a}} e^{(b^2-4ac)/4a}$$

$$18.76. \int_0^\infty x^n e^{-ax} dx = \frac{\Gamma(n+1)}{a^{n+1}}$$

$$18.77. \int_0^\infty x^m e^{-ax^2} dx = \frac{\Gamma[(m+1)/2]}{2a^{(m+1)/2}}$$

$$18.78. \int_0^\infty e^{-(ax^2+b/x^2)} dx = \frac{1}{2} \sqrt{\frac{\pi}{a}} e^{-2\sqrt{ab}}$$

$$18.79. \int_0^\infty \frac{x dx}{e^x - 1} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots = \frac{\pi^2}{6}$$

$$18.80. \int_0^\infty \frac{x^{n-1}}{e^x - 1} dx = \Gamma(n) \left( \frac{1}{1^n} + \frac{1}{2^n} + \frac{1}{3^n} + \dots \right)$$

For even  $n$  this can be summed in terms of Bernoulli numbers (See pages 142–143).

$$18.81. \int_0^\infty \frac{x dx}{e^x + 1} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots = \frac{\pi^2}{12}$$

$$18.82. \int_0^\infty \frac{x^{n-1}}{e^x + 1} dx = \Gamma(n) \left( \frac{1}{1^n} - \frac{1}{2^n} + \frac{1}{3^n} - \dots \right)$$

For some positive integer values of  $n$  the series can be summed (See 23.10).

$$18.83. \int_0^\infty \frac{\sin mx}{e^{2\pi x} - 1} dx = \frac{1}{4} \coth \frac{m}{2} - \frac{1}{2m}$$

$$18.84. \int_0^\infty \left( \frac{1}{1+x} - e^{-x} \right) \frac{dx}{x} = \gamma$$

$$18.85. \int_0^\infty \frac{e^{-x^2} - e^{-x}}{x} dx = \frac{1}{2} \gamma$$

$$18.86. \int_0^\infty \left( \frac{1}{e^x - 1} - \frac{e^{-x}}{x} \right) dx = \gamma$$

$$18.87. \int_0^\infty \frac{e^{-ax} - e^{-bx}}{x \sec px} dx = \frac{1}{2} \ln \left( \frac{b^2 + p^2}{a^2 + p^2} \right)$$

$$18.88. \int_0^\infty \frac{e^{-ax} - e^{-bx}}{x \csc px} dx = \tan^{-1} \frac{b}{p} - \tan^{-1} \frac{a}{p}$$

$$18.89. \int_0^\infty \frac{e^{-ax} (1 - \cos x)}{x^2} dx = \cot^{-1} a - \frac{a}{2} \ln(a^2 + 1)$$

### Definite Integrals Involving Logarithmic Functions

---

$$18.90. \int_0^1 x^m (\ln x)^n dx = \frac{(-1)^n n!}{(m+1)^{n+1}} \quad m > -1, \quad n = 0, 1, 2, \dots$$

If  $n \neq 0, 1, 2, \dots$  replace  $n!$  by  $\Gamma(n+1)$ .

$$18.91. \int_0^1 \frac{\ln x}{1+x} dx = -\frac{\pi^2}{12}$$

$$18.92. \int_0^1 \frac{\ln x}{1-x} dx = -\frac{\pi^2}{6}$$

$$18.93. \int_0^1 \frac{\ln(1+x)}{x} dx = \frac{\pi^2}{12}$$

$$18.94. \int_0^1 \frac{\ln(1-x)}{x} dx = -\frac{\pi^2}{6}$$

$$18.95. \int_0^1 \ln x \ln(1+x) dx = 2 - 2 \ln 2 - \frac{\pi^2}{12}$$

$$18.96. \int_0^1 \ln x \ln(1-x) dx = 2 - \frac{\pi^2}{6}$$

$$18.97. \int_0^\infty \frac{x^{p-1} \ln x}{1+x} dx = -\pi^2 \csc p\pi \cot p\pi \quad 0 < p < 1$$

$$18.98. \int_0^1 \frac{x^m - x^n}{\ln x} dx = \ln \frac{m+1}{n+1}$$

$$18.99. \int_0^\infty e^{-x} \ln x dx = -\gamma$$

$$18.100. \int_0^\infty e^{-x^2} \ln x dx = -\frac{\sqrt{\pi}}{4}(\gamma + 2 \ln 2)$$

$$18.101. \int_0^\infty \ln \left( \frac{e^x + 1}{e^x - 1} \right) dx = \frac{\pi^2}{4}$$

$$18.102. \int_0^{\pi/2} \ln \sin x dx = \int_0^{\pi/2} \ln \cos x dx = -\frac{\pi}{2} \ln 2$$

$$18.103. \int_0^{\pi/2} (\ln \sin x)^2 dx = \int_0^{\pi/2} (\ln \cos x)^2 dx = \frac{\pi}{2} (\ln 2)^2 + \frac{\pi^3}{24}$$

$$18.104. \int_0^\pi x \ln \sin x dx = -\frac{\pi^2}{2} \ln 2$$

$$18.105. \int_0^{\pi/2} \sin x \ln \sin x dx = \ln 2 - 1$$

$$18.106. \int_0^{2\pi} \ln(a + b \sin x) dx = \int_0^{2\pi} \ln(a + b \cos x) dx = 2\pi \ln(a + \sqrt{a^2 - b^2})$$

$$18.107. \int_0^\pi \ln(a + b \cos x) dx = \pi \ln \left( \frac{a + \sqrt{a^2 - b^2}}{2} \right)$$

$$18.108. \int_0^\pi \ln(a^2 - 2ab \cos x + b^2) dx = \begin{cases} 2\pi \ln a, & a \geq b > 0 \\ 2\pi \ln b, & b \geq a > 0 \end{cases}$$

$$18.109. \int_0^{\pi/4} \ln(1 + \tan x) dx = \frac{\pi}{8} \ln 2$$

$$18.110. \int_0^{\pi/2} \sec x \ln \left( \frac{1+b \cos x}{1+a \cos x} \right) dx = \frac{1}{2} \{ (\cos^{-1} a)^2 - (\cos^{-1} b)^2 \}$$

$$18.111. \int_0^a \ln \left( 2 \sin \frac{x}{2} \right) dx = - \left( \frac{\sin a}{1^2} + \frac{\sin 2a}{2^2} + \frac{\sin 3a}{3^2} + \dots \right)$$

See also 18.102.

### Definite Integrals Involving Hyperbolic Functions

---

$$18.112. \int_0^\infty \frac{\sin ax}{\sinh bx} dx = \frac{\pi}{2b} \tanh \frac{a\pi}{2b}$$

$$18.113. \int_0^\infty \frac{\cos ax}{\cosh bx} dx = \frac{\pi}{2b} \operatorname{sech} \frac{a\pi}{2b}$$

$$18.114. \int_0^\infty \frac{x dx}{\sinh ax} = \frac{\pi^2}{4a^2}$$

$$18.115. \int_0^\infty \frac{x^n dx}{\sinh ax} = \frac{2^{n+1}-1}{2^n a^{n+1}} \Gamma(n+1) \left\{ \frac{1}{1^{n+1}} + \frac{1}{2^{n+1}} + \frac{1}{3^{n+1}} + \dots \right\}$$

If  $n$  is an odd positive integer, the series can be summed.

$$18.116. \int_0^\infty \frac{\sinh ax}{e^{bx} + 1} dx = \frac{\pi}{2b} \csc \frac{a\pi}{b} - \frac{1}{2a}$$

$$18.117. \int_0^\infty \frac{\sinh ax}{e^{bx} - 1} dx = \frac{1}{2a} - \frac{\pi}{2b} \cot \frac{a\pi}{b}$$

### Miscellaneous Definite Integrals

---

$$18.118. \int_0^\infty \frac{f(ax) - f(bx)}{x} dx = \{f(0) - f(\infty)\} \ln \frac{b}{a}$$

This is called *Frullani's integral*. It holds if  $f'(x)$  is continuous and  $\int_0^\infty \frac{f(x) - f(\infty)}{x} dx$  converges.

$$18.119. \int_0^1 \frac{dx}{x^x} = \frac{1}{1^1} + \frac{1}{2^2} + \frac{1}{3^3} + \dots$$

$$18.120. \int_{-a}^a (a+x)^{m-1} (a-x)^{n-1} dx = (2a)^{m+n-1} \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$$

## Section V: Differential Equations and Vector Analysis

# 19 BASIC DIFFERENTIAL EQUATIONS and SOLUTIONS

DIFFERENTIAL EQUATION	SOLUTION
19.1. Separation of variables	
$f_1(x) g_1(y) dx + f_2(x) g_2(y) dy = 0$	$\int \frac{f_1(x)}{f_2(x)} dx + \int \frac{g_2(y)}{g_1(y)} dy = c$
19.2. Linear first order equation	
$\frac{dy}{dx} + p(x)y = Q(x)$	$ye^{\int P dx} = \int Q e^{\int P dx} dx + c$
19.3. Bernoulli's equation	
$\frac{dy}{dx} + P(x)y = Q(x)y^n$	$ve^{(1-n)\int P dx} = (1-n) \int Q e^{(1-n)\int P dx} dx + c$ where $v = y^{1-n}$ . If $n = 1$ , the solution is $\ln y = \int (Q - P) dx + c$
19.4. Exact equation	
$M(x, y)dx + N(x, y)dy = 0$ where $\partial M / \partial y = \partial N / \partial x$ .	$\int M \partial x + \int \left( N - \frac{\partial}{\partial y} \int M \partial x \right) dy = c$ where $\partial x$ indicates that the integration is to be performed with respect to $x$ keeping $y$ constant.
19.5 Homogeneous equation	
$\frac{dy}{dx} = F\left(\frac{y}{x}\right)$	$\ln x = \int \frac{dv}{F(v) - v} + c$ where $v = y/x$ . If $F(v) = v$ , the solution is $y = cx$ .

<b>19.6.</b> $y F(xy) dx + x G(xy) dy = 0$	$\ln x = \int \frac{G(v) dv}{v \{G(v) - F(v)\}} + c$ where $v = xy$ . If $G(v) = F(v)$ , the solution is $xy = c$ .
<b>19.7.</b> Linear, homogeneous second order equation $\frac{d^2y}{dx^2} + a \frac{dy}{dx} + by = 0$ $a, b$ are real constants.	Let $m_1, m_2$ be the roots of $m^2 + am + b = 0$ . Then there are 3 cases. <b>Case 1.</b> $m_1, m_2$ real and distinct: $y = c_1 e^{m_1 x} + c_2 e^{m_2 x}$ <b>Case 2.</b> $m_1, m_2$ real and equal: $y = c_1 e^{m_1 x} + c_2 x e^{m_1 x}$ <b>Case 3.</b> $m_1 = p + qi, m_2 = p - qi$ : $y = e^{px} (c_1 \cos qx + c_2 \sin qx)$ where $p = -a/2, q = \sqrt{b - a^2/4}$ .
<b>19.8.</b> Linear, nonhomogeneous second order equation $\frac{d^2y}{dx^2} + a \frac{dy}{dx} + by = R(x)$ $a, b$ are real constants.	There are 3 cases corresponding to those of entry 19.7 above. <b>Case 1.</b> $y = c_1 e^{m_1 x} + c_2 e^{m_2 x} + \frac{e^{m_1 x}}{m_1 - m_2} \int e^{-m_1 x} R(x) dx + \frac{e^{m_2 x}}{m_2 - m_1} \int e^{-m_2 x} R(x) dx$  <b>Case 2.</b> $y = c_1 e^{m_1 x} + c_2 x e^{m_1 x} + x e^{m_1 x} \int e^{-m_1 x} R(x) dx - e^{m_1 x} \int x e^{-m_1 x} R(x) dx$  <b>Case 3.</b> $y = e^{px} (c_1 \cos qx + c_2 \sin qx) + \frac{e^{px} \sin qx}{q} \int e^{-px} R(x) \cos qx dx - \frac{e^{px} \cos qx}{q} \int e^{-px} R(x) \sin qx dx$

<b>19.9.</b> Euler or Cauchy equation $x^2 \frac{d^2y}{dx^2} + ax \frac{dy}{dx} + by = S(x)$	Putting $x = e^t$ , the equation becomes $\frac{d^2y}{dt^2} + (a-1) \frac{dy}{dt} + by = S(e^t)$ and can then be solved as in entries 19.7 and 19.8 above.
<b>19.10.</b> Bessel's equation $x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + (\lambda^2 x^2 - n^2)y = 0$	$y = c_1 J_n(\lambda x) + c_2 Y_n(\lambda x)$ See 27.1 to 27.15.
<b>19.11.</b> Transformed Bessel's equation $x^2 \frac{d^2y}{dx^2} + (2p+1)x \frac{dy}{dx} + (a^2 x^{2r} + \beta^2)y = 0$	$y = x^{-p} \left\{ c_1 J_{q/r} \left( \frac{\alpha}{r} x^r \right) + c_2 Y_{q/r} \left( \frac{\alpha}{r} x^r \right) \right\}$ where $q = \sqrt{p^2 - \beta^2}$ .
<b>19.12.</b> Legendre's equation $(1-x^2) \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} + n(n+1)y = 0$	$y = c_1 P_n(x) + c_2 Q_n(x)$ See 28.1 to 28.48.

# 20 FORMULAS from VECTOR ANALYSIS

## Vectors and Scalars

Various quantities in physics such as temperature, volume, and speed can be specified by a real number. Such quantities are called *scalars*.

Other quantities such as force, velocity, and momentum require for their specification a direction as well as magnitude. Such quantities are called *vectors*. A vector is represented by an arrow or directed line segment indicating direction. The magnitude of the vector is determined by the length of the arrow, using an appropriate unit.

## Notation for Vectors

A vector is denoted by a bold faced letter such as  $\mathbf{A}$  (Fig. 20-1). The magnitude is denoted by  $|\mathbf{A}|$  or  $A$ . The tail end of the arrow is called the *initial point*, while the head is called the *terminal point*.

## Fundamental Definitions

- Equality of vectors.** Two vectors are equal if they have the same magnitude and direction. Thus,  $\mathbf{A} = \mathbf{B}$  in (Fig. 20-1).
- Multiplication of a vector by a scalar.** If  $m$  is any real number (scalar), then  $m\mathbf{A}$  is a vector whose magnitude is  $|m|$  times the magnitude of  $\mathbf{A}$  and whose direction is the same as or opposite to  $\mathbf{A}$  according as  $m > 0$  or  $m < 0$ . If  $m = 0$ , then  $m\mathbf{A} = \mathbf{0}$  is called the *zero* or *null* vector.
- Sums of vectors.** The sum or resultant of  $\mathbf{A}$  and  $\mathbf{B}$  is a vector  $\mathbf{C} = \mathbf{A} + \mathbf{B}$  formed by placing the initial point  $\mathbf{B}$  on the terminal point  $\mathbf{A}$  and joining the initial point of  $\mathbf{A}$  to the terminal point of  $\mathbf{B}$  as in Fig. 20-2b. This definition is equivalent to the parallelogram law for vector addition as indicated in Fig. 20-2c. The vector  $\mathbf{A} - \mathbf{B}$  is defined as  $\mathbf{A} + (-\mathbf{B})$ .




Fig. 20-1




Fig. 20-2

Extension to sums of more than two vectors are immediate. Thus, Fig. 20-3 shows how to obtain the sum  $\mathbf{E}$  of the vectors  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{C}$ , and  $\mathbf{D}$ .




Fig. 20-3

- 4. Unit vectors.** A *unit vector* is a vector with unit magnitude. If  $\mathbf{A}$  is a vector, then a unit vector in the direction of  $\mathbf{A}$  is  $\mathbf{a} = \mathbf{A}/A$  where  $A > 0$ .

### Laws of Vector Algebra

If  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{C}$  are vectors and  $m$ ,  $n$  are scalars, then:

$$\mathbf{20.1.} \quad \mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A} \quad \text{Commutative law for addition}$$

$$\mathbf{20.2.} \quad \mathbf{A} + (\mathbf{B} + \mathbf{C}) = (\mathbf{A} + \mathbf{B}) + \mathbf{C} \quad \text{Associative law for addition}$$

$$\mathbf{20.3.} \quad m(n\mathbf{A}) = (mn)\mathbf{A} = n(m\mathbf{A}) \quad \text{Associative law for scalar multiplication}$$

$$\mathbf{20.4.} \quad (m+n)\mathbf{A} = m\mathbf{A} + n\mathbf{A} \quad \text{Distributive law}$$

$$\mathbf{20.5.} \quad m(\mathbf{A} + \mathbf{B}) = m\mathbf{A} + m\mathbf{B} \quad \text{Distributive law}$$

### Components of a Vector

A vector  $\mathbf{A}$  can be represented with initial point at the origin of a rectangular coordinate system. If  $\mathbf{i}$ ,  $\mathbf{j}$ ,  $\mathbf{k}$  are unit vectors in the directions of the positive  $x$ ,  $y$ ,  $z$  axes, then

$$\mathbf{20.6.} \quad \mathbf{A} = A_1\mathbf{i} + A_2\mathbf{j} + A_3\mathbf{k}$$

where  $A_1\mathbf{i}$ ,  $A_2\mathbf{j}$ ,  $A_3\mathbf{k}$  are called *component vectors* of  $\mathbf{A}$  in the  $\mathbf{i}$ ,  $\mathbf{j}$ ,  $\mathbf{k}$  directions and  $A_1$ ,  $A_2$ ,  $A_3$  are called the *components* of  $\mathbf{A}$ .




Fig. 20-4

### Dot or Scalar Product

$$\mathbf{20.7.} \quad \mathbf{A} \cdot \mathbf{B} = AB \cos \theta \quad 0 \leq \theta \leq \pi$$

where  $\theta$  is the angle between  $\mathbf{A}$  and  $\mathbf{B}$ .

Fundamental results follow:

**20.8.**  $\mathbf{A} \cdot \mathbf{B} = \mathbf{B} \cdot \mathbf{A}$  Commutative law

**20.9.**  $\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) = \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C}$  Distributive law

**20.10.**  $\mathbf{A} \cdot \mathbf{B} = A_1 B_1 + A_2 B_2 + A_3 B_3$

where  $\mathbf{A} = A_1 \mathbf{i} + A_2 \mathbf{j} + A_3 \mathbf{k}$ ,  $\mathbf{B} = B_1 \mathbf{i} + B_2 \mathbf{j} + B_3 \mathbf{k}$ .

### Cross or Vector Product

**20.11.**  $\mathbf{A} \times \mathbf{B} = AB \sin \theta \mathbf{u}$   $0 \leq \theta \leq \pi$

where  $\theta$  is the angle between  $\mathbf{A}$  and  $\mathbf{B}$  and  $\mathbf{u}$  is a unit vector perpendicular to the plane of  $\mathbf{A}$  and  $\mathbf{B}$  such that  $\mathbf{A}$ ,  $\mathbf{B}$ ,  $\mathbf{u}$  form a *right-handed system* (i.e., a right-threaded screw rotated through an angle less than  $180^\circ$  from  $\mathbf{A}$  to  $\mathbf{B}$  will advance in the direction of  $\mathbf{u}$  as in Fig. 20-5).

Fundamental results follow:

$$\begin{aligned} \mathbf{20.12.} \quad \mathbf{A} \times \mathbf{B} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \end{vmatrix} \\ &= (A_2 B_3 - A_3 B_2) \mathbf{i} + (A_3 B_1 - A_1 B_3) \mathbf{j} + (A_1 B_2 - A_2 B_1) \mathbf{k} \end{aligned}$$

**20.13.**  $\mathbf{A} \times \mathbf{B} = -(\mathbf{B} \times \mathbf{A})$

**20.14.**  $\mathbf{A} \times (\mathbf{B} + \mathbf{C}) = \mathbf{A} \times \mathbf{B} + \mathbf{A} \times \mathbf{C}$

**20.15.**  $|\mathbf{A} \times \mathbf{B}| = \text{area of parallelogram having sides } \mathbf{A} \text{ and } \mathbf{B}$




Fig. 20-5

### Miscellaneous Formulas Involving Dot and Cross Products

$$\mathbf{20.16.} \quad \mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \begin{vmatrix} A_1 & A_2 & A_3 \\ B_1 & B_2 & B_3 \\ C_1 & C_2 & C_3 \end{vmatrix} = A_1 B_2 C_3 + A_2 B_3 C_1 + A_3 B_1 C_2 - A_3 B_2 C_1 - A_2 B_1 C_3 - A_1 B_3 C_2$$

**20.17.**  $|\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})| = \text{volume of parallelepiped with sides } \mathbf{A}, \mathbf{B}, \mathbf{C}$

**20.18.**  $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$

**20.19.**  $(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{A}(\mathbf{B} \cdot \mathbf{C})$

**20.20.**  $(\mathbf{A} \times \mathbf{B}) \cdot (\mathbf{C} \times \mathbf{D}) = (\mathbf{A} \cdot \mathbf{C})(\mathbf{B} \cdot \mathbf{D}) - (\mathbf{A} \cdot \mathbf{D})(\mathbf{B} \cdot \mathbf{C})$

$$\begin{aligned} \mathbf{20.21.} \quad (\mathbf{A} \times \mathbf{B}) \times (\mathbf{C} \times \mathbf{D}) &= \mathbf{C}\{\mathbf{A} \cdot (\mathbf{B} \times \mathbf{D})\} - \mathbf{D}\{\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})\} \\ &= \mathbf{B}\{\mathbf{A} \cdot (\mathbf{C} \times \mathbf{D})\} - \mathbf{A}\{\mathbf{B} \cdot (\mathbf{C} \times \mathbf{D})\} \end{aligned}$$

## Derivatives of Vectors

---

The derivative of a vector function  $\mathbf{A}(u) = A_1(u)\mathbf{i} + A_2(u)\mathbf{j} + A_3(u)\mathbf{k}$  of the scalar variable  $u$  is given by

$$20.22. \quad \frac{d\mathbf{A}}{du} = \lim_{\Delta u \rightarrow 0} \frac{\mathbf{A}(u + \Delta u) - \mathbf{A}(u)}{\Delta u} = \frac{dA_1}{du}\mathbf{i} + \frac{dA_2}{du}\mathbf{j} + \frac{dA_3}{du}\mathbf{k}$$

Partial derivatives of a vector function  $\mathbf{A}(x, y, z)$  are similarly defined. We assume that all derivatives exist unless otherwise specified.

## Formulas Involving Derivatives

---

$$20.23. \quad \frac{d}{du}(\mathbf{A} \cdot \mathbf{B}) = \mathbf{A} \cdot \frac{d\mathbf{B}}{du} + \frac{d\mathbf{A}}{du} \cdot \mathbf{B}$$

$$20.24. \quad \frac{d}{du}(\mathbf{A} \times \mathbf{B}) = \mathbf{A} \times \frac{d\mathbf{B}}{du} + \frac{d\mathbf{A}}{du} \times \mathbf{B}$$

$$20.25. \quad \frac{d}{du}\{\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})\} = \frac{d\mathbf{A}}{du} \cdot (\mathbf{B} \times \mathbf{C}) + \mathbf{A} \cdot \left( \frac{d\mathbf{B}}{du} \times \mathbf{C} \right) + \mathbf{A} \cdot \left( \mathbf{B} \times \frac{d\mathbf{C}}{du} \right)$$

$$20.26. \quad \mathbf{A} \cdot \frac{d\mathbf{A}}{du} = A \frac{dA}{du}$$

$$20.27. \quad \mathbf{A} \cdot \frac{d\mathbf{A}}{du} = 0 \quad \text{if } |\mathbf{A}| \text{ is a constant}$$

## The Del Operator

---

The operator *del* is defined by

$$20.28. \quad \nabla = \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z}$$

In the following results we assume that  $U = U(x, y, z)$ ,  $V = V(x, y, z)$ ,  $\mathbf{A} = \mathbf{A}(x, y, z)$  and  $\mathbf{B} = \mathbf{B}(x, y, z)$  have partial derivatives.

## The Gradient

---

$$20.29. \quad \text{Gradient of } U = \text{grad } U = \nabla U = \left( \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) U = \frac{\partial U}{\partial x} \mathbf{i} + \frac{\partial U}{\partial y} \mathbf{j} + \frac{\partial U}{\partial z} \mathbf{k}$$

## The Divergence

---

$$\begin{aligned} 20.30. \quad \text{Divergence of } \mathbf{A} = \text{div } \mathbf{A} = \nabla \cdot \mathbf{A} &= \left( \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \cdot (A_1 \mathbf{i} + A_2 \mathbf{j} + A_3 \mathbf{k}) \\ &= \frac{\partial A_1}{\partial x} + \frac{\partial A_2}{\partial y} + \frac{\partial A_3}{\partial z} \end{aligned}$$

### The Curl

---

**20.31.** Curl of  $\mathbf{A} = \operatorname{curl} \mathbf{A} = \nabla \times \mathbf{A}$

$$\begin{aligned}
 &= \left( \mathbf{i} \frac{\partial}{\partial x} + \mathbf{j} \frac{\partial}{\partial y} + \mathbf{k} \frac{\partial}{\partial z} \right) \times (A_1 \mathbf{i} + A_2 \mathbf{j} + A_3 \mathbf{k}) \\
 &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ A_1 & A_2 & A_3 \end{vmatrix} \\
 &= \left( \frac{\partial A_3}{\partial y} - \frac{\partial A_2}{\partial z} \right) \mathbf{i} + \left( \frac{\partial A_1}{\partial z} - \frac{\partial A_3}{\partial x} \right) \mathbf{j} + \left( \frac{\partial A_2}{\partial x} - \frac{\partial A_1}{\partial y} \right) \mathbf{k}
 \end{aligned}$$

### The Laplacian

---

**20.32.** Laplacian of  $U = \nabla^2 U = \nabla \cdot (\nabla U) = \frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} + \frac{\partial^2 U}{\partial z^2}$

**20.33.** Laplacian of  $\mathbf{A} = \nabla^2 \mathbf{A} = \frac{\partial^2 \mathbf{A}}{\partial x^2} + \frac{\partial^2 \mathbf{A}}{\partial y^2} + \frac{\partial^2 \mathbf{A}}{\partial z^2}$

### The Biharmonic Operator

---

**20.34.** Biharmonic operator on  $U = \nabla^4 U = \nabla^2(\nabla^2 U)$

$$= \frac{\partial^4 U}{\partial x^4} + \frac{\partial^4 U}{\partial y^4} + \frac{\partial^4 U}{\partial z^4} + 2 \frac{\partial^4 U}{\partial x^2 \partial y^2} + 2 \frac{\partial^4 U}{\partial y^2 \partial z^2} + 2 \frac{\partial^4 U}{\partial x^2 \partial z^2}$$

### Miscellaneous Formulas Involving $\nabla$

---

**20.35.**  $\nabla(U + V) = \nabla U + \nabla V$

**20.36.**  $\nabla \cdot (\mathbf{A} + \mathbf{B}) = \nabla \cdot \mathbf{A} + \nabla \cdot \mathbf{B}$

**20.37.**  $\nabla \times (\mathbf{A} + \mathbf{B}) = \nabla \times \mathbf{A} + \nabla \times \mathbf{B}$

**20.38.**  $\nabla \cdot (U\mathbf{A}) = (\nabla U) \cdot \mathbf{A} + U(\nabla \cdot \mathbf{A})$

**20.39.**  $\nabla \times (U\mathbf{A}) = (\nabla U) \times \mathbf{A} + U(\nabla \times \mathbf{A})$

**20.40.**  $\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B})$

**20.41.**  $\nabla \times (\mathbf{A} \times \mathbf{B}) = (\mathbf{B} \cdot \nabla) \mathbf{A} - \mathbf{B}(\nabla \cdot \mathbf{A}) - (\mathbf{A} \cdot \nabla) \mathbf{B} + \mathbf{A}(\nabla \cdot \mathbf{B})$

**20.42.**  $\nabla(\mathbf{A} \cdot \mathbf{B}) = (\mathbf{B} \cdot \nabla) \mathbf{A} + (\mathbf{A} \cdot \nabla) \mathbf{B} + \mathbf{B} \times (\nabla \times \mathbf{A}) + \mathbf{A} \times (\nabla \times \mathbf{B})$

**20.43.**  $\nabla \times (\nabla U) = 0$ , that is, the curl of the gradient of  $U$  is zero.

**20.44.**  $\nabla \cdot (\nabla \times \mathbf{A}) = 0$ , that is, the divergence of the curl of  $\mathbf{A}$  is zero.

**20.45.**  $\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}$

## Integrals Involving Vectors

If  $\mathbf{A}(u) = \frac{d}{du} \mathbf{B}(u)$ , then the *indefinite integral* of  $\mathbf{A}(u)$  is as follows:

$$20.46. \quad \int \mathbf{A}(u) du = \mathbf{B}(u) + \mathbf{c}, \quad c = \text{constant vector}$$

The *definite integral* of  $\mathbf{A}(u)$  from  $u = a$  to  $u = b$  in this case is given by

$$20.47. \quad \int_a^b \mathbf{A}(u) du = \mathbf{B}(b) - \mathbf{B}(a)$$

The definite integral can be defined as in 18.1.

## Line Integrals

Consider a space curve  $C$  joining two points  $P_1(a_1, a_2, a_3)$  and  $P_2(b_1, b_2, b_3)$  as in Fig. 20-6. Divide the curve into  $n$  parts by points of subdivision  $(x_1, y_1, z_1), \dots, (x_{n-1}, y_{n-1}, z_{n-1})$ . Then the *line integral* of a vector  $\mathbf{A}(x, y, z)$  along  $C$  is defined as

$$20.48. \quad \int_C \mathbf{A} \cdot d\mathbf{r} = \int_{P_1}^{P_2} \mathbf{A} \cdot d\mathbf{r} = \lim_{n \rightarrow \infty} \sum_{p=1}^n \mathbf{A}(x_p, y_p, z_p) \cdot \Delta \mathbf{r}_p$$




Fig. 20-6

where  $\Delta \mathbf{r}_p = \Delta x_p \mathbf{i} + \Delta y_p \mathbf{j} + \Delta z_p \mathbf{k}$ ,  $\Delta x_p = x_{p+1} - x_p$ ,  $\Delta y_p = y_{p+1} - y_p$ ,  $\Delta z_p = z_{p+1} - z_p$  and where it is assumed that as  $n \rightarrow \infty$  the largest of the magnitudes  $|\Delta \mathbf{r}_p|$  approaches zero. The result 20.48 is a generalization of the ordinary definite integral (see 18.1).

The line integral 20.48 can also be written as

$$20.49. \quad \int_C \mathbf{A} \cdot d\mathbf{r} = \int_C (A_1 dx + A_2 dy + A_3 dz)$$

using  $\mathbf{A} = A_1 \mathbf{i} + A_2 \mathbf{j} + A_3 \mathbf{k}$  and  $d\mathbf{r} = dx \mathbf{i} + dy \mathbf{j} + dz \mathbf{k}$ .

## Properties of Line Integrals

$$20.50. \quad \int_{P_1}^{P_2} \mathbf{A} \cdot d\mathbf{r} = - \int_{P_2}^{P_1} \mathbf{A} \cdot d\mathbf{r}$$

$$20.51. \quad \int_{P_1}^{P_2} \mathbf{A} \cdot d\mathbf{r} = \int_{P_1}^{P_3} \mathbf{A} \cdot d\mathbf{r} + \int_{P_3}^{P_2} \mathbf{A} \cdot d\mathbf{r}$$

## Independence of the Path

In general, a line integral has a value that depends on the particular path  $C$  joining points  $P_1$  and  $P_2$  in a region  $\mathcal{R}$ . However, in the case of  $\mathbf{A} = \nabla \phi$  or  $\nabla \times \mathbf{A} = 0$  where  $\phi$  and its partial derivatives are continuous in  $\mathcal{R}$ , the line integral  $\int_C \mathbf{A} \cdot d\mathbf{r}$  is independent of the path. In such a case,

$$20.52. \quad \int_C \mathbf{A} \cdot d\mathbf{r} = \int_{P_1}^{P_2} \mathbf{A} \cdot d\mathbf{r} = \phi(P_2) - \phi(P_1)$$

where  $\phi(P_1)$  and  $\phi(P_2)$  denote the values of  $\phi$  at  $P_1$  and  $P_2$ , respectively. In particular if  $C$  is a closed curve,

**20.53.**  $\int_C \mathbf{A} \cdot d\mathbf{r} = \oint_C \mathbf{A} \cdot d\mathbf{r} = 0$

where the circle on the integral sign is used to emphasize that  $C$  is closed.

## Multiple Integrals

Let  $F(x, y)$  be a function defined in a region  $\mathcal{R}$  of the  $xy$  plane as in Fig. 20-7. Subdivide the region into  $n$  parts by lines parallel to the  $x$  and  $y$  axes as indicated. Let  $\Delta A_p = \Delta x_p \Delta y_p$  denote an area of one of these parts. Then the integral of  $F(x, y)$  over  $\mathcal{R}$  is defined as

**20.54.**  $\int_{\mathcal{R}} F(x, y) dA = \lim_{n \rightarrow \infty} \sum_{p=1}^n F(x_p, y_p) \Delta A_p$

provided this limit exists.

In such a case, the integral can also be written as

$$\begin{aligned} \text{20.55. } & \int_{x=a}^b \int_{y=f_1(x)}^{f_2(x)} F(x, y) dy dx \\ & = \int_{x=a}^b \left\{ \int_{y=f_1(x)}^{f_2(x)} F(x, y) dy \right\} dx \end{aligned}$$

where  $y = f_1(x)$  and  $y = f_2(x)$  are the equations of curves  $PHQ$  and  $PGQ$ , respectively, and  $a$  and  $b$  are the  $x$  coordinates of points  $P$  and  $Q$ . The result can also be written as

**20.56.**  $\int_{y=c}^d \int_{x=g_1(y)}^{g_2(y)} F(x, y) dx dy = \int_{y=c}^d \left\{ \int_{x=g_1(y)}^{g_2(y)} F(x, y) dx \right\} dy$

where  $x = g_1(y)$ ,  $x = g_2(y)$  are the equations of curves  $HPG$  and  $HQG$ , respectively, and  $c$  and  $d$  are the  $y$  coordinates of  $H$  and  $G$ .

These are called *double integrals* or *area integrals*. The ideas can be similarly extended to *triple* or *volume integrals* or to higher *multiple integrals*.

## Surface Integrals

Subdivide the surface  $S$  (see Fig. 20-8) into  $n$  elements of area  $\Delta S_p$ ,  $p = 1, 2, \dots, n$ . Let  $\mathbf{A}(x_p, y_p, z_p) = \mathbf{A}_p$  where  $(x_p, y_p, z_p)$  is a point  $P$  in  $\Delta S_p$ . Let  $\mathbf{N}_p$  be a unit normal to  $\Delta S_p$  at  $P$ . Then the surface integral of the normal component of  $\mathbf{A}$  over  $S$  is defined as

**20.57.**  $\int_S \mathbf{A} \cdot \mathbf{N} dS = \lim_{n \rightarrow \infty} \sum_{p=1}^n \mathbf{A}_p \cdot \mathbf{N}_p \Delta S_p$




Fig. 20-7




Fig. 20-8

## Relation Between Surface and Double Integrals

If  $\mathcal{R}$  is the projection of  $S$  on the  $xy$  plane, then (see Fig. 20-8)

$$20.58. \int_S \mathbf{A} \cdot \mathbf{N} dS = \int_{\mathcal{R}} \int \mathbf{A} \cdot \mathbf{N} \frac{dx dy}{|\mathbf{N} \cdot \mathbf{k}|}$$

## The Divergence Theorem

Let  $S$  be a closed surface bounding a region of volume  $V$ ; and suppose  $\mathbf{N}$  is the positive (outward drawn) normal and  $d\mathbf{S} = \mathbf{N} dS$ . Then (see Fig. 20-9)

$$20.59. \int_V \nabla \cdot \mathbf{A} dV = \int_S \mathbf{A} \cdot d\mathbf{S}$$

The result is also called *Gauss' theorem* or *Green's theorem*.




Fig. 20-9




Fig. 20-10

## Stokes' Theorem

Let  $S$  be an open two-sided surface bounded by a closed non-intersecting curve  $C$  (simple closed curve) as in Fig. 20-10. Then

$$20.60. \oint_C \mathbf{A} \cdot d\mathbf{r} = \int_S (\nabla \times \mathbf{A}) \cdot d\mathbf{S}$$

where the circle on the integral is used to emphasize that  $C$  is closed.

## Green's Theorem in the Plane

$$20.61. \oint_C (P dx + Q dy) = \int_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

where  $R$  is the area bounded by the closed curve  $C$ . This result is a special case of the divergence theorem or Stokes' theorem.

### Green's First Identity

$$20.62. \int_V \{(\phi \nabla^2 \psi + (\nabla \phi) \cdot (\nabla \psi))\} dV = \int (\phi \nabla \psi) \cdot dS$$

where  $\phi$  and  $\psi$  are scalar functions.

### Green's Second Identity

$$20.63. \int_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV = \int_S (\phi \nabla \psi - \psi \nabla \phi) \cdot dS$$

### Miscellaneous Integral Theorems

$$20.64. \int_V \nabla \times \mathbf{A} dV = \int_S dS \times \mathbf{A}$$

$$20.65. \int_C \phi d\mathbf{r} = \int_S dS \times \nabla \phi$$

### Curvilinear Coordinates

A point  $P$  in space (see Fig. 20-11) can be located by rectangular coordinates  $(x, y, z)$  or curvilinear coordinates  $(u_1, u_2, u_3)$  where the transformation equations from one set of coordinates to the other are given by

$$20.66. \quad x = x(u_1, u_2, u_3)$$

$$y = y(u_1, u_2, u_3)$$

$$z = z(u_1, u_2, u_3)$$

If  $u_2$  and  $u_3$  are constant, then as  $u_1$  varies, the position vector  $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  of  $P$  describes a curve called the  $u_1$  coordinate curve. Similarly, we define the  $u_2$  and  $u_3$  coordinate curves through  $P$ . The vectors  $\frac{\partial \mathbf{r}}{\partial u_1}$ ,  $\frac{\partial \mathbf{r}}{\partial u_2}$ ,  $\frac{\partial \mathbf{r}}{\partial u_3}$  represent tangent vectors to the  $u_1$ ,  $u_2$ ,  $u_3$  coordinate curves. Letting  $\mathbf{e}_1$ ,  $\mathbf{e}_2$ ,  $\mathbf{e}_3$  be unit tangent vectors to these curves, we have

$$20.67. \quad \frac{\partial \mathbf{r}}{\partial u_1} = h_1 \mathbf{e}_1, \quad \frac{\partial \mathbf{r}}{\partial u_2} = h_2 \mathbf{e}_2, \quad \frac{\partial \mathbf{r}}{\partial u_3} = h_3 \mathbf{e}_3$$

where

$$20.68. \quad h_1 = \left| \frac{\partial \mathbf{r}}{\partial u_1} \right|, \quad h_2 = \left| \frac{\partial \mathbf{r}}{\partial u_2} \right|, \quad h_3 = \left| \frac{\partial \mathbf{r}}{\partial u_3} \right|$$

are called *scale factors*. If  $\mathbf{e}_1$ ,  $\mathbf{e}_2$ ,  $\mathbf{e}_3$  are mutually perpendicular, the curvilinear coordinate system is called *orthogonal*.




Fig. 20-11

### Formulas Involving Orthogonal Curvilinear Coordinates

---

$$20.69. \quad d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial u_1} du_1 + \frac{\partial \mathbf{r}}{\partial u_2} du_2 + \frac{\partial \mathbf{r}}{\partial u_3} du_3 = h_1 du_1 \mathbf{e}_1 + h_2 du_2 \mathbf{e}_2 + h_3 du_3 \mathbf{e}_3$$

$$20.70. \quad ds^2 = d\mathbf{r} \bullet d\mathbf{r} = h_1^2 du_1^2 + h_2^2 du_2^2 + h_3^2 du_3^2$$

where  $ds$  is the element of length.

If  $dV$  is the element of volume, then

$$20.71. \quad dV = |(h_1 \mathbf{e}_1 du_1) \bullet (h_2 \mathbf{e}_2 du_2) \times (h_3 \mathbf{e}_3 du_3)| = h_1 h_2 h_3 du_1 du_2 du_3$$

$$= \left| \frac{\partial \mathbf{r}}{\partial u_1} \bullet \frac{\partial \mathbf{r}}{\partial u_2} \times \frac{\partial \mathbf{r}}{\partial u_3} \right| du_1 du_2 du_3 = \left| \frac{\partial(x, y, z)}{\partial(u_1, u_2, u_3)} \right| du_1 du_2 du_3$$

where

$$20.72. \quad \frac{\partial(x, y, z)}{\partial(u_1, u_2, u_3)} = \begin{vmatrix} \frac{\partial x}{\partial u_1} & \frac{\partial x}{\partial u_2} & \frac{\partial x}{\partial u_3} \\ \frac{\partial y}{\partial u_1} & \frac{\partial y}{\partial u_2} & \frac{\partial y}{\partial u_3} \\ \frac{\partial z}{\partial u_1} & \frac{\partial z}{\partial u_2} & \frac{\partial z}{\partial u_3} \end{vmatrix}$$

sometimes written  $J(x, y, z; u_1, u_2, u_3)$ , is called the *Jacobian* of the transformation.

### Transformation of Multiple Integrals

---

Result 20.72 can be used to transform multiple integrals from rectangular to curvilinear coordinates. For example, we have

$$20.73. \quad \iiint_{\mathcal{R}} F(x, y, z) dx dy dz = \iiint_{\mathcal{R}'} G(u_1, u_2, u_3) \left| \frac{\partial(x, y, z)}{\partial(u_1, u_2, u_3)} \right| du_1 du_2 du_3$$

where  $\mathcal{R}'$  is the region into which  $\mathcal{R}$  is mapped by the transformation and  $G(u_1, u_2, u_3)$  is the value of  $F(x, y, z)$  corresponding to the transformation.

### Gradient, Divergence, Curl, and Laplacian

---

In the following,  $\Phi$  is a scalar function and  $\mathbf{A} = A_1 \mathbf{e}_1 + A_2 \mathbf{e}_2 + A_3 \mathbf{e}_3$  is a vector function of orthogonal curvilinear coordinates  $u_1, u_2, u_3$ .

$$20.74. \quad \text{Gradient of } \Phi = \text{grad } \Phi = \nabla \Phi = \frac{\mathbf{e}_1}{h_1} \frac{\partial \Phi}{\partial u_1} + \frac{\mathbf{e}_2}{h_2} \frac{\partial \Phi}{\partial u_2} + \frac{\mathbf{e}_3}{h_3} \frac{\partial \Phi}{\partial u_3}$$

$$20.75. \quad \text{Divergence of } \mathbf{A} = \text{div } \mathbf{A} = \nabla \bullet \mathbf{A} = \frac{1}{h_1 h_2 h_3} \left[ \frac{\partial}{\partial u_1} (h_2 h_3 A_1) + \frac{\partial}{\partial u_2} (h_3 h_1 A_2) + \frac{\partial}{\partial u_3} (h_1 h_2 A_3) \right]$$

$$20.76. \quad \text{Curl of } \mathbf{A} = \text{curl } \mathbf{A} = \nabla \times \mathbf{A} = \frac{1}{h_1 h_2 h_3} \begin{vmatrix} h_1 \mathbf{e}_1 & h_2 \mathbf{e}_2 & h_3 \mathbf{e}_3 \\ \frac{\partial}{\partial u_1} & \frac{\partial}{\partial u_2} & \frac{\partial}{\partial u_3} \\ h_1 A_1 & h_2 A_2 & h_3 A_3 \end{vmatrix}$$

$$= \frac{1}{h_2 h_3} \left[ \frac{\partial}{\partial u_2} (h_3 A_3) - \frac{\partial}{\partial u_3} (h_2 A_2) \right] \mathbf{e}_1 + \frac{1}{h_1 h_3} \left[ \frac{\partial}{\partial u_3} (h_1 A_1) - \frac{\partial}{\partial u_1} (h_3 A_3) \right] \mathbf{e}_2$$

$$+ \frac{1}{h_1 h_2} \left[ \frac{\partial}{\partial u_1} (h_2 A_2) - \frac{\partial}{\partial u_2} (h_1 A_1) \right] \mathbf{e}_3$$

**20.77.** Laplacian of  $\Phi = \nabla^2 \Phi = \frac{1}{h_1 h_2 h_3} \left[ \frac{\partial}{\partial u_1} \left( \frac{h_2 h_3}{h_1} \frac{\partial \Phi}{\partial u_1} \right) + \frac{\partial}{\partial u_2} \left( \frac{h_3 h_1}{h_2} \frac{\partial \Phi}{\partial u_2} \right) + \frac{\partial}{\partial u_3} \left( \frac{h_1 h_2}{h_3} \frac{\partial \Phi}{\partial u_3} \right) \right]$

Note that the biharmonic operator  $\nabla^4 \Phi = \nabla^2(\nabla^2 \Phi)$  can be obtained from 20.77.

## Special Orthogonal Coordinate Systems

### Cylindrical Coordinates $(r, \theta, z)$ (See Fig. 20-12)

**20.78.**  $x = r \cos \theta, \quad y = r \sin \theta, \quad z = z$

**20.79.**  $h_1^2 = 1, \quad h_2^2 = r^2, \quad h_3^2 = 1$

**20.80.**  $\nabla^2 \Phi = \frac{\partial^2 \Phi}{\partial r^2} + \frac{1}{r} \frac{\partial \Phi}{\partial r} + \frac{1}{r^2} \frac{\partial^2 \Phi}{\partial \theta^2} + \frac{\partial^2 \Phi}{\partial z^2}$




Fig. 20-12. Cylindrical coordinates.




Fig. 20-13. Spherical coordinates.

### Spherical Coordinates $(r, \theta, \phi)$ (See Fig. 20-13)

**20.81.**  $x = r \sin \theta \cos \phi, \quad y = r \sin \theta \sin \phi, \quad z = r \cos \theta$

**20.82.**  $h_1^2 = 1, \quad h_2^2 = r^2, \quad h_3^2 = r^2 \sin^2 \theta$

**20.83.**  $\nabla^2 \Phi = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \Phi}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial \Phi}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2 \Phi}{\partial \phi^2}$

### Parabolic Cylindrical Coordinates $(u, v, z)$

**20.84.**  $x = \frac{1}{2}(u^2 - v^2), \quad y = uv, \quad z = z$

**20.85.**  $h_1^2 = h_2^2 = u^2 + v^2, \quad h_3^2 = 1$

**20.86.**  $\nabla^2 \Phi = \frac{1}{u^2 + v^2} \left( \frac{\partial^2 \Phi}{\partial u^2} + \frac{\partial^2 \Phi}{\partial v^2} \right) + \frac{\partial^2 \Phi}{\partial z^2}$

The traces of the coordinate surfaces on the  $xy$  plane are shown in Fig. 20-14. They are confocal parabolas with a common axis.




Fig. 20-14

### Paraboloidal Coordinates ( $u, v, \phi$ )

**20.87.**  $x = uv \cos \phi, \quad y = uv \sin \phi, \quad z = \frac{1}{2}(u^2 - v^2)$

where  $u \geq 0, \quad v \geq 0, \quad 0 \leq \phi < 2\pi$

**20.88.**  $h_1^2 = h_2^2 = u^2 + v^2, \quad h_3^2 = u^2 v^2$

**20.89.**  $\nabla^2 \Phi = \frac{1}{u(u^2 + v^2)} \frac{\partial}{\partial u} \left( u \frac{\partial \Phi}{\partial u} \right) + \frac{1}{v(u^2 + v^2)} \frac{\partial}{\partial v} \left( v \frac{\partial \Phi}{\partial v} \right) + \frac{1}{u^2 v^2} \frac{\partial^2 \Phi}{\partial \phi^2}$

Two sets of coordinate surfaces are obtained by revolving the parabolas of Fig. 20-14 about the  $x$  axis which is then relabeled the  $z$  axis.

### Elliptic Cylindrical Coordinates ( $u, v, z$ )

**20.90.**  $x = a \cosh u \cos v, \quad y = a \sinh u \sin v, \quad z = z$

where  $u \geq 0, \quad 0 \leq v < 2\pi, \quad -\infty < z < \infty$

**20.91.**  $h_1^2 = h_2^2 = a^2(\sinh^2 u + \sin^2 v), \quad h_3^2 = 1$

**20.92.**  $\nabla^2 \Phi = \frac{1}{a^2(\sinh^2 u + \sin^2 v)} \left( \frac{\partial^2 \Phi}{\partial u^2} + \frac{\partial^2 \Phi}{\partial v^2} \right) + \frac{\partial^2 \Phi}{\partial z^2}$

The traces of the coordinate surfaces on the  $xy$  plane are shown in Fig. 20-15. They are confocal ellipses and hyperbolas.




Fig. 20-15. Elliptic cylindrical coordinates.

### Prolate Spheroidal Coordinates ( $\xi, \eta, \phi$ )

**20.93.**  $x = a \sinh \xi \sin \eta \cos \phi, \quad y = a \sinh \xi \sin \eta \sin \phi, \quad z = a \cosh \xi \cos \eta$

where

$$\xi \geq 0, \quad 0 \leq \eta \leq \pi, \quad 0 \leq \phi < 2\pi$$

**20.94.**  $h_1^2 = h_2^2 = a^2(\sinh^2 \xi \sin^2 \eta), \quad h_3^2 = a^2 \sinh^2 \xi \sin^2 \eta$

**20.95.** 
$$\nabla^2 \Phi = \frac{1}{a^2(\sinh^2 \xi + \sin^2 \eta) \sinh \xi} \frac{\partial}{\partial \xi} \left( \sinh \xi \frac{\partial \Phi}{\partial \xi} \right)$$

$$+ \frac{1}{a^2(\sinh^2 \xi + \sin^2 \eta) \sin \eta} \frac{\partial}{\partial \eta} \left( \sin \eta \frac{\partial \Phi}{\partial \eta} \right) + \frac{1}{a^2 \sinh^2 \xi \sin^2 \eta} \frac{\partial^2 \Phi}{\partial \phi^2}$$

Two sets of coordinate surfaces are obtained by revolving the curves of Fig. 20-15 about the  $x$  axis which is relabeled the  $z$  axis. The third set of coordinate surfaces consists of planes passing through this axis.

### Oblate Spheroidal Coordinates ( $\xi, \eta, \phi$ )

**20.96.**  $x = a \cosh \xi \cos \eta \cos \phi, \quad y = a \cosh \xi \cos \eta \sin \phi, \quad z = a \sinh \xi \sin \eta$

where

$$\xi \geq 0, \quad -\pi/2 \leq \eta \leq \pi/2, \quad 0 \leq \phi < 2\pi$$

**20.97.**  $h_1^2 = h_2^2 = a^2(\sinh^2 \xi + \sin^2 \eta), \quad h_3^2 = a^2 \cosh^2 \xi \cos^2 \eta$

**20.98.** 
$$\nabla^2 \Phi = \frac{1}{a^2(\sinh^2 \xi + \sin^2 \eta) \cosh \xi} \frac{\partial}{\partial \xi} \left( \cosh \xi \frac{\partial \Phi}{\partial \xi} \right)$$

$$+ \frac{1}{a^2(\sinh^2 \xi + \sin^2 \eta) \cos \eta} \frac{\partial}{\partial \eta} \left( \cos \eta \frac{\partial \Phi}{\partial \eta} \right) + \frac{1}{a^2 \cosh^2 \xi \cos^2 \eta} \frac{\partial^2 \Phi}{\partial \phi^2}$$

Two sets of coordinate surfaces are obtained by revolving the curves of Fig. 20-15 about the  $y$  axis which is relabeled the  $z$  axis. The third set of coordinate surfaces are planes passing through this axis.

### Bipolar Coordinates ( $u, v, z$ )

**20.99.**  $x = \frac{a \sinh v}{\cosh v - \cos u}, \quad y = \frac{a \sin u}{\cosh v - \cos u}, \quad z = z$

where

$$0 \leq u < 2\pi, \quad -\infty < v < \infty, \quad -\infty < z < \infty$$

or

**20.100.**  $x^2 + (y - a \cot u)^2 = a^2 \csc^2 u, \quad (x - a \coth v)^2 + y^2 = a^2 \operatorname{csch}^2 v, \quad z = z$

**20.101.**  $h_1^2 = h_2^2 = \frac{a^2}{(\cosh v - \cos u)^2}, \quad h_3^2 = 1$

**20.102.**  $\nabla^2 \Phi = \frac{(\cosh v - \cos u)^2}{a^2} \left( \frac{\partial^2 \Phi}{\partial u^2} + \frac{\partial^2 \Phi}{\partial v^2} \right) + \frac{\partial^2 \Phi}{\partial z^2}$

The traces of the coordinate surfaces on the  $xy$  plane are shown in Fig. 20-16.




Fig. 20-16. Bipolar coordinates.

#### Toroidal Coordinates ( $u, v, \phi$ )

**20.103.**  $x = \frac{a \sinh v \cos \phi}{\cosh v - \cos u}, \quad y = \frac{a \sinh v \sin \phi}{\cosh v - \cos u}, \quad z = \frac{a \sin u}{\cosh v - \cos u}$

**20.104.**  $h_1^2 = h_2^2 = \frac{a^2}{(\cosh v - \cos u)^2}, \quad h_3^2 = \frac{a^2 \sinh^2 v}{(\cosh v - \cos u)^2}$

**20.105.**  $\nabla^2 \Phi = \frac{(\cosh v - \cos u)^3}{a^2} \frac{\partial}{\partial u} \left( \frac{1}{\cosh v - \cos u} \frac{\partial \Phi}{\partial u} \right)$   
 $+ \frac{(\cosh v - \cos u)^3}{a^2 \sinh v} \frac{\partial}{\partial v} \left( \frac{\sinh v}{\cosh v - \cos u} \frac{\partial \Phi}{\partial v} \right) + \frac{(\cosh v - \cos u)^2}{a^2 \sinh^2 v} \frac{\partial^2 \Phi}{\partial \phi^2}$

The coordinate surfaces are obtained by revolving the curves of Fig. 20.16 about the  $y$  axis which is relabeled the  $z$  axis.

#### Conical Coordinates ( $\lambda, \mu, \nu$ )

**20.106.**  $x = \frac{\lambda \mu v}{ab}, \quad y = \frac{\lambda}{a} \sqrt{\frac{(\mu^2 - a^2)(v^2 - a^2)}{a^2 - b^2}}, \quad z = \frac{\lambda}{b} \sqrt{\frac{(\mu^2 - b^2)(v^2 - b^2)}{b^2 - a^2}}$

**20.107.**  $h_1^2 = 1, \quad h_2^2 = \frac{\lambda^2(\mu^2 - v^2)}{(\mu^2 - a^2)(b^2 - \mu^2)}, \quad h_3^2 = \frac{\lambda^2(\mu^2 - v^2)}{(v^2 - a^2)(v^2 - b^2)}$

Confocal Ellipsoidal Coordinates  $(\lambda, \mu, v)$ 

$$20.108. \quad \begin{cases} \frac{x^2}{a^2 - \lambda} + \frac{y^2}{b^2 - \lambda} + \frac{z^2}{c^2 - \lambda} = 1, & \lambda < c^2 < b^2 < a^2 \\ \frac{x^2}{a^2 - \mu} + \frac{y^2}{b^2 - \mu} + \frac{z^2}{c^2 - \mu} = 1, & c^2 < \mu < b^2 < a^2 \\ \frac{x^2}{a^2 - v} + \frac{y^2}{b^2 - v} + \frac{z^2}{c^2 - v} = 1, & c^2 < b^2 < v < a^2 \end{cases}$$

or

$$20.109. \quad \begin{cases} x^2 = \frac{(a^2 - \lambda)(a^2 - \mu)(a^2 - v)}{(a^2 - b^2)(a^2 - c^2)} \\ y^2 = \frac{(b^2 - \lambda)(b^2 - \mu)(b^2 - v)}{(b^2 - a^2)(b^2 - c^2)} \\ z^2 = \frac{(c^2 - \lambda)(c^2 - \mu)(c^2 - v)}{(c^2 - a^2)(c^2 - b^2)} \end{cases}$$

$$20.110. \quad \begin{cases} h_1^2 = \frac{(\mu - \lambda)(v - \lambda)}{4(a^2 - \lambda)(b^2 - \lambda)(c^2 - \lambda)} \\ h_2^2 = \frac{(v - \mu)(\lambda - \mu)}{4(a^2 - \mu)(b^2 - \mu)(c^2 - \mu)} \\ h_3^2 = \frac{(\lambda - v)(\mu - v)}{4(a^2 - v)(b^2 - v)(c^2 - v)} \end{cases}$$

Confocal Paraboloidal Coordinates  $(\lambda, \mu, v)$ 

$$20.111. \quad \begin{cases} \frac{x^2}{a^2 - \lambda} + \frac{h^2}{b^2 - \lambda} = z - \lambda, & -\infty < \lambda < b^2 \\ \frac{x^2}{a^2 - \mu} + \frac{y^2}{b^2 - \mu} = z - \mu, & b^2 < \mu < a^2 \\ \frac{x^2}{a^2 - v} + \frac{y^2}{b^2 - v} = z - v, & a^2 < v < \infty \end{cases}$$

or

$$20.112. \quad \begin{cases} x^2 = \frac{(a^2 - \lambda)(a^2 - \mu)(a^2 - v)}{b^2 - a^2} \\ y^2 = \frac{(b^2 - \lambda)(b^2 - \mu)(b^2 - v)}{a^2 - b^2} \\ z = \lambda + \mu + v - a^2 - b^2 \end{cases}$$

$$20.113. \quad \begin{cases} h_1^2 = \frac{(\mu - \lambda)(v - \lambda)}{4(a^2 - \lambda)(b^2 - \lambda)} \\ h_2^2 = \frac{(v - \mu)(\lambda - \mu)}{4(a^2 - \mu)(b^2 - \mu)} \\ h_3^2 = \frac{(\lambda - v)(\mu - v)}{16(a^2 - v)(b^2 - v)} \end{cases}$$

## Section VI: Series

# 21 SERIES of CONSTANTS

### Arithmetic Series

$$21.1. \quad a + (a+d) + (a+2d) + \cdots + \{a + (n-1)d\} = \frac{1}{2}n\{2a + (n-1)d\} = \frac{1}{2}n(a+l)$$

where  $l = a + (n-1)d$  is the last term.

Some special cases are

$$21.2. \quad 1 + 2 + 3 + \cdots + n = \frac{1}{2}n(n+1)$$

$$21.3. \quad 1 + 3 + 5 + \cdots + (2n-1) = n^2$$

### Geometric Series

$$21.4. \quad a + ar + ar^2 + ar^3 + \cdots + ar^{n-1} = \frac{a(1-r^n)}{1-r} = \frac{a - rl}{1-r}$$

where  $l = ar^{n-1}$  is the last term and  $r \neq 1$ .

If  $-1 < r < 1$ , then

$$21.5. \quad a + ar + ar^2 + ar^3 + \cdots = \frac{a}{1-r}$$

### Arithmetic-Geometric Series

$$21.6. \quad a + (a+d)r + (a+2d)r^2 + \cdots + \{a + (n-1)d\}r^{n-1} = \frac{a(1-r^n)}{1-r} + \frac{rd\{1-nr^{n-1} + (n-1)r^n\}}{(1-r)^2}$$

where  $r \neq 1$ .

If  $-1 < r < 1$ , then

$$21.7. \quad a + (a+d)r + (d+2d)r^2 + \cdots = \frac{a}{1-r} + \frac{rd}{(1-r)^2}$$

### Sums of Powers of Positive Integers

$$21.8. \quad 1^p + 2^p + 3^p + \cdots + n^p = \frac{n^{p+1}}{p+1} + \frac{1}{2}n^p + \frac{B_1 p n^{p-1}}{2!} - \frac{B_2 p(p-1)(p-2)n^{p-3}}{4!} + \cdots$$

where the series terminates at  $n^2$  or  $n$  according as  $p$  is odd or even, and  $B_k$  are the *Bernoulli numbers* (see page 142).

Some special cases are

$$21.9. \quad 1+2+3+\cdots+n = \frac{n(n+1)}{2}$$

$$21.10. \quad 1^2 + 2^2 + 3^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}$$

$$21.11. \quad 1^3 + 2^3 + 3^3 + \cdots + n^3 = \frac{n^2(n+1)^2}{4} = (1+2+3+\cdots+n)^2$$

$$21.12. \quad 1^4 + 2^4 + 3^4 + \cdots + n^4 = \frac{n(n+1)(2n+1)(3n^2+3n-1)}{30}$$

If  $S_k = 1^k + 2^k + 3^k + \cdots + n^k$  where  $k$  and  $n$  are positive integers, then

$$21.13. \quad \binom{k+1}{1} S_1 + \binom{k+1}{2} S_2 + \cdots + \binom{k+1}{k} S_k = (n+1)^{k+1} - (n+1)$$

### **Series Involving Reciprocals of Powers of Positive Integers**

---

$$21.14. \quad 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \cdots = \ln 2$$

$$21.15. \quad 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots = \frac{\pi}{4}$$

$$21.16. \quad 1 - \frac{1}{4} + \frac{1}{7} - \frac{1}{10} + \frac{1}{13} - \cdots = \frac{\pi\sqrt{3}}{9} + \frac{1}{3}\ln 2$$

$$21.17. \quad 1 - \frac{1}{5} + \frac{1}{9} - \frac{1}{13} + \frac{1}{17} - \cdots = \frac{\pi\sqrt{2}}{8} + \frac{\sqrt{2}\ln(1+\sqrt{2})}{4}$$

$$21.18. \quad \frac{1}{2} - \frac{1}{5} + \frac{1}{8} - \frac{1}{11} + \frac{1}{14} - \cdots = \frac{\pi\sqrt{3}}{9} + \frac{1}{3}\ln 2$$

$$21.19. \quad \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots = \frac{\pi^2}{6}$$

$$21.20. \quad \frac{1}{1^4} + \frac{1}{2^4} + \frac{1}{3^4} + \frac{1}{4^4} + \cdots = \frac{\pi^4}{90}$$

$$21.21. \quad \frac{1}{1^6} + \frac{1}{2^6} + \frac{1}{3^6} + \frac{1}{4^6} + \cdots = \frac{\pi^6}{945}$$

$$21.22. \quad \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \cdots = \frac{\pi^2}{12}$$

$$21.23. \quad \frac{1}{1^4} - \frac{1}{2^4} + \frac{1}{3^4} - \frac{1}{4^4} + \cdots = \frac{7\pi^4}{720}$$

$$21.24. \quad \frac{1}{1^6} - \frac{1}{2^6} + \frac{1}{3^6} - \frac{1}{4^6} + \cdots = \frac{31\pi^6}{30,240}$$

$$21.25. \quad \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \cdots = \frac{\pi^2}{8}$$

$$21.26. \quad \frac{1}{1^4} + \frac{1}{3^4} + \frac{1}{5^4} + \frac{1}{7^4} + \cdots = \frac{\pi^4}{96}$$

21.27.  $\frac{1}{1^6} + \frac{1}{3^6} + \frac{1}{5^6} + \frac{1}{7^6} + \dots = \frac{\pi^6}{960}$

21.28.  $\frac{1}{1^3} - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \dots = \frac{\pi^3}{32}$

21.29.  $\frac{1}{1^3} + \frac{1}{3^3} - \frac{1}{5^3} - \frac{1}{7^3} + \dots = \frac{3\pi^3\sqrt{2}}{128}$

21.30.  $\frac{1}{1 \cdot 3} + \frac{1}{3 \cdot 5} + \frac{1}{5 \cdot 7} + \frac{1}{7 \cdot 9} + \dots = \frac{1}{2}$

21.31.  $\frac{1}{1 \cdot 3} + \frac{1}{2 \cdot 4} + \frac{1}{3 \cdot 5} + \frac{1}{4 \cdot 6} + \dots = \frac{3}{4}$

21.32.  $\frac{1}{1^2 \cdot 3^2} + \frac{1}{3^2 \cdot 5^2} + \frac{1}{5^2 \cdot 7^2} + \frac{1}{7^2 \cdot 9^2} + \dots = \frac{\pi^2 - 8}{16}$

21.33.  $\frac{1}{1^2 \cdot 2^2 \cdot 3^2} + \frac{1}{2^2 \cdot 3^2 \cdot 4^2} + \frac{1}{3^2 \cdot 4^2 \cdot 5^2} + \dots = \frac{4\pi^2 - 39}{16}$

21.34.  $\frac{1}{a} - \frac{1}{a+d} + \frac{1}{a+2d} - \frac{1}{a+3d} + \dots = \int_0^1 \frac{u^{a-1} du}{1+u^d}$

21.35.  $\frac{1}{1^{2p}} + \frac{1}{2^{2p}} + \frac{1}{3^{2p}} + \frac{1}{4^{2p}} + \dots = \frac{2^{2p-1} \pi^{2p} B_p}{(2p)!}$

21.36.  $\frac{1}{1^{2p}} + \frac{1}{3^{2p}} + \frac{1}{5^{2p}} + \frac{1}{7^{2p}} + \dots = \frac{(2^{2p}-1) \pi^{2p} B_p}{2(2p)!}$

21.37.  $\frac{1}{1^{2p}} - \frac{1}{2^{2p}} + \frac{1}{3^{2p}} - \frac{1}{4^{2p}} + \dots = \frac{(2^{2p-1}-1) \pi^{2p} B_p}{(2p)!}$

21.38.  $\frac{1}{1^{2p+1}} - \frac{1}{3^{2p+1}} + \frac{1}{5^{2p+1}} - \frac{1}{7^{2p+1}} + \dots = \frac{\pi^{2p+1} E_p}{2^{2p+2}(2p)!}$

### Miscellaneous Series

21.39.  $\frac{1}{2} + \cos \alpha + \cos 2\alpha + \dots + \cos n\alpha = \frac{\sin(n+1/2)\alpha}{2 \sin(\alpha/2)}$

21.40.  $\sin \alpha + \sin 2\alpha + \sin 3\alpha + \dots + \sin n\alpha = \frac{\sin[1/2(n+1)]\alpha \sin 1/2n\alpha}{\sin(\alpha/2)}$

21.41.  $1 + r \cos \alpha + r^2 \cos 2\alpha + r^3 \cos 3\alpha + \dots = \frac{1 - r \cos \alpha}{1 - 2r \cos \alpha + r^2}, |r| < 1$

21.42.  $r \sin \alpha + r^2 \sin 2\alpha + r^3 \sin 3\alpha + \dots = \frac{r \sin \alpha}{1 - 2r \cos \alpha + r^2}, |r| < 1$

21.43.  $1 + r \cos \alpha + r^2 \cos 2\alpha + \dots + r^n \cos n\alpha = \frac{r^{n+2} \cos n\alpha - r^{n+1} \cos(n+1)\alpha - r \cos \alpha + 1}{1 - 2r \cos \alpha + r^2}$

21.44.  $r \sin \alpha + r^2 \sin 2\alpha + \dots + r^n \sin n\alpha = \frac{r \sin \alpha - r^{n+1} \sin(n+1)\alpha + r^{n+2} \sin n\alpha}{1 - 2r \cos \alpha + r^2}$

**The Euler-Maclaurin Summation Formula**

---

**21.45.** 
$$\sum_{k=1}^{n-1} F(k) = \int_0^n F(k) dk - \frac{1}{2}\{F(0) + F(n)\}$$

$$+ \frac{1}{12}\{F'(n) - F(0)\} - \frac{1}{720}\{F'''(n) - F'''(0)\}$$

$$+ \frac{1}{30,240}\{F^{(v)}(n) - F^{(v)}(0)\} - \frac{1}{1,209,600}\{F^{(vii)}(n) - F^{(vii)}(0)\}$$

$$+ \cdots (-1)^{p-1} \frac{B_p}{(2p)!} \{F^{(2p-1)}(n) - F^{(2p-1)}(0)\} + \cdots$$

**The Poisson Summation Formula**

---

**21.46.** 
$$\sum_{k=-\infty}^{\infty} F(k) = \sum_{m=-\infty}^{\infty} \left\{ \int_{-\infty}^{\infty} e^{2\pi i mx} F(x) dx \right\}$$

# 22

## TAYLOR SERIES

### Taylor Series for Functions of One Variable

$$22.1. \quad f(x) = f(a) + f'(a)(x-a) + \frac{f''(a)(x-a)^2}{2!} + \cdots + \frac{f^{(n-1)}(a)(x-a)^{n-1}}{(n-1)!} + R_n$$

where  $R_n$ , the remainder after  $n$  terms, is given by either of the following forms:

$$22.2. \quad \text{Lagrange's form: } R_n = \frac{f^{(n)}(\xi)(x-a)^n}{n!}$$

$$22.3. \quad \text{Cauchy's form: } R_n = \frac{f^{(n)}(\xi)(x-\xi)^{n-1}(x-a)}{(n-1)!}$$

The value  $\xi$ , which may be different in the two forms, lies between  $a$  and  $x$ . The result holds if  $f(x)$  has continuous derivatives of order  $n$  at least.

If  $\lim_{n \rightarrow \infty} R_n = 0$ , the infinite series obtained is called the *Taylor series* for  $f(x)$  about  $x = a$ . If  $a = 0$ , the series is often called a *Maclaurin series*. These series, often called power series, generally converge for all values of  $x$  in some interval called the *interval of convergence* and diverge for all  $x$  outside this interval.

Some series contain the Bernoulli numbers  $B_n$  and the Euler numbers  $E_n$  defined in Chapter 23, pages 142–143.

### Binomial Series

$$22.4. \quad (a+x)^n = a^n + na^{n-1}x + \frac{n(n-1)}{2!}a^{n-2}x^2 + \frac{n(n-1)(n-2)}{3!}a^{n-3}x^3 + \cdots \\ = a^n + \binom{n}{1}a^{n-1}x + \binom{n}{2}a^{n-2}x^2 + \binom{n}{3}a^{n-3}x^3 + \cdots$$

Special cases are

$$22.5. \quad (a+x)^2 = a^2 + 2ax + x^2$$

$$22.6. \quad (a+x)^3 = a^3 + 3a^2x + 3ax^2 + x^3$$

$$22.7. \quad (a+x)^4 = a^4 + 4a^3x + 6a^2x^2 + 4ax^3 + x^4$$

$$22.8. \quad (1+x)^{-1} = 1 - x + x^2 - x^3 + x^4 - \cdots \quad -1 < x < 1$$

$$22.9. \quad (1+x)^{-2} = 1 - 2x + 3x^2 - 4x^3 + 5x^4 - \cdots \quad -1 < x < 1$$

$$22.10. \quad (1+x)^{-3} = 1 - 3x + 6x^2 - 10x^3 + 15x^4 - \cdots \quad -1 < x < 1$$

- 22.11.**  $(1+x)^{-1/2} = 1 - \frac{1}{2}x + \frac{1 \cdot 3}{2 \cdot 4}x^2 - \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6}x^3 + \dots$   $-1 < x \leq 1$
- 22.12.**  $(1+x)^{1/2} = 1 + \frac{1}{2}x - \frac{1}{2 \cdot 4}x^2 + \frac{1 \cdot 3}{2 \cdot 4 \cdot 6}x^3 - \dots$   $-1 < x \leq 1$
- 22.13.**  $(1+x)^{-1/3} = 1 - \frac{1}{3}x + \frac{1 \cdot 4}{3 \cdot 6}x^2 - \frac{1 \cdot 4 \cdot 7}{3 \cdot 6 \cdot 9}x^3 + \dots$   $-1 < x \leq 1$
- 22.14.**  $(1+x)^{1/3} = 1 + \frac{1}{3}x - \frac{2}{3 \cdot 6}x^2 + \frac{2 \cdot 5}{3 \cdot 6 \cdot 9}x^3 - \dots$   $-1 < x \leq 1$

### Series for Exponential and Logarithmic Functions

---

- 22.15.**  $e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$   $-\infty < x < \infty$
- 22.16.**  $a^x = e^{x \ln a} = 1 + x \ln a + \frac{(x \ln a)^2}{2!} + \frac{(x \ln a)^3}{3!} + \dots$   $-\infty < x < \infty$
- 22.17.**  $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$   $-1 < x \leq 1$
- 22.18.**  $\frac{1}{2} \ln\left(\frac{1+x}{1-x}\right) = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \dots$   $-1 < x < 1$
- 22.19.**  $\ln x = 2 \left\{ \left( \frac{x-1}{x+1} \right) + \frac{1}{3} \left( \frac{x-1}{x+1} \right)^3 + \frac{1}{5} \left( \frac{x-1}{x+1} \right)^5 + \dots \right\}$   $x > 0$
- 22.20.**  $\ln x = \left( \frac{x-1}{x} \right) + \frac{1}{2} \left( \frac{x-1}{x} \right)^2 + \frac{1}{3} \left( \frac{x-1}{x} \right)^3 + \dots$   $x \geq \frac{1}{2}$

### Series for Trigonometric Functions

---

- 22.21.**  $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$   $-\infty < x < \infty$
- 22.22.**  $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$   $-\infty < x < \infty$
- 22.23.**  $\tan x = x + \frac{x^3}{3} + \frac{2x^5}{15} + \frac{17x^7}{315} + \dots + \frac{2^{2n}(2^{2n}-1)B_n x^{2n-1}}{(2n)!} + \dots$   $|x| < \frac{\pi}{2}$
- 22.24.**  $\cot x = \frac{1}{x} - \frac{x}{3} - \frac{x^3}{45} - \frac{2x^5}{945} - \dots - \frac{2^{2n} B_n x^{2n-1}}{(2n)!} - \dots$   $0 < |x| < \pi$
- 22.25.**  $\sec x = 1 + \frac{x^2}{2} + \frac{5x^4}{24} + \frac{61x^6}{720} + \dots + \frac{E_n x^{2n}}{(2n)!} + \dots$   $|x| < \frac{\pi}{2}$
- 22.26.**  $\csc x = \frac{1}{x} + \frac{x}{6} + \frac{7x^3}{360} + \frac{31x^5}{15,120} + \dots + \frac{2(2^{2n-1}-1)B_n x^{2n-1}}{(2n)!} + \dots$   $0 < |x| < \pi$
- 22.27.**  $\sin^{-1} x = x + \frac{1}{2} \frac{x^3}{3} + \frac{1 \cdot 3}{2 \cdot 4} \frac{x^5}{5} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \frac{x^7}{7} + \dots$   $|x| < 1$
- 22.28.**  $\cos^{-1} x = \frac{\pi}{2} - \sin^{-1} x = \frac{\pi}{2} - \left( x + \frac{1}{2} \frac{x^3}{3} + \frac{1 \cdot 3}{2 \cdot 4} \frac{x^5}{5} + \dots \right)$   $|x| < 1$

- 22.29.  $\tan^{-1} x = \begin{cases} x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots & |x| < 1 \\ \pm \frac{\pi}{2} - \frac{1}{x} + \frac{1}{3x^3} - \frac{1}{5x^5} + \dots & (+ \text{ if } x \geq 1, - \text{ if } x \leq -1) \end{cases}$
- 22.30.  $\cot^{-1} x = \frac{\pi}{2} - \tan^{-1} x = \begin{cases} \frac{\pi}{2} - \left( x - \frac{x^3}{3} + \frac{x^5}{5} - \dots \right) & |x| < 1 \\ p\pi + \frac{1}{x} - \frac{1}{3x^3} + \frac{1}{5x^5} - \dots & (p = 0 \text{ if } x > 1, p = 1 \text{ if } x < -1) \end{cases}$
- 22.31.  $\sec^{-1} x = \cos^{-1}(1/x) = \frac{\pi}{2} - \left( \frac{1}{x} + \frac{1}{2 \cdot 3x^3} + \frac{1 \cdot 3}{2 \cdot 4 \cdot 5x^5} + \dots \right)$   $|x| > 1$
- 22.32.  $\csc^{-1} x = \sin^{-1}(1/x) = \frac{1}{x} + \frac{1}{2 \cdot 3x^3} + \frac{1 \cdot 3}{2 \cdot 4 \cdot 5x^5} + \dots$   $|x| > 1$

### Series for Hyperbolic Functions

---

- 22.33.  $\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \frac{x^7}{7!} + \dots$   $-\infty < x < \infty$
- 22.34.  $\cosh x = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \frac{x^6}{6!} + \dots$   $-\infty < x < \infty$
- 22.35.  $\tanh x = x - \frac{x^3}{3} + \frac{2x^5}{15} - \frac{17x^7}{315} + \dots \frac{(-1)^{n-1} 2^{2n} (2^{2n}-1) B_n x^{2n-1}}{(2n)!} + \dots$   $|x| < \frac{\pi}{2}$
- 22.36.  $\coth x = \frac{1}{x} + \frac{x}{3} - \frac{x^3}{45} + \frac{2x^5}{945} + \dots \frac{(-1)^{n-1} 2^{2n} B_n x^{2n-1}}{(2n)!} + \dots$   $0 < |x| < \pi$
- 22.37.  $\operatorname{sech} x = 1 - \frac{x^2}{2} + \frac{5x^4}{24} - \frac{61x^6}{720} + \dots \frac{(-1)^n E_n x^{2n}}{(2n)!} + \dots$   $|x| < \frac{\pi}{2}$
- 22.38.  $\operatorname{csch} x = \frac{1}{x} - \frac{x}{6} + \frac{7x^3}{360} - \frac{31x^5}{15,120} + \dots \frac{(-1)^n 2(2^{2n-1}-1) B_n x^{2n-1}}{(2n)!} + \dots$   $0 < |x| < \pi$
- 22.39.  $\sinh^{-1} x = \begin{cases} x - \frac{x^3}{2 \cdot 3} + \frac{1 \cdot 3x^5}{2 \cdot 4 \cdot 5} - \frac{1 \cdot 3 \cdot 5x^7}{2 \cdot 4 \cdot 6 \cdot 7} + \dots & |x| < 1 \\ \pm \left( \ln |2x| + \frac{1}{2 \cdot 2x^2} - \frac{1 \cdot 3}{2 \cdot 4 \cdot 4x^4} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 6x^6} - \dots \right) & \begin{bmatrix} + \text{ if } x \geq 1 \\ - \text{ if } x \leq -1 \end{bmatrix} \end{cases}$
- 22.40.  $\cosh^{-1} x = \pm \left\{ \ln(2x) - \left( \frac{1}{2 \cdot 2x^2} + \frac{1 \cdot 3}{2 \cdot 4 \cdot 4x^4} + \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6 \cdot 6x^6} + \dots \right) \right\}$   $\begin{bmatrix} + \text{ if } \cosh^{-1} x > 0, x \geq 1 \\ - \text{ if } \cosh^{-1} x < 0, x \geq 1 \end{bmatrix}$
- 22.41.  $\tanh^{-1} x = x + \frac{x^3}{3} + \frac{x^5}{5} + \frac{x^7}{7} + \dots$   $|x| < 1$
- 22.42.  $\coth^{-1} x = \frac{1}{x} + \frac{1}{3x^3} + \frac{1}{5x^5} + \frac{1}{7x^7} + \dots$   $|x| > 1$

### Miscellaneous Series

---

- 22.43.  $e^{\sin x} = 1 + x + \frac{x^2}{2} - \frac{x^4}{8} - \frac{x^5}{15} + \dots$   $-\infty < x < \infty$
- 22.44.  $e^{\cos x} = e \left( 1 - \frac{x^2}{2} + \frac{x^4}{6} - \frac{31x^6}{720} + \dots \right)$   $-\infty < x < \infty$

- 22.45.**  $e^{\tan x} = 1 + x + \frac{x^2}{2} + \frac{x^3}{2} + \frac{3x^4}{8} + \dots$   $|x| < \frac{\pi}{2}$
- 22.46.**  $e^x \sin x = x + x^2 + \frac{x^3}{3} - \frac{x^5}{30} - \frac{x^6}{90} + \dots + \frac{2^{n/2} \sin(n\pi/4)x^n}{n!} + \dots$   $-\infty < x < \infty$
- 22.47.**  $e^x \cos x = 1 + x - \frac{x^3}{3} - \frac{x^4}{6} + \dots + \frac{2^{n/2} \cos(n\pi/4)x^n}{n!} + \dots$   $-\infty < x < \infty$
- 22.48.**  $\ln |\sin x| = \ln |x| - \frac{x^2}{6} - \frac{x^4}{180} - \frac{x^6}{2835} - \dots - \frac{2^{2n-1} B_n x^{2n}}{n(2n)!} + \dots$   $0 < |x| < \pi$
- 22.49.**  $\ln |\cos x| = -\frac{x^2}{2} - \frac{x^4}{12} - \frac{x^6}{45} - \frac{17x^8}{2520} - \dots - \frac{2^{2n-1}(2^{2n}-1)B_n x^{2n}}{n(2n)!} + \dots$   $|x| < \frac{\pi}{2}$
- 22.50.**  $\ln |\tan x| = \ln |x| + \frac{x^2}{3} + \frac{7x^4}{90} + \frac{62x^6}{2835} + \dots + \frac{2^{2n}(2^{2n-1}-1)B_n x^{2n}}{n(2n)!} + \dots$   $0 < |x| < \frac{\pi}{2}$
- 22.51.**  $\frac{\ln(1+x)}{1+x} = x - (1 + \frac{1}{2})x^2 + (1 + \frac{1}{2} + \frac{1}{3})x^3 - \dots$   $|x| < 1$

### Reversion of Power Series

---

Suppose

$$\text{22.52. } y = C_1 x + C_2 x^2 + C_3 x^3 + C_4 x^4 + C_5 x^5 + C_6 x^6 + \dots$$

then

$$\text{22.53. } x = C_1 y + C_2 y^2 + C_3 y^3 + C_4 y^4 + C_5 y^5 + C_6 y^6 + \dots$$

where

$$\text{22.54. } c_1 C_1 = 1$$

$$\text{22.55. } c_1^3 C_2 = -c_2$$

$$\text{22.56. } c_1^5 C_3 = 2c_2^2 - c_1 c_3$$

$$\text{22.57. } c_1^7 C_4 = 5c_1 c_2 c_3 - 5c_2^3 - c_1^2 c_4$$

$$\text{22.58. } c_1^9 C_5 = 6c_1^2 c_2 c_4 + 3c_1^2 c_3^2 - c_1^3 c_5 + 14c_2^4 - 21c_1 c_2^2 c_3$$

$$\text{22.59. } c_1^{11} C_6 = 7c_1^3 c_2 c_5 + 84c_1 c_2^3 c_3 + 7c_1^3 c_3 c_4 - 28c_1^2 c_2 c_3^2 - c_1^4 c_6 - 28c_1^2 c_2^2 c_4 - 42c_2^5$$

### Taylor Series for Functions of Two Variables

---

$$\begin{aligned} \text{22.60. } f(x, y) &= f(a, b) + (x-a)f_x(a, b) + (y-b)f_y(a, b) \\ &\quad + \frac{1}{2!}\{(x-a)^2 f_{xx}(a, b) + 2(x-a)(y-b)f_{xy}(a, b) + (y-b)^2 f_{yy}(a, b)\} + \dots \end{aligned}$$

where  $f_x(a, b), f_y(a, b), \dots$  denote partial derivatives with respect to  $x, y, \dots$  evaluated at  $x=a, y=b$ .

# 23 BERNOULLI and EULER NUMBERS

## Definition of Bernoulli Numbers

The *Bernoulli numbers*  $B_1, B_2, B_3, \dots$  are defined by the series

$$23.1. \quad \frac{x}{e^x - 1} = 1 - \frac{x}{2} + \frac{B_1 x^2}{2!} - \frac{B_2 x^4}{4!} + \frac{B_3 x^6}{6!} - \dots \quad |x| < 2\pi$$

$$23.2. \quad 1 - \frac{x}{2} \cot \frac{x}{2} = \frac{B_1 x^2}{2!} + \frac{B_3 x^4}{4!} + \frac{B_5 x^6}{6!} + \dots \quad |x| < \pi$$

## Definition of Euler Numbers

The *Euler numbers*  $E_1, E_2, E_3, \dots$  are defined by the series

$$23.3. \quad \operatorname{sech} x = 1 - \frac{E_1 x^2}{2!} + \frac{E_2 x^4}{4!} - \frac{E_3 x^6}{6!} + \dots \quad |x| < \frac{\pi}{2}$$

$$23.4. \quad \sec x = 1 + \frac{E_1 x^2}{2!} + \frac{E_2 x^4}{4!} - \frac{E_3 x^6}{6!} + \dots \quad |x| < \frac{\pi}{2}$$

## Table of First Few Bernoulli and Euler Numbers

Bernoulli Numbers	Euler Numbers
$B_1 = 1/6$	$E_1 = 1$
$B_2 = 1/30$	$E_2 = 5$
$B_3 = 1/42$	$E_3 = 61$
$B_4 = 1/30$	$E_4 = 1385$
$B_5 = 5/66$	$E_5 = 50,521$
$B_6 = 691/2730$	$E_6 = 2,702,765$
$B_7 = 7/6$	$E_7 = 199,360,981$
$B_8 = 3617/510$	$E_8 = 19,391,512,145$
$B_9 = 43,867/798$	$E_9 = 2,404,879,675,441$
$B_{10} = 174,611/330$	$E_{10} = 370,371,188,237,525$
$B_{11} = 854,513/138$	$E_{11} = 69,348,874,393,137,901$
$B_{12} = 236,364,091/2730$	$E_{12} = 15,514,534,163,557,086,905$

---

**Relationships of Bernoulli and Euler Numbers**


---

$$23.5. \quad \binom{2n+1}{2} 2^2 B_1 - \binom{2n+1}{4} 2^4 B_2 + \binom{2n+1}{6} 2^6 B_3 - \cdots (-1)^{n-1} (2n+1) 2^{2n} B_n = 2n$$

$$23.6. \quad E_n = \binom{2n}{2} E_{n-1} - \binom{2n}{4} E_{n-2} + \binom{2n}{6} E_{n-3} - \cdots (-1)^n$$

$$23.7. \quad B_n = \frac{2n}{2^{2n}(2^{2n}-1)} \left\{ \binom{2n-1}{1} E_{n-1} - \binom{2n-1}{3} E_{n-2} + \binom{2n-1}{5} E_{n-3} - \cdots (-1)^{n-1} \right\}$$

---

**Series Involving Bernoulli and Euler Numbers**


---

$$23.8. \quad B_n = \frac{(2n)!}{2^{2n-1} \pi^{2n}} \left\{ 1 + \frac{1}{2^{2n}} + \frac{1}{3^{2n}} + \cdots \right\}$$

$$23.9. \quad B_n = \frac{2(2n)!}{(2^{2n}-1)\pi^{2n}} \left\{ 1 + \frac{1}{3^{2n}} + \frac{1}{5^{2n}} + \cdots \right\}$$

$$23.10. \quad B_n = \frac{2(2n)!}{(2^{2n-1}-1)\pi^{2n}} \left\{ 1 - \frac{1}{2^{2n}} + \frac{1}{3^{2n}} - \cdots \right\}$$

$$23.11. \quad E_n = \frac{2^{2n+2}(2n)!}{\pi^{2n+1}} \left\{ 1 - \frac{1}{3^{2n+1}} + \frac{1}{5^{2n+1}} - \cdots \right\}$$

---

**Asymptotic Formula for Bernoulli Numbers**


---

$$23.12. \quad B_n \sim 4n^{2n} (\pi e)^{-2n} \sqrt{\pi n}$$

# 24 FOURIER SERIES

## Definition of a Fourier Series

The Fourier series corresponding to a function  $f(x)$  defined in the interval  $c \leq x \leq c + 2L$  where  $c$  and  $L > 0$  are constants, is defined as

$$24.1. \quad \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right)$$

where

$$24.2. \quad \begin{cases} a_n = \frac{1}{L} \int_c^{c+2L} f(x) \cos \frac{n\pi x}{L} dx \\ b_n = \frac{1}{L} \int_c^{c+2L} f(x) \sin \frac{n\pi x}{L} dx \end{cases}$$

If  $f(x)$  and  $f'(x)$  are piecewise continuous and  $f(x)$  is defined by periodic extension of period  $2L$ , i.e.,  $f(x + 2L) = f(x)$ , then the series converges to  $f(x)$  if  $x$  is a point of continuity and to  $\frac{1}{2}\{f(x+0) + f(x-0)\}$  if  $x$  is a point of discontinuity.

## Complex Form of Fourier Series

Assuming that the series 24.1 converges to  $f(x)$ , we have

$$24.3. \quad f(x) = \sum_{n=-\infty}^{\infty} c_n e^{inx/L}$$

where

$$24.4. \quad c_n = \frac{1}{2L} \int_c^{c+2L} f(x) e^{-inx/L} dx = \begin{cases} \frac{1}{2}(a_n - ib_n) & n > 0 \\ \frac{1}{2}(a_{-n} + ib_{-n}) & n < 0 \\ \frac{1}{2}a_0 & n = 0 \end{cases}$$

## Parseval's Identity

$$24.5. \quad \frac{1}{L} \int_c^{c+2L} \{f(x)\}^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2)$$

## Generalized Parseval Identity

$$24.6. \quad \frac{1}{L} \int_c^{c+2L} f(x)g(x) dx = \frac{a_0 c_0}{2} + \sum_{n=1}^{\infty} (a_n c_n + b_n d_n)$$

where  $a_n, b_n$  and  $c_n, d_n$  are the Fourier coefficients corresponding to  $f(x)$  and  $g(x)$ , respectively.

## Special Fourier Series and Their Graphs

24.7.  $f(x) = \begin{cases} 1 & 0 < x < \pi \\ -1 & -\pi < x < 0 \end{cases}$

$$\frac{4}{\pi} \left( \frac{\sin x}{1} + \frac{\sin 3x}{3} + \frac{\sin 5x}{5} + \dots \right)$$




Fig. 24-1

24.8.  $f(x) = |x| = \begin{cases} x & 0 < x < \pi \\ -x & -\pi < x < 0 \end{cases}$

$$\frac{\pi}{2} - \frac{4}{\pi} \left( \frac{\cos x}{1^2} + \frac{\cos 3x}{3^2} + \frac{\cos 5x}{5^2} + \dots \right)$$




Fig. 24-2

24.9.  $f(x) = x, \quad -\pi < x < \pi$

$$2 \left( \frac{\sin x}{1} - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \dots \right)$$




Fig. 24-3

24.10.  $f(x) = x, \quad 0 < x < 2\pi$

$$\pi - 2 \left( \frac{\sin x}{1} + \frac{\sin 2x}{2} + \frac{\sin 3x}{3} + \dots \right)$$




Fig. 24-4

24.11.  $f(x) = |\sin x|, \quad -\pi < x < \pi$

$$\frac{2}{\pi} - \frac{4}{\pi} \left( \frac{\cos 2x}{1 \cdot 3} + \frac{\cos 4x}{3 \cdot 5} + \frac{\cos 6x}{5 \cdot 7} + \dots \right)$$




Fig. 24-5

**24.12.**  $f(x) = \begin{cases} \sin x & 0 < x < \pi \\ 0 & \pi < x < 2\pi \end{cases}$

$$\frac{1}{\pi} + \frac{1}{2}\sin x - \frac{2}{\pi} \left( \frac{\cos 2x}{1 \cdot 3} + \frac{\cos 4x}{3 \cdot 5} + \frac{\cos 6x}{5 \cdot 7} + \dots \right)$$




Fig. 24-6

**24.13.**  $f(x) = \begin{cases} \cos x & 0 < x < \pi \\ -\cos x & -\pi < x < 0 \end{cases}$

$$\frac{8}{\pi} \left( \frac{\sin 2x}{1 \cdot 3} + \frac{2\sin 4x}{3 \cdot 5} + \frac{3\sin 6x}{5 \cdot 7} + \dots \right)$$




Fig. 24-7

**24.14.**  $f(x) = x^2, \quad -\pi < x < \pi$

$$\frac{\pi^2}{3} - 4 \left( \frac{\cos x}{1^2} - \frac{\cos 2x}{2^2} + \frac{\cos 3x}{3^2} - \dots \right)$$




Fig. 24-8

**24.15.**  $f(x) = x(\pi - x), \quad 0 < x < \pi$

$$\frac{\pi^2}{6} - \left( \frac{\cos 2x}{1^2} + \frac{\cos 4x}{2^2} + \frac{\cos 6x}{3^2} + \dots \right)$$




Fig. 24-9

**24.16.**  $f(x) = x(\pi - x)(\pi + x), \quad -\pi < x < \pi$

$$12 \left( \frac{\sin x}{1^3} - \frac{\sin 2x}{2^3} + \frac{\sin 3x}{3^3} - \dots \right)$$




Fig. 24-10

**24.17.**  $f(x) = \begin{cases} 0 & 0 < x < \pi - \alpha \\ 1 & \pi - \alpha < x < \pi + \alpha \\ 0 & \pi + \alpha < x < 2\pi \end{cases}$

$$\frac{\alpha}{\pi} - \frac{2}{\pi} \left( \frac{\sin \alpha \cos x}{1} - \frac{\sin 2\alpha \cos 2x}{2} + \frac{\sin 3\alpha \cos 3x}{3} - \dots \right)$$




Fig. 24-11

**24.18.**  $f(x) = \begin{cases} x(\pi - x) & 0 < x < \pi \\ -x(\pi - x) & -\pi < x < 0 \end{cases}$

$$\frac{8}{\pi} \left( \frac{\sin x}{1^3} + \frac{\sin 3x}{3^3} + \frac{\sin 5x}{5^3} + \dots \right)$$




Fig. 24-12

## Miscellaneous Fourier Series

**24.19.**  $f(x) = \sin \mu x, \quad -\pi < x < \pi, \quad \mu \neq \text{integer}$

$$\frac{2 \sin \mu \pi}{\pi} \left( \frac{\sin x}{1^2 - \mu^2} - \frac{2 \sin 2x}{2^2 - \mu^2} + \frac{3 \sin 3x}{3^2 - \mu^2} - \dots \right)$$

**24.20.**  $f(x) = \cos \mu x, \quad -\pi < x < \pi, \quad \mu \neq \text{integer}$

$$\frac{2\mu \sin \mu \pi}{\pi} \left( \frac{1}{2\mu^2} + \frac{\cos x}{1^2 - \mu^2} - \frac{\cos 2x}{2^2 - \mu^2} + \frac{\cos 3x}{3^2 - \mu^2} - \dots \right)$$

**24.21.**  $f(x) = \tan^{-1}[(a \sin x) / (1 - a \cos x)], \quad -\pi < x < \pi, \quad |a| < 1$

$$a \sin x + \frac{a^2}{2} \sin 2x + \frac{a^3}{3} \sin 3x + \dots$$

**24.22.**  $f(x) = \ln(1 - 2a \cos x + a^2), \quad -\pi < x < \pi, \quad |a| < 1$

$$-2 \left( a \cos x + \frac{a^2}{2} \cos 2x + \frac{a^3}{3} \cos 3x + \dots \right)$$

**24.23.**  $f(x) = \frac{1}{2} \tan^{-1}[(2a \sin x) / (1 - a^2)], \quad -\pi < x < \pi, \quad |a| < 1$

$$a \sin x + \frac{a^3}{3} \sin 3x + \frac{a^5}{5} \sin 5x + \dots$$

**24.24.**  $f(x) = \frac{1}{2} \tan^{-1}[(2a \cos x) / (1 - a^2)], \quad -\pi < x < \pi, \quad |a| < 1$

$$a \cos x - \frac{a^3}{3} \cos 3x + \frac{a^5}{5} \cos 5x - \dots$$

**24.25.**  $f(x) = e^{\mu x}, -\pi < x < \pi$

$$\frac{2\sinh \mu\pi}{\pi} \left( \frac{1}{2\mu} + \sum_{n=1}^{\infty} \frac{(-1)^n (\mu \cos nx - n \sin nx)}{\mu^2 + n^2} \right)$$

**24.26.**  $f(x) = \sinh \mu x, -\pi < x < \pi$

$$\frac{2\sinh \mu\pi}{\pi} \left( \frac{\sin x}{1^2 + \mu^2} - \frac{2\sin 2x}{2^2 + \mu^2} + \frac{3\sin 3x}{3^2 + \mu^2} - \dots \right)$$

**24.27.**  $f(x) = \cosh \mu x, -\pi < x < \pi$

$$\frac{2\mu \sinh \mu\pi}{\pi} \left( \frac{1}{2\mu^2} - \frac{\cos x}{1^2 + \mu^2} + \frac{\cos 2x}{2^2 + \mu^2} - \frac{\cos 3x}{3^2 + \mu^2} + \dots \right)$$

**24.28.**  $f(x) = \ln |\sin \frac{1}{2}x|, 0 < x < \pi$

$$-\left( \ln 2 + \frac{\cos x}{1} + \frac{\cos 2x}{2} + \frac{\cos 3x}{3} + \dots \right)$$

**24.29.**  $f(x) = \ln |\cos \frac{1}{2}x|, -\pi < x < \pi$

$$-\left( \ln 2 - \frac{\cos x}{1} + \frac{\cos 2x}{2} - \frac{\cos 3x}{3} + \dots \right)$$

**24.30.**  $f(x) = \frac{1}{6}\pi^2 - \frac{1}{2}\pi x + \frac{1}{4}x^2, 0 \leq x \leq 2\pi$

$$\frac{\cos x}{1^2} + \frac{\cos 2x}{2^2} + \frac{\cos 3x}{3^2} + \dots$$

**24.31.**  $f(x) = \frac{1}{12}x(x - \pi)(x - 2\pi), 0 \leq x \leq 2\pi$

$$\frac{\sin x}{1^3} + \frac{\sin 2x}{2^3} + \frac{\sin 3x}{3^3} + \dots$$

**24.32.**  $f(x) = \frac{1}{90}\pi^4 - \frac{1}{12}\pi^2 x^2 + \frac{1}{12}\pi x^3 - \frac{1}{48}x^4, 0 \leq x \leq 2\pi$

$$\frac{\cos x}{1^4} + \frac{\cos 2x}{2^4} + \frac{\cos 3x}{3^4} + \dots$$

## Section VII: Special Functions and Polynomials

# 25 THE GAMMA FUNCTION

### Definition of the Gamma Function $\Gamma(n)$ for $n > 0$

$$25.1. \quad \Gamma(n) = \int_0^{\infty} t^{n-1} e^{-t} dt \quad n > 0$$

### Recursion Formula

$$25.2. \quad \Gamma(n+1) = n\Gamma(n)$$

If  $n = 0, 1, 2, \dots$ , a nonnegative integer, we have the following (where  $0! = 1$ ):

$$25.3. \quad \Gamma(n+1) = n!$$

### The Gamma Function for $n < 0$

For  $n < 0$  the gamma function can be defined by using 25.2, that is,

$$25.4. \quad \Gamma(n) = \frac{\Gamma(n+1)}{n}$$

### Graph of the Gamma Function




Fig. 25-1

### Special Values for the Gamma Function

---

$$25.5. \quad \Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

$$25.6. \quad \Gamma\left(m + \frac{1}{2}\right) = \frac{1 \cdot 3 \cdot 5 \cdots (2m-1)}{2^m} \sqrt{\pi} \quad m = 1, 2, 3, \dots$$

$$25.7. \quad \Gamma\left(-m + \frac{1}{2}\right) = \frac{(-1)^m 2^m \sqrt{\pi}}{1 \cdot 3 \cdot 5 \cdots (2m-1)} \quad m = 1, 2, 3, \dots$$

### Relationships Among Gamma Functions

---

$$25.8. \quad \Gamma(p)\Gamma(1-p) = \frac{\pi}{\sin p\pi}$$

$$25.9. \quad 2^{2x-1} \Gamma(x)\Gamma(x + \frac{1}{2}) = \sqrt{\pi} \Gamma(2x)$$

This is called the *duplication formula*.

$$25.10. \quad \Gamma(x)\Gamma\left(x + \frac{1}{m}\right)\Gamma\left(x + \frac{2}{m}\right)\cdots\Gamma\left(x + \frac{m-1}{m}\right) = m^{1/2-mx} (2\pi)^{(m-1)/2} \Gamma(mx)$$

For  $m = 2$  this reduces to 25.9.

### Other Definitions of the Gamma Function

---

$$25.11. \quad \Gamma(x+1) = \lim_{k \rightarrow \infty} \frac{1 \cdot 2 \cdot 3 \cdots k}{(x+1)(x+2)\cdots(x+k)} k^x$$

$$25.12. \quad \frac{1}{\Gamma(x)} = xe^{\gamma x} \prod_{m=1}^{\infty} \left\{ \left(1 + \frac{x}{m}\right) e^{-x/m} \right\}$$

This is an infinite product representation for the gamma function where  $\gamma$  is Euler's constant defined in 1.3, page 3.

### Derivatives of the Gamma Function

---

$$25.13. \quad \Gamma'(1) = \int_0^\infty e^{-x} \ln x dx = -\gamma$$

$$25.14. \quad \frac{\Gamma'(x)}{\Gamma(x)} = -\gamma + \left(\frac{1}{1} - \frac{1}{x}\right) + \left(\frac{1}{2} - \frac{1}{x+1}\right) + \cdots + \left(\frac{1}{n} - \frac{1}{x+n-1}\right) + \cdots$$

Here again is Euler's constant  $\gamma$ .

### Asymptotic Expansions for the Gamma Function

---

$$25.15. \quad \Gamma(x+1) = \sqrt{2\pi}xx^x e^{-x} \left\{ 1 + \frac{1}{12x} + \frac{1}{288x^2} - \frac{139}{51,840x^3} + \dots \right\}$$

This is called *Stirling's asymptotic series*.

If we let  $x = n$  a positive integer in 25.15, then a useful approximation for  $n!$  where  $n$  is large (e.g.,  $n > 10$ ) is given by *Stirling's formula*

$$25.16. \quad n! \sim \sqrt{2\pi n} n^n e^{-n}$$

where  $\sim$  is used to indicate that the ratio of the terms on each side approaches 1 as  $n \rightarrow \infty$ .

### Miscellaneous Results

---

$$25.17. \quad |\Gamma(ix)|^2 = \frac{\pi}{x \sinh \pi x}$$

# 26 THE BETA FUNCTION

## Definition of the Beta Function $B(m, n)$

---

$$26.1. \quad B(m, n) = \int_0^1 t^{m-1} (1-t)^{n-1} dt \quad m > 0, n > 0$$

## Relationship of Beta Function to Gamma Function

---

$$26.2. \quad B(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}$$

Extensions of  $B(m, n)$  to  $m < 0, n < 0$  are provided by using 25.4.

## Some Important Results

---

$$26.3. \quad B(m, n) = B(n, m)$$

$$26.4. \quad B(m, n) = 2 \int_0^{\pi/2} \sin^{2m-1} \theta \cos^{2n-1} \theta d\theta$$

$$26.5. \quad B(m, n) = \int_0^\infty \frac{t^{m-1}}{(1+t)^{m+n}} dt$$

$$26.6. \quad B(m, n) = r^n (r+1)^m \int_0^1 \frac{t^{m-1} (1-t)^{n-1}}{(r+t)^{m+n}} dt$$

# 27 BESSEL FUNCTIONS

## Bessel's Differential Equation

---

$$27.1. \quad x^2 y'' + xy' + (x^2 - n^2)y = 0 \quad n \geq 0$$

Solutions of this equation are called *Bessel functions of order n*.

### Bessel Functions of the First Kind of Order n

---

$$27.2. \quad J_n(x) = \frac{x^n}{2^n \Gamma(n+1)} \left\{ 1 - \frac{x^2}{2(2n+2)} + \frac{x^4}{2 \cdot 4(2n+2)(2n+4)} - \dots \right\}$$

$$= \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{n+2k}}{k! \Gamma(n+k+1)}$$

$$27.3. \quad J_{-n}(x) = \frac{x^{-n}}{2^{-n} \Gamma(1-n)} \left\{ 1 - \frac{x^2}{2(2-2n)} + \frac{x^4}{2 \cdot 4(2-2n)(4-2n)} - \dots \right\}$$

$$= \sum_{k=0}^{\infty} \frac{(-1)^k (x/2)^{2k-n}}{k! \Gamma(k+1-n)}$$

$$27.4. \quad J_{-n}(x) = (-1)^n J_n(x) \quad n = 0, 1, 2, \dots$$

If  $n \neq 0, 1, 2, \dots$ ,  $J_n(x)$  and  $J_{-n}(x)$  are linearly independent.

If  $n \neq 0, 1, 2, \dots$ ,  $J_n(x)$  is bounded at  $x = 0$  while  $J_{-n}(x)$  is unbounded.

For  $n = 0, 1$  we have

$$27.5. \quad J_0(x) = 1 - \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} - \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$

$$27.6. \quad J_1(x) = \frac{x}{2} - \frac{x^3}{2^2 \cdot 4} + \frac{x^5}{2^2 \cdot 4^2 \cdot 6^2} - \frac{x^7}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8} + \dots$$

$$27.7. \quad J'_0(x) = -J_1(x)$$

### Bessel Functions of the Second Kind of Order n

---

$$27.8. \quad Y_n(x) = \begin{cases} \frac{J_n(x) \cos n\pi - J_{-n}(x)}{\sin n\pi} & n \neq 0, 1, 2, \dots \\ \lim_{p \rightarrow n} \frac{J_p(x) \cos p\pi - J_{-p}(x)}{\sin p\pi} & n = 0, 1, 2, \dots \end{cases}$$

This is also called *Weber's function* or *Neumann's function* [also denoted by  $N_n(x)$ ].

For  $n = 0, 1, 2, \dots$ , L' Hospital's rule yields

$$27.9. \quad Y_n(x) = \frac{2}{\pi} \{ \ln(x/2) + \gamma \} J_n(x) - \frac{1}{\pi} \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} (x/2)^{2k-n} \\ - \frac{1}{\pi} \sum_{k=0}^{\infty} (-1)^k \{ \Phi(k) + \Phi(n+k) \} \frac{(x/2)^{2k+n}}{k!(n+k)!}$$

where  $\gamma = .5772156 \dots$  is Euler's constant (see 1.20) and

$$27.10. \quad \Phi(p) = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{P}, \quad \Phi(0) = 0$$

For  $n = 0$ ,

$$27.11. \quad Y_0(x) = \frac{2}{\pi} \{ \ln(x/2) + \gamma \} J_0(x) + \frac{2}{\pi} \left\{ \frac{x^2}{2^2} - \frac{x^4}{2^2 4^2} \left( 1 + \frac{1}{2} \right) + \frac{x^6}{2^2 4^2 6^2} \left( 1 + \frac{1}{2} + \frac{1}{3} \right) - \dots \right\}$$

$$27.12. \quad Y_{-n}(x) = (-1)^n Y_n(x) \quad n = 0, 1, 2, \dots$$

For any value  $n \geq 0$ ,  $J_n(x)$  is bounded at  $x = 0$  while  $Y_n(x)$  is unbounded.

### General Solution of Bessel's Differential Equation

---

$$27.13. \quad y = AJ_n(x) + BJ_{-n}(x) \quad n \neq 0, 1, 2, \dots$$

$$27.14. \quad y = AJ_n(x) + BY_n(x) \quad \text{all } n$$

$$27.15. \quad y = AJ_n(x) + BJ_n(x) \int \frac{dx}{xJ_n^2(x)} \quad \text{all } n$$

where  $A$  and  $B$  are arbitrary constants.

### Generating Function for $J_n(x)$

---

$$27.16. \quad e^{x(t-1/t)/2} = \sum_{n=-\infty}^{\infty} J_n(x)t^n$$

### Recurrence Formulas for Bessel Functions

---

$$27.17. \quad J_{n+1}(x) = \frac{2n}{x} J_n(x) - J_{n-1}(x)$$

$$27.18. \quad J'_n(x) = \frac{1}{2} \{ J_{n-1}(x) - J_{n+1}(x) \}$$

$$27.19. \quad xJ'_n(x) = xJ_{n-1}(x) - nJ_n(x)$$

$$27.20. \quad xJ'_n(x) = nJ_n(x) - xJ_{n+1}(x)$$

$$27.21. \quad \frac{d}{dx} \{x^n J_n(x)\} = x^n J_{n-1}(x)$$

$$27.22. \quad \frac{d}{dx} \{x^{-n} J_n(x)\} = -x^{-n} J_{n+1}(x)$$

The functions  $Y_n(x)$  satisfy identical relations.

### Bessel Functions of Order Equal to Half an Odd Integer

---

In this case the functions are expressible in terms of sines and cosines.

$$27.23. \quad J_{1/2}(x) = \sqrt{\frac{2}{\pi x}} \sin x$$

$$27.26. \quad J_{-3/2}(x) = \sqrt{\frac{2}{\pi x}} \left( \frac{\cos x}{x} + \sin x \right)$$

$$27.24. \quad J_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cos x$$

$$27.27. \quad J_{5/2}(x) = \sqrt{\frac{2}{\pi x}} \left\{ \left( \frac{3}{x^2} - 1 \right) \sin x - \frac{3}{x} \cos x \right\}$$

$$27.25. \quad J_{3/2}(x) = \sqrt{\frac{2}{\pi x}} \left( \frac{\sin x}{x} - \cos x \right)$$

$$27.28. \quad J_{-5/2}(x) = \sqrt{\frac{2}{\pi x}} \left\{ \frac{3}{x} \sin x + \left( \frac{3}{x} - 1 \right) \cos x \right\}$$

For further results use the recurrence formula. Results for  $Y_{1/2}(x), Y_{3/2}(x), \dots$  are obtained from 27.8.

### Hankel Functions of First and Second Kinds of Order $n$

---

$$27.29. \quad H_n^{(1)}(x) = J_n(x) + iY_n(x)$$

$$27.30. \quad H_n^{(2)}(x) = J_n(x) - iY_n(x)$$

### Bessel's Modified Differential Equation

---

$$27.31. \quad x^2 y'' + xy' - (x^2 + n^2)y = 0 \quad n \geq 0$$

Solutions of this equation are called *modified Bessel functions of order  $n$* .

### Modified Bessel Functions of the First Kind of Order $n$

---

$$27.32. \quad I_n(x) = i^{-n} J_n(ix) = e^{-n\pi i/2} J_n(ix) \\ = \frac{x^n}{2^n \Gamma(n+1)} \left\{ 1 + \frac{x^2}{2(2n+2)} + \frac{x^4}{2 \cdot 4(2n+2)(2n+4)} + \dots \right\} = \sum_{k=0}^{\infty} \frac{(x/2)^{n+2k}}{k! \Gamma(n+k+1)}$$

$$27.33. \quad I_{-n}(x) = i^n J_{-n}(ix) = e^{n\pi i/2} J_{-n}(ix) \\ = \frac{x^{-n}}{2^{-n} \Gamma(1-n)} \left\{ 1 + \frac{x^2}{2(2-2n)} + \frac{x^4}{2 \cdot 4(2-2n)(4-2n)} + \dots \right\} = \sum_{k=0}^{\infty} \frac{(x/2)^{2k-n}}{k! \Gamma(k+1-n)}$$

$$27.34. \quad I_{-n}(x) = I_n(x) \quad n = 0, 1, 2, \dots$$

If  $n \neq 0, 1, 2, \dots$ , then  $I_n(x)$  and  $I_{-n}(x)$  are linearly independent.

For  $n = 0, 1$ , we have

$$27.35. \quad I_0(x) = 1 + \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} + \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} + \dots$$

$$27.36. \quad I_1(x) = \frac{x}{2} + \frac{x^3}{2^2 \cdot 4} + \frac{x^5}{2^2 \cdot 4^2 \cdot 6} + \frac{x^7}{2^2 \cdot 4^2 \cdot 6^2 \cdot 8} + \dots$$

$$27.37. \quad I'_0(x) = I_1(x)$$

### Modified Bessel Functions of the Second Kind of Order $n$

---

$$27.38. \quad K_n(x) = \begin{cases} \frac{\pi}{2 \sin n\pi} \{I_{-n}(x) - I_n(x)\} & n \neq 0, 1, 2, \dots \\ \lim_{p \rightarrow n} \frac{\pi}{2 \sin p\pi} \{I_{-p}(x) - I_p(x)\} & n = 0, 1, 2, \dots \end{cases}$$

For  $n = 0, 1, 2, \dots$ , L' Hospital's rule yields

$$27.39. \quad K_n(x) = (-1)^{n+1} \{ \ln(x/2) + \gamma \} I_n(x) + \frac{1}{2} \sum_{k=0}^{n-1} (-1)^k (n-k-1)! (x/2)^{2k-n} + \frac{(-1)^n}{2} \sum_{k=0}^{\infty} \frac{(x/2)^{n+2k}}{k!(n+k)!} \{ \Phi(k) + \Phi(n+k) \}$$

where  $\Phi(p)$  is given by 27.10.

For  $n = 0$ ,

$$27.40. \quad K_0(x) = -\{ \ln(x/2) + \gamma \} I_0(x) + \frac{x^2}{2^2} + \frac{x^4}{2^2 \cdot 4^2} \left( 1 + \frac{1}{2} \right) + \frac{x^6}{2^2 \cdot 4^2 \cdot 6^2} \left( 1 + \frac{1}{2} + \frac{1}{3} \right) + \dots$$

$$27.41. \quad K_{-n}(x) = K_n(x) \quad n = 0, 1, 2, \dots$$

### General Solution of Bessel's Modified Equation

---

$$27.42. \quad y = AI_n(x) + BI_{-n}(x) \quad n \neq 0, 1, 2, \dots$$

$$27.43. \quad y = AI_n(x) + BK_n(x) \quad \text{all } n$$

$$27.44. \quad y = AI_n(x) + BI_n(x) \int \frac{dx}{x I_n^2(x)} \quad \text{all } n$$

where  $A$  and  $B$  are arbitrary constants.

### Generating Function for $I_n(x)$

---

$$27.45. \quad e^{x(t+1/t)/2} = \sum_{n=-\infty}^{\infty} I_n(x) t^n$$

### Recurrence Formulas for Modified Bessel Functions

---

$$27.46. \quad I_{n+1}(x) = I_{n-1}(x) - \frac{2n}{x} I_n(x)$$

$$27.52. \quad K_{n+1}(x) = K_{n-1}(x) + \frac{2n}{x} K_n(x)$$

$$27.47. \quad I'_n(x) = \frac{1}{2} \{I_{n-1}(x) + I_{n+1}(x)\}$$

$$27.53. \quad K'_n(x) = -\frac{1}{2} \{K_{n-1}(x) + K_{n+1}(x)\}$$

$$27.48. \quad xI'_n(x) = xI_{n-1}(x) - nI_n(x)$$

$$27.54. \quad xK'_n(x) = -xK_{n-1}(x) - nK_n(x)$$

$$27.49. \quad xI'_n(x) = xI_{n+1}(x) + nI_n(x)$$

$$27.55. \quad xK'_n(x) = nK_n(x) - xK_{n+1}(x)$$

$$27.50. \quad \frac{d}{dx} \{x^n I_n(x)\} = x^n I_{n-1}(x)$$

$$27.56. \quad \frac{d}{dx} \{x^n K_n(x)\} = -x^n K_{n-1}(x)$$

$$27.51. \quad \frac{d}{dx} \{x^{-n} I_n(x)\} = x^{-n} I_{n+1}(x)$$

$$27.57. \quad \frac{d}{dx} \{x^{-n} K_n(x)\} = -x^{-n} K_{n+1}(x)$$

### Modified Bessel Functions of Order Equal to Half an Odd Integer

---

In this case the functions are expressible in terms of hyperbolic sines and cosines.

$$27.58. \quad I_{1/2}(x) = \sqrt{\frac{2}{\pi x}} \sinh x$$

$$27.61. \quad I_{-3/2}(x) = \sqrt{\frac{2}{\pi x}} \left( \sinh x - \frac{\cosh x}{x} \right)$$

$$27.59. \quad I_{-1/2}(x) = \sqrt{\frac{2}{\pi x}} \cosh x$$

$$27.62. \quad I_{5/2}(x) = \sqrt{\frac{2}{\pi x}} \left\{ \left( \frac{3}{x^2} + 1 \right) \sinh x - \frac{3}{x} \cosh x \right\}$$

$$27.60. \quad I_{3/2}(x) = \sqrt{\frac{2}{\pi x}} \left( \cosh x - \frac{\sinh x}{x} \right)$$

$$27.63. \quad I_{-5/2}(x) = \sqrt{\frac{2}{\pi x}} \left\{ \left( \frac{3}{x^2} + 1 \right) \cosh x - \frac{3}{x} \sinh x \right\}$$

For further results use the recurrence formula 27.46. Results for  $K_{1/2}(x)$ ,  $K_{3/2}(x)$ , ... are obtained from 27.38.

### Ber and Bei Functions

---

The real and imaginary parts of  $J_n(xe^{3\pi i/4})$  are denoted by  $\text{Ber}_n(x)$  and  $\text{Bei}_n(x)$  where

$$27.64. \quad \text{Ber}_n(x) = \sum_{k=0}^{\infty} \frac{(x/2)^{2k+n}}{k! \Gamma(n+k+1)} \cos \frac{(3n+2k)\pi}{4}$$

$$27.65. \quad \text{Bei}_n(x) = \sum_{k=0}^{\infty} \frac{(x/2)^{2k+n}}{k! \Gamma(n+k+1)} \sin \frac{(3n+2k)\pi}{4}$$

If  $n = 0$ .

$$27.66. \quad \text{Ber}(x) = 1 - \frac{(x/2)^4}{2!^2} + \frac{(x/2)^8}{4!^2} - \dots$$

$$27.67. \quad \text{Bei}(x) = (x/2)^2 - \frac{(x/2)^6}{3!^2} + \frac{(x/2)^{10}}{5!^2} - \dots$$

### Ker and Kei Functions

The real and imaginary parts of  $e^{-n\pi i/2} K_n(xe^{\pi i/4})$  are denoted by  $\text{Ker}_n(x)$  and  $\text{Kei}_n(x)$  where

$$27.68. \quad \text{Ker}_n(x) = -\{\ln(x/2) + \gamma\} \text{Ber}_n(x) + \frac{1}{4}\pi \text{Bei}_n(x)$$

$$\begin{aligned} &+ \frac{1}{2} \sum_{k=0}^{n-1} \frac{(n-k-1)!(x/2)^{2k-n}}{k!} \cos \frac{(3n+2k)\pi}{4} \\ &+ \frac{1}{2} \sum_{k=0}^{\infty} \frac{(x/2)^{n+2k}}{k!(n+k)!} \{\Phi(k) + \Phi(n+k)\} \cos \frac{(3n+2k)\pi}{4} \end{aligned}$$

$$27.69. \quad \text{Kei}_n(x) = -\{\ln(x/2) + \gamma\} \text{Bei}_n(x) - \frac{1}{4}\pi \text{Ber}_n(x)$$

$$\begin{aligned} &- \frac{1}{2} \sum_{k=0}^{n-1} \frac{(n-k-1)!(x/2)^{2k-n}}{k!} \sin \frac{(3n+2k)\pi}{4} \\ &+ \frac{1}{2} \sum_{k=0}^{\infty} \frac{(x/2)^{n+2k}}{k!(n+k)!} \{\Phi(k) + \Phi(n+k)\} \sin \frac{(3n+2k)\pi}{4} \end{aligned}$$

and  $\Phi$  is given by 27.10.

If  $n = 0$ ,

$$27.70. \quad \text{Ker}(x) = -\{\ln(x/2) + \gamma\} \text{Ber}(x) + \frac{\pi}{4} \text{Bei}(x) + 1 - \frac{(x/2)^4}{2!^2} (1 + \frac{1}{2}) + \frac{(x/2)^8}{4!^2} (1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}) - \dots$$

$$27.71. \quad \text{Kei}(x) = -\{\ln(x/2) + \gamma\} \text{Bei}(x) - \frac{\pi}{4} \text{Ber}(x) + (x/2)^2 - \frac{(x/2)^6}{3!^2} (1 + \frac{1}{2} + \frac{1}{3}) + \dots$$

### Differential Equation For Ber, Bei, Ker, Kei Functions

$$27.72. \quad x^2 y'' + xy' - (ix^2 + n^2)y = 0$$

The general solution of this equation is

$$27.73. \quad y = A\{\text{Ber}_n(x) + i\text{Bei}_n(x)\} + B\{\text{Ker}_n(x) + i\text{Kei}_n(x)\}$$

### Graphs of Bessel Functions




Fig. 27-1




Fig. 27-2




Fig. 27-3




Fig. 27-4




Fig. 27-5




Fig. 27-6

### Indefinite Integrals Involving Bessel Functions

$$27.74. \quad \int x J_0(x) dx = x J_1(x)$$

$$27.75. \quad \int x^2 J_0(x) dx = x^2 J_1(x) + x J_0(x) - \int J_0(x) dx$$

$$27.76. \quad \int x^m J_0(x) dx = x^m J_1(x) + (m-1)x^{m-1} J_0(x) - (m-1)^2 \int x^{m-2} J_0(x) dx$$

$$27.77. \quad \int \frac{J_0(x)}{x^2} dx = J_1(x) - \frac{J_0(x)}{x} - \int J_0(x) dx$$

$$27.78. \quad \int \frac{J_0(x)}{x^m} dx = \frac{J_1(x)}{(m-1)^2 x^{m-2}} - \frac{J_0(x)}{(m-1)x^{m-1}} - \frac{1}{(m-1)^2} \int \frac{J_0(x)}{x^{m-2}} dx$$

$$27.79. \quad \int J_1(x) dx = -J_0(x)$$

$$27.80. \quad \int x J_1(x) dx = -x J_0(x) + \int J_0(x) dx$$

$$27.81. \quad \int x^m J_1(x) dx = -x^m J_0(x) + m \int x^{m-1} J_0(x) dx$$

$$27.82. \int \frac{J_1(x)}{x} dx = -J_1(x) + \int J_0(x) dx$$

$$27.83. \int \frac{J_1(x)}{x^m} dx = -\frac{J_1(x)}{mx^{m-1}} + \frac{1}{m} \int \frac{J_0(x)}{x^{m-1}} dx$$

$$27.84. \int x^n J_{n-1}(x) dx = x^n J_n(x)$$

$$27.85. \int x^{-n} J_{n+1}(x) dx = -x^{-n} J_n(x)$$

$$27.86. \int x^m J_n(x) dx = -x^m J_{n-1}(x) + (m+n-1) \int x^{m-1} J_{n-1}(x) dx$$

$$27.87. \int x J_n(\alpha x) J_n(\beta x) dx = \frac{x \{ \alpha J_n(\beta x) J'_n(\alpha x) - \beta J_n(\alpha x) J'_n(\beta x) \}}{\beta^2 - \alpha^2}$$

$$27.88. \int x J_n^2(\alpha x) dx = \frac{x^2}{2} \{ J'_n(\alpha x) \}^2 + \frac{x^2}{2} \left( 1 - \frac{n^2}{\alpha^2 x^2} \right) \{ J_n(\alpha x) \}^2$$

The above results also hold if we replace  $J_n(x)$  by  $Y_n(x)$  or, more generally,  $AJ_n(x) + BY_n(x)$  where  $A$  and  $B$  are constants.

### Definite Integrals Involving Bessel Functions

---

$$27.89. \int_0^\infty e^{-ax} J_0(bx) dx = \frac{1}{\sqrt{a^2 + b^2}}$$

$$27.90. \int_0^\infty e^{-ax} J_n(bx) dx = \frac{(\sqrt{a^2 + b^2} - a)^n}{b^n \sqrt{a^2 + b^2}} \quad n > -1$$

$$27.91. \int_0^\infty \cos ax J_0(bx) dx = \begin{cases} \frac{1}{\sqrt{a^2 - b^2}} & a > b \\ 0 & a < b \end{cases}$$

$$27.92. \int_0^\infty J_n(bx) dx = \frac{1}{b}, \quad n > -1$$

$$27.93. \int_0^\infty \frac{J_n(bx)}{x} dx = \frac{1}{n}, \quad n = 1, 2, 3, \dots$$

$$27.94. \int_0^\infty e^{-ax} J_0(b\sqrt{x}) dx = \frac{e^{-b^2/4a}}{a}$$

$$27.95. \int_0^1 x J_n(\alpha x) J_n(\beta x) dx = \frac{\alpha J_n(\beta) J'_n(\alpha) - \beta J_n(\alpha) J'_n(\beta)}{\beta^2 - \alpha^2}$$

$$27.96. \int_0^1 x J_n^2(\alpha x) dx = \frac{1}{2} \{ J'_n(\alpha) \}^2 + \frac{1}{2} (1 - n^2/\alpha^2) \{ J_n(\alpha) \}^2$$

$$27.97. \int_0^1 x J_0(\alpha x) I_0(\beta x) dx = \frac{\beta J_0(\alpha) I'_0(\beta) - \alpha J'_0(\alpha) I_0(\beta)}{\alpha^2 + \beta^2}$$

### Integral Representations for Bessel Functions

---

$$27.98. \quad J_0(x) = \frac{1}{\pi} \int_0^\pi \cos(x \sin \theta) d\theta$$

$$27.99. \quad J_n(x) = \frac{1}{\pi} \int_0^\pi \cos(n\theta - x \sin \theta) d\theta \quad n = \text{integer}$$

$$27.100. \quad J_n(x) = \frac{x^n}{2^n \sqrt{\pi} \Gamma(n + \frac{1}{2})} \int_0^\pi \cos(x \sin \theta) \cos^{2n} \theta d\theta, \quad n > -\frac{1}{2}$$

$$27.101. \quad Y_0(x) = -\frac{2}{\pi} \int_0^\infty \cos(x \cosh u) du$$

$$27.102. \quad I_0(x) = \frac{1}{\pi} \int_0^\pi \cosh(x \sin \theta) d\theta = \frac{1}{2\pi} \int_0^{2\pi} e^{x \sin \theta} d\theta$$

### Asymptotic Expansions

---

$$27.103. \quad J_n(x) \sim \sqrt{\frac{2}{\pi x}} \cos\left(x - \frac{n\pi}{2} - \frac{\pi}{4}\right) \quad \text{where } x \text{ is large}$$

$$27.104. \quad Y_n(x) \sim \sqrt{\frac{2}{\pi x}} \sin\left(x - \frac{n\pi}{2} - \frac{\pi}{4}\right) \quad \text{where } x \text{ is large}$$

$$27.105. \quad J_n(x) \sim \frac{1}{\sqrt{2\pi n}} \left(\frac{ex}{2n}\right)^n \quad \text{where } n \text{ is large}$$

$$27.106. \quad Y_n(x) \sim -\sqrt{\frac{2}{\pi n}} \left(\frac{ex}{2n}\right)^{-n} \quad \text{where } n \text{ is large}$$

$$27.107. \quad I_n(x) \sim \frac{e^x}{\sqrt{2\pi x}} \quad \text{where } x \text{ is large}$$

$$27.108. \quad K_n(x) \sim \frac{e^{-x}}{\sqrt{2\pi x}} \quad \text{where } x \text{ is large}$$

### Orthogonal Series of Bessel Functions

---

Let  $\lambda_1, \lambda_2, \lambda_3, \dots$  be the positive roots of  $RJ_n(x) + SxJ'_n(x) = 0$ ,  $n > -1$ . Then the following series expansions hold under the conditions indicated.

$S = 0, R \neq 0$ , i.e., $\lambda_1, \lambda_2, \lambda_3, \dots$ are positive roots of $J_N(x) = 0$
---

$$27.109. \quad f(x) = A_1 J_n(\lambda_1 x) + A_2 J_n(\lambda_2 x) + A_3 J_n(\lambda_3 x) + \dots$$

where

$$27.110. \quad A_k = \frac{2}{J_{n+1}^2(\lambda_k)} \int_0^1 xf(x) J_n(\lambda_k x) dx$$

In particular if  $n = 0$ ,

$$27.111. \quad f(x) = A_1 J_0(\lambda_1 x) + A_2 J_0(\lambda_2 x) + A_3 J_0(\lambda_3 x) + \dots$$

where

$$27.112. \quad A_k = \frac{2}{J_1^2(\lambda_k)} \int_0^1 xf(x) J_0(\lambda_k x) dx$$

$R/S > -n$ 

$$27.113. \quad f(x) = A_1 J_n(\lambda_1 x) + A_2 J_n(\lambda_2 x) + A_3 J_n(\lambda_3 x) + \dots$$

where

$$27.114. \quad A_k = \frac{2}{J_n^2(\lambda_k) - J_{n-1}(\lambda_k)J_{n+1}(\lambda_k)} \int_0^1 x f(x) J_n(\lambda_k x) dx$$

In particular if  $n = 0$ .

$$27.115. \quad f(x) = A_1 J_0(\lambda_1 x) + A_2 J_0(\lambda_2 x) + A_3 J_0(\lambda_3 x) + \dots$$

where

$$27.116. \quad A_k = \frac{2}{J_0^2(\lambda_k) + J_1^2(\lambda_k)} \int_0^1 x f(x) J_0(\lambda_k x) dx$$

The next formulas refer to the expansion of Bessel functions where  $S \neq 0$ .

 $R/S = -n$ 

$$27.117. \quad f(x) = A_0 x^n + A_1 J_n(\lambda_1 x) + A_2 J_n(\lambda_2 x) + \dots$$

where

$$27.118. \quad \begin{cases} A_0 = 2(n+1) \int_0^1 x^{n+1} f(x) dx \\ A_k = \frac{2}{J_n^2(\lambda_k) - J_{n-1}(\lambda_k)J_{n+1}(\lambda_k)} \int_0^1 x f(x) J_n(\lambda_k x) dx \end{cases}$$

In particular if  $n = 0$  so that  $R = 0$  [i.e.,  $\lambda_1, \lambda_2, \lambda_3, \dots$  are the positive roots of  $J_1(x) = 0$ ],

$$27.119. \quad f(x) = A_0 + A_1 J_0(\lambda_1 x) + A_2 J_0(\lambda_2 x) + \dots$$

where

$$27.120. \quad \begin{cases} A_0 = 2 \int_0^1 x f(x) dx \\ A_k = \frac{2}{J_0^2(\lambda_k)} \int_0^1 x f(x) J_0(\lambda_k x) dx \end{cases}$$

 $R/S < -N$ 

In this case there are two pure imaginary roots  $\pm i\lambda_0$  as well as the positive roots  $\lambda_1, \lambda_2, \lambda_3, \dots$  and we have

$$27.121. \quad f(x) = A_0 I_n(\lambda_0 x) + A_1 J_n(\lambda_1 x) + A_2 J_n(\lambda_2 x) + \dots$$

where

$$27.122. \quad \begin{cases} A_0 = \frac{2}{I_n^2(\lambda_0) + I_{n-1}(\lambda_0)I_{n+1}(\lambda_0)} \int_0^1 x f(x) I_n(\lambda_0 x) dx \\ A_k = \frac{2}{J_n^2(\lambda_k) - J_{n-1}(\lambda_k)J_{n+1}(\lambda_k)} \int_0^1 x f(x) J_n(\lambda_k x) dx \end{cases}$$

### Miscellaneous Results

---

**27.123.**  $\cos(x \sin \theta) = J_0(x) + 2J_2(x) \cos 2\theta + 2J_4(x) \cos 4\theta + \dots$

**27.124.**  $\sin(x \sin \theta) = 2J_1(x) \sin \theta + 2J_3(x) \sin 3\theta + 2J_5(x) \sin 5\theta + \dots$

**27.125.**  $J_n(x+y) = \sum_{k=-\infty}^{\infty} J_k(x) J_{n-k}(y) \quad n = 0, \pm 1, \pm 2, \dots$

This is called the *addition formula* for Bessel functions.

**27.126.**  $1 = J_0(x) + 2J_2(x) + \dots + 2J_{2n}(x) + \dots$

**27.127.**  $x = 2\{J_1(x) + 3J_3(x) + 5J_5(x) + \dots + (2n+1)J_{2n+1}(x) + \dots\}$

**27.128.**  $x^2 = 2\{4J_2(x) + 16J_4(x) + 36J_6(x) + \dots + (2n)^2 J_{2n}(x) + \dots\}$

**27.129.**  $\frac{xJ_1(x)}{4} = J_2(x) - 2J_4(x) + 3J_6(x) - \dots$

**27.130.**  $1 = J_0^2(x) + 2J_1^2(x) + 2J_2^2(x) + 2J_3^2(x) + \dots$

**27.131.**  $J_n''(x) = \frac{1}{4}\{J_{n-2}(x) - 2J_n(x) + J_{n+2}(x)\}$

**27.132.**  $J_n'''(x) = \frac{1}{8}\{J_{n-3}(x) - 3J_{n-1}(x) + 3J_{n+1}(x) - J_{n+3}(x)\}$

Formulas 27.131 and 27.132 can be generalized.

**27.133.**  $J'_n(x)J_{-n}(x) - J'_{-n}J_n(x) = \frac{2 \sin n\pi}{\pi x}$

**27.134.**  $J_n(x)J_{-n+1}(x) + J_{-n}(x)J_{n-1}(x) = \frac{2 \sin n\pi}{\pi x}$

**27.135.**  $J_{n+1}(x)Y_n(x) - J_n(x)Y_{n+1}(x) = J_n(x)Y'_n(x) - J'_n(x)Y_n(x) = \frac{2}{\pi x}$

**27.136.**  $\sin x = 2\{J_1(x) - J_3(x) + J_5(x) - \dots\}$

**27.137.**  $\cos x = J_0(x) - 2J_2(x) + 2J_4(x) - \dots$

**27.138.**  $\sinh x = 2\{I_1(x) + I_3(x) + I_5(x) + \dots\}$

**27.139.**  $\cosh x = I_0(x) + 2\{I_2(x) + I_4(x) + I_6(x) + \dots\}$

# 28

## LEGENDRE and ASSOCIATED LEGENDRE FUNCTIONS

### Legendre's Differential Equation

---

$$28.1. \quad (1-x^2)y'' - 2xy' + n(n+1)y = 0$$

Solutions of this equation are called *Legendre functions of order n*.

### Legendre Polynomials

---

If  $n = 0, 1, 2, \dots$ , a solution of 28.1 is the Legendre polynomial  $P_n(x)$  given by *Rodrigues' formula*

$$28.2. \quad P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n$$

### Special Legendre Polynomials

---

$$28.3. \quad P_0(x) = 1$$

$$28.7. \quad P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3)$$

$$28.4. \quad P_1(x) = x$$

$$28.8. \quad P_5(x) = \frac{1}{8}(63x^5 - 70x^3 + 15x)$$

$$28.5. \quad P_2(x) = \frac{1}{2}(3x^2 - 1)$$

$$28.9. \quad P_6(x) = \frac{1}{16}(231x^6 - 315x^4 + 105x^2 - 5)$$

$$28.6. \quad P_3(x) = \frac{1}{2}(5x^3 - 3x)$$

$$28.10. \quad P_7(x) = \frac{1}{16}(429x^7 - 693x^5 + 315x^3 - 35x)$$

### Legendre Polynomials in Terms of $\theta$ where $x = \cos \theta$

---

$$28.11. \quad P_0(\cos \theta) = 1$$

$$28.12. \quad P_1(\cos \theta) = \cos \theta$$

$$28.13. \quad P_2(\cos \theta) = \frac{1}{4}(1 + 3\cos 2\theta)$$

$$28.14. \quad P_3(\cos \theta) = \frac{1}{8}(3\cos \theta + 5\cos 3\theta)$$

$$28.15. \quad P_4(\cos \theta) = \frac{1}{64}(9 + 20\cos 2\theta + 35\cos 4\theta)$$

28.16.  $P_5(\cos \theta) = \frac{1}{128}(30 \cos \theta + 35 \cos 3\theta + 63 \cos 5\theta)$

28.17.  $P_6(\cos \theta) = \frac{1}{512}(50 + 105 \cos 2\theta + 126 \cos 4\theta + 231 \cos 6\theta)$

28.18.  $P_7(\cos \theta) = \frac{1}{1024}(175 \cos \theta + 189 \cos 3\theta + 231 \cos 5\theta + 429 \cos 7\theta)$

### Generating Function for Legendre Polynomials

---

28.19.  $\frac{1}{\sqrt{1-2tx+t^2}} = \sum_{n=0}^{\infty} P_n(x)t^n$

### Recurrence Formulas for Legendre Polynomials

---

28.20.  $(n+1)P_{n+1}(x) - (2n+1)xP_n(x) + nP_{n-1}(x) = 0$

28.21.  $P'_{n+1}(x) - xP'_n(x) = (n+1)P_n(x)$

28.22.  $xP'_n(x) - P'_{n-1}(x) = nP_n(x)$

28.23.  $P'_{n+1}(x) - P'_{n-1}(x) = (2n+1)P_n(x)$

28.24.  $(x^2 - 1)P'_n(x) - nxP_n(x) - nP_{n-1}(x)$

### Orthogonality of Legendre Polynomials

---

28.25.  $\int_{-1}^1 P_m(x)P_n(x)dx = 0 \quad m \neq n$

28.26.  $\int_{-1}^1 \{P_n(x)\}^2 dx = \frac{2}{2n+1}$

Because of 28.25,  $P_m(x)$  and  $P_n(x)$  are called *orthogonal* in  $-1 \leq x \leq 1$ .

### Orthogonal Series of Legendre Polynomials

---

28.27.  $f(x) = A_0P_0(x) + A_1P_1(x) + A_2P_2(x) + \dots$

where

28.28.  $A_k = \frac{2k+1}{2} \int_{-1}^1 f(x)P_k(x)dx$

### Special Results Involving Legendre Polynomials

---

28.29.  $P_n(1) = 1$

28.30.  $P_n(-1) = (-1)^n$

28.31.  $P_n(-x) = (-1)^n P_n(x)$

28.32.  $P_n(0) = \begin{cases} 0 & n \text{ odd} \\ (-1)^{n/2} \frac{1 \cdot 3 \cdot 5 \cdots (n-1)}{2 \cdot 4 \cdot 6 \cdots n} & n \text{ even} \end{cases}$

28.33.  $P_n(x) = \frac{1}{\pi} \int_0^\pi (x + \sqrt{x^2 - 1} \cos \phi)^n d\phi$

28.34.  $\int P_n(x) dx = \frac{P_{n+1}(x) - P_{n-1}(x)}{2n+1}$

28.35.  $|P_n(x)| \leq 1$

28.36.  $P_n(x) = \frac{1}{2^{n+1} \pi i} \oint_C \frac{(z^2 - 1)^n}{(z - x)^{n+1}} dz$

where  $C$  is a simple closed curve having  $x$  as interior point.

### General Solution of Legendre's Equation

---

The general solution of Legendre's equation is

28.37.  $y = AU_n(x) + BV_n(x)$

where

28.38.  $U_n(x) = 1 - \frac{n(n+1)}{2!} x^2 + \frac{n(n-2)(n+1)(n+3)}{4!} x^4 - \dots$

28.39.  $V_n(x) = x - \frac{(n-1)(n+2)}{3!} x^3 + \frac{(n-1)(n-3)(n+2)(n+4)}{5!} x^5 - \dots$

These series converge for  $-1 < x < 1$ .

### Legendre Functions of the Second Kind

---

If  $n = 0, 1, 2, \dots$  one of the series 28.38, 28.39 terminates. In such cases,

28.40.  $P_n(x) = \begin{cases} U_n(x)/U_n(1) & n = 0, 2, 4, \dots \\ V_n(x)/V_n(1) & n = 1, 3, 5, \dots \end{cases}$

where

28.41.  $U_n(1) = (-1)^{n/2} 2^n \left[ \left( \frac{n}{2} \right)! \right]^2 / n! \quad n = 0, 2, 4, \dots$

$$28.42. \quad V_n(1) = (-1)^{(n-1)/2} 2^{n-1} \left[ \left( \frac{n-1}{2} \right)! \right]^2 / n! \quad n = 1, 3, 5, \dots$$

The nonterminating series in such a case with a suitable multiplicative constant is denoted by  $Q_n(x)$  and is called *Legendre's function of the second kind of order n*. We define

$$28.43. \quad Q_n(x) = \begin{cases} U_n(1)V_n(x) & n = 0, 2, 4, \dots \\ -V_n(1)U_n(x) & n = 1, 3, 5, \dots \end{cases}$$

### Special Legendre Functions of the Second Kind

---

$$28.44. \quad Q_0(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$$

$$28.45. \quad Q_1(x) = \frac{x}{2} \ln \left( \frac{1+x}{1-x} \right) - 1$$

$$28.46. \quad Q_2(x) = \frac{3x^2 - 1}{4} \ln \left( \frac{1+x}{1-x} \right) - \frac{3x}{2}$$

$$28.47. \quad Q_3(x) = \frac{5x^3 - 3x}{4} \ln \left( \frac{1+x}{1-x} \right) - \frac{5x^2}{2} + \frac{2}{3}$$

The functions  $Q_n(x)$  satisfy recurrence formulas exactly analogous to 28.20 through 28.24. Using these, the general solution of Legendre's equation can also be written as

$$28.48. \quad y = AP_n(x) + BQ_n(x)$$

### Legendre's Associated Differential Equation

---

$$28.49. \quad (1-x^2)y'' - 2xy' + \left\{ n(n+1) - \frac{m^2}{1-x^2} \right\} y = 0$$

Solutions of this equation are called *associated Legendre functions*. We restrict ourselves to the important case where  $m, n$  are nonnegative integers.

### Associated Legendre Functions of the First Kind

---

$$28.50. \quad P_n^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} P_n(x) = \frac{(1-x^2)^{m/2}}{2^n n!} \frac{d^{m+n}}{dx^{m+n}} (x^2 - 1)^n$$

where  $P_n(x)$  are Legendre polynomials (page 164). We have

$$28.51. \quad P_n^0(x) = P_n(x)$$

$$28.52. \quad P_n^m(x) = 0 \quad \text{if } m > n$$

### Special Associated Legendre Functions of the First Kind

---

$$28.53. \quad P_1^1(x) = (1-x^2)^{1/2}$$

$$28.56. \quad P_3^1(x) = \frac{3}{2}(5x^2 - 1)(1-x^2)^{1/2}$$

$$28.54. \quad P_2^1(x) = 3x(1-x^2)^{1/2}$$

$$28.57. \quad P_3^2(x) = 15x(1-x^2)$$

$$28.55. \quad P_2^2(x) = 3(1-x^2)$$

$$28.58. \quad P_3^3(x) = 15(1-x^2)^{3/2}$$

### Generating Function for $P_n^m(x)$

---

$$28.59. \quad \frac{(2m)!(1-x^2)^{m/2}t^m}{2^m m!(1-2tx+t^2)^{m+1/2}} = \sum_{n=m}^{\infty} P_n^m(x)t^n$$

### Recurrence Formulas

---

$$28.60. \quad (n+1-m)P_{n+1}^m(x) - (2n+1)xP_n^m(x) + (n+m)P_{n-1}^m(x) = 0$$

$$28.61. \quad P_n^{m+2}(x) - \frac{2(m+1)x}{(1-x^2)^{1/2}} P_n^{m+1}(x) + (n-m)(n+m+1)P_n^m(x) = 0$$

### Orthogonality of $P_n^m(x)$

---

$$28.62. \quad \int_{-1}^1 P_l^m(x)P_l^m(x) dx = 0 \quad \text{if } n \neq l$$

$$28.63. \quad \int_{-1}^1 \{P_n^m(x)\}^2 dx = \frac{2}{2n+1} \frac{(n+m)!}{(n-m)!}$$

### Orthogonal Series

---

$$28.64. \quad f(x) = A_m P_m^m(x) + A_{m+1} P_{m+1}^m(x) + A_{m+2} P_{m+2}^m(x) + \dots$$

where

$$28.65. \quad A_k = \frac{2k+1}{2} \frac{(k-m)!}{(k+m)!} \int_{-1}^1 f(x)P_k^m(x) dx$$

### Associated Legendre Functions of the Second Kind

---

$$28.66. \quad Q_n^m(x) = (1-x^2)^{m/2} \frac{d^m}{dx^m} Q_n(x)$$

where  $Q_n(x)$  are Legendre functions of the second kind (page 166).

These functions are unbounded at  $x = \pm 1$ , whereas  $P_n^m(x)$  are bounded at  $x = \pm 1$ .

The functions  $Q_n^m(x)$  satisfy the same recurrence relations as  $P_n^m(x)$  (see 28.60 and 28.61).

### General Solution of Legendre's Associated Equation

---

$$28.67. \quad y = AP_n^m(x) + BQ_n^m(x)$$

# 29 HERMITE POLYNOMIALS

## Hermite's Differential Equation

---

$$29.1. \quad y'' - 2xy' + 2ny = 0$$

## Hermite Polynomials

---

If  $n = 0, 1, 2, \dots$ , then a solution of Hermite's equation is the Hermite polynomial  $H_n(x)$  given by *Rodrigue's formula*.

$$29.2. \quad H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$$

## Special Hermite Polynomials

---

$$29.3. \quad H_0(x) = 1$$

$$29.7. \quad H_4(x) = 16x^4 - 48x^2 + 12$$

$$29.4. \quad H_1(x) = 2x$$

$$29.8. \quad H_5(x) = 32x^5 - 160x^3 + 120x$$

$$29.5. \quad H_2(x) = 4x^2 - 2$$

$$29.9. \quad H_6(x) = 64x^6 - 480x^4 + 720x^2 - 120$$

$$29.6. \quad H_3(x) = 8x^3 - 12x$$

$$29.10. \quad H_7(x) = 128x^7 - 1344x^5 + 3360x^3 - 1680x$$

## Generating Function

---

$$29.11. \quad e^{2tx-t^2} = \sum_{n=0}^{\infty} \frac{H_n(x)t^n}{n!}$$

## Recurrence Formulas

---

$$29.12. \quad H_{n+1}(x) = 2xH_n(x) - 2nH_{n-1}(x)$$

$$29.13. \quad H'_n(x) = 2nH_{n-1}(x)$$

## Orthogonality of Hermite Polynomials

---

$$29.14. \quad \int_{-\infty}^{\infty} e^{-x^2} H_m(x) H_n(x) dx = 0 \quad m \neq n$$

$$29.15. \quad \int_{-\infty}^{\infty} e^{-x^2} \{H_n(x)\}^2 dx = 2^n n! \sqrt{\pi}$$

### Orthogonal Series

---

**29.16.**  $f(x) = A_0 H_0(x) + A_1 H_1(x) + A_2 H_2(x) + \dots$

where

**29.17.**  $A_k = \frac{1}{2^k k! \sqrt{\pi}} \int_{-\infty}^{\infty} e^{-x^2} f(x) H_k(x) dx$

### Special Results

---

**29.18.**  $H_n(x) = (2x)^n - \frac{n(n-1)}{1!}(2x)^{n-2} + \frac{n(n-1)(n-2)(n-3)}{2!}(2x)^{n-4} - \dots$

**29.19.**  $H_n(-x) = (-1)^n H_n(x)$

**29.20.**  $H_{2n-1}(0) = 0$

**29.21.**  $H_{2n}(0) = (-1)^n 2^n \cdot 1 \cdot 3 \cdot 5 \cdots (2n-1)$

**29.22.**  $\int_0^x H_n(t) dt = \frac{H_{n+1}(x)}{2(n+1)} - \frac{H_{n+1}(0)}{2(n+1)}$

**29.23.**  $\frac{d}{dx} \{e^{-x^2} H_n(x)\} = -e^{-x^2} H_{n+1}(x)$

**29.24.**  $\int_0^x e^{-t^2} H_n(t) dt = H_{n-1}(0) - e^{-x^2} H_{n-1}(x)$

**29.25.**  $\int_{-\infty}^{\infty} t^n e^{-t^2} H_n(xt) dt = \sqrt{\pi} n! P_n(x)$

**29.26.**  $H_n(x+y) = \sum_{k=0}^n \frac{1}{2^{n/2}} \binom{n}{k} H_k(x\sqrt{2}) H_{n-k}(y\sqrt{2})$

This is called the *addition formula* for Hermite polynomials.

**29.27.**  $\sum_{k=0}^n \frac{H_k(x) H_k(y)}{2^k k!} = \frac{H_{n+1}(x) H_n(y) - H_n(x) H_{n+1}(y)}{2^{n+1} n!(x-y)}$

# 30

## LAGUERRE and ASSOCIATED LAGUERRE POLYNOMIALS

### Laguerre's Differential Equation

---

$$30.1. \quad xy'' + (1-x)y' + ny = 0$$

### Laguerre Polynomials

---

If  $n = 0, 1, 2, \dots$ , then a solution of Laguerre's equation is the Laguerre polynomial  $L_n(x)$  given by *Rodrigues' formula*

$$30.2. \quad L_n(x) = e^x \frac{d^n}{dx^n} (x^n e^{-x})$$

### Special Laguerre Polynomials

---

$$30.3. \quad L_0(x) = 1$$

$$30.4. \quad L_1(x) = -x + 1$$

$$30.5. \quad L_2(x) = x^2 - 4x + 2$$

$$30.6. \quad L_3(x) = -x^3 + 9x^2 - 18x + 6$$

$$30.7. \quad L_4(x) = x^4 - 16x^3 + 72x^2 - 96x + 24$$

$$30.8. \quad L_5(x) = -x^5 + 25x^4 - 200x^3 + 600x^2 - 600x + 120$$

$$30.9. \quad L_6(x) = x^6 - 36x^5 + 450x^4 - 2400x^3 + 5400x^2 - 4320x + 720$$

$$30.10. \quad L_7(x) = -x^7 + 49x^6 - 882x^5 + 7350x^4 - 29,400x^3 + 52,920x^2 - 35,280x + 5040$$

### Generating Function

---

$$30.11. \quad \frac{e^{-xt/(1-t)}}{1-t} = \sum_{n=0}^{\infty} \frac{L_n(x)t^n}{n!}$$

### Recurrence Formulas

---

**30.12.**  $L_{n+1}(x) - (2n+1-x)L_n(x) + n^2L_{n-1}(x) = 0$

**30.13.**  $L'_n(x) - nL'_{n-1}(x) + nL_{n-1}(x) = 0$

**30.14.**  $xL'_n(x) = nL_n(x) - n^2L_{n-1}(x)$

### Orthogonality of Laguerre Polynomials

---

**30.15.**  $\int_0^\infty e^{-x} L_m(x) L_n(x) dx = 0 \quad m \neq n$

**30.16.**  $\int_0^\infty e^{-x} \{L_n(x)\}^2 dx = (n!)^2$

### Orthogonal Series

---

**30.17.**  $f(x) = A_0 L_0(x) + A_1 L_1(x) + A_2 L_2(x) + \dots$

where

**30.18.**  $A_k = \frac{1}{(k!)^2} \int_0^\infty e^{-x} f(x) L_k(x) dx$

### Special Results

---

**30.19.**  $L_n(0) = n!$

**30.20.**  $\int_0^x L_n(t) dt = L_n(x) - \frac{L_{n+1}(x)}{n+1}$

**30.21.**  $L_n(x) = (-1)^n \left\{ x^n - \frac{n^2 x^{n-1}}{1!} + \frac{n^2(n-1)^2 x^{n-2}}{2!} - \dots - (-1)^n n! \right\}$

**30.22.**  $\int_0^\infty x^p e^{-x} L_n(x) dx = \begin{cases} 0 & \text{if } p < n \\ (-1)^n (n!)^2 & \text{if } p = n \end{cases}$

**30.23.**  $\sum_{k=0}^n \frac{L_k(x) L_k(y)}{(k!)^2} = \frac{L_n(x) L_{n+1}(y) - L_{n+1}(x) L_n(y)}{(n!)^2 (x-y)}$

**30.24.**  $\sum_{k=0}^{\infty} \frac{t^k L_k(x)}{(k!)^2} = e^t J_0(2\sqrt{xt})$

**30.25.**  $L_n(x) = \int_0^\infty u^n e^{x-u} J_0(2\sqrt{xu}) du$

### Laguerre's Associated Differential Equation

---

30.26.  $xy'' + (m+1-x)y' + (n-m)y = 0$

### Associated Laguerre Polynomials

---

Solutions of 30.26 for nonnegative integers  $m$  and  $n$  are given by the associated Laguerre polynomials

30.27.  $L_n^m(x) = \frac{d^m}{dx^m} L_n(x)$

where  $L_n(x)$  are Laguerre polynomials (see page 171).

30.28.  $L_n^0(x) = L_n(x)$

30.29.  $L_n^m(x) = 0 \quad \text{if } m > n$

### Special Associated Laguerre Polynomials

---

30.30.  $L_1^1(x) = -1$

30.35.  $L_3^3(x) = -6$

30.31.  $L_2^1(x) = 2x - 4$

30.36.  $L_4^1(x) = 4x^3 - 48x^2 + 144x - 96$

30.32.  $L_2^2(x) = 2$

30.37.  $L_4^2(x) = 12x^2 - 96x + 144$

30.33.  $L_3^1(x) = -3x^2 + 18x - 18$

30.38.  $L_4^3(x) = 24x - 96$

30.34.  $L_3^2(x) = -6x + 18$

30.39.  $L_4^4(x) = 24$

### Generating Function for $L_n^m(x)$

---

30.40.  $\frac{(-1)^m t^m}{(1-t)^{m+1}} e^{-xt/(1-t)} = \sum_{n=m}^{\infty} \frac{L_n^m(x)}{n!} t^n$

### Recurrence Formulas

---

30.41.  $\frac{n-m+1}{n+1} L_{n+1}^m(x) + (x+m-2n-1) L_n^m(x) + n^2 L_{n-1}^m(x) = 0$

30.42.  $\frac{d}{dx} \{L_n^m(x)\} = L_n^{m+1}(x)$

30.43.  $\frac{d}{dx} \{x^m e^{-x} L_n^m(x)\} = (m-n-1)x^{m-1} e^{-x} L_n^{m-1}(x)$

30.44.  $x \frac{d}{dx} \{L_n^m(x)\} = (x-m)L_n^m(x) + (m-n-1)L_n^{m-1}(x)$

### Orthogonality

---

$$30.45. \int_0^\infty x^m e^{-x} L_n^m(x) L_p^m(x) dx = 0 \quad p \neq n$$

$$30.46. \int_0^\infty x^m e^{-x} \{L_n^m(x)\}^2 dx = \frac{(n!)^3}{(n-m)!}$$

### Orthogonal Series

---

$$30.47. f(x) = A_m L_m^m(x) + A_{m+1} L_{m+1}^m(x) + A_{m+2} L_{m+2}^m(x) + \dots$$

where

$$30.48. A_k = \frac{(k-m)!}{(k!)^3} \int_0^\infty x^m e^{-x} L_k^m(x) f(x) dx$$

### Special Results

---

$$30.49. L_n^m(x) = (-1)^n \frac{n!}{(n-m)!} \left\{ x^{n-m} - \frac{n(n-m)}{1!} x^{n-m-1} + \frac{n(n-1)(n-m)(n-m-1)}{2!} x^{n-m-2} + \dots \right\}$$

$$30.50. \int_0^\infty x^{m+1} e^{-x} \{L_n^m(x)\}^2 dx = \frac{(2n-m+1)(n!)^3}{(n-m)!}$$

# 31 CHEBYSHEV POLYNOMIALS

## Chebyshev's Differential Equation

---

$$31.1. \quad (1-x^2)y'' - xy' + n^2y = 0 \quad n = 0, 1, 2, \dots$$

## Chebyshev Polynomials of the First Kind

---

A solution of 31.1 is given by

$$31.2. \quad T_n(x) = \cos(n \cos^{-1} x) = x^n - \binom{n}{2} x^{n-2}(1-x^2) + \binom{n}{4} x^{n-4}(1-x^2)^2 - \dots$$

## Special Chebyshev Polynomials of The First Kind

---

$$31.3. \quad T_0(x) = 1$$

$$31.7. \quad T_4(x) = 8x^4 - 8x^2 + 1$$

$$31.4. \quad T_1(x) = x$$

$$31.8. \quad T_5(x) = 16x^5 - 20x^3 + 5x$$

$$31.5. \quad T_2(x) = 2x^2 - 1$$

$$31.9. \quad T_6(x) = 32x^6 - 48x^4 + 18x^2 - 1$$

$$31.6. \quad T_3(x) = 4x^3 - 3x$$

$$31.10. \quad T_7(x) = 64x^7 - 112x^5 + 56x^3 - 7x$$

## Generating Function for $T_n(x)$

---

$$31.11. \quad \frac{1-tx}{1-2tx+t^2} = \sum_{n=0}^{\infty} T_n(x)t^n$$

## Special Values

---

$$31.12. \quad T_n(-x) = (-1)^n T_n(x)$$

$$31.14. \quad T_n(-1) = (-1)^n$$

$$31.16. \quad T_{2n+1}(0) = 0$$

$$31.13. \quad T_n(1) = 1$$

$$31.15. \quad T_{2n}(0) = (-1)^n$$

## Recursion Formula for $T_n(x)$

---

$$31.17. \quad T_{n+1}(x) - 2xT_n(x) + T_{n-1}(x) = 0$$

### Orthogonality

---

$$31.18. \int_{-1}^1 \frac{T_m(x)T_n(x)}{\sqrt{1-x^2}} dx = 0 \quad m \neq n$$

$$31.19. \int_{-1}^1 \frac{\{T_n(x)\}^2}{\sqrt{1-x^2}} dx = \begin{cases} \pi & \text{if } n = 0 \\ \pi/2 & \text{if } n = 1, 2, \dots \end{cases}$$

### Orthogonal Series

---

$$31.20. f(x) = \frac{1}{2}A_0 T_0(x) + A_1 T_1(x) + A_2 T_2(x) + \dots$$

where

$$31.21. A_k = \frac{2}{\pi} \int_{-1}^1 \frac{f(x)T_k(x)}{\sqrt{1-x^2}} dx$$

### Chebyshev Polynomials of The Second Kind

---

$$31.22. U_n(x) = \frac{\sin\{(n+1)\cos^{-1}x\}}{\sin(\cos^{-1}x)}$$

$$= \binom{n+1}{1} x^n - \binom{n+1}{3} x^{n-2}(1-x^2) + \binom{n+1}{5} x^{n-4}(1-x^2)^2 - \dots$$

### Special Chebyshev Polynomials of The Second Kind

---

$$31.23. U_0(x) = 1$$

$$31.27. U_4(x) = 16x^4 - 12x^2 + 1$$

$$31.24. U_1(x) = 2x$$

$$31.28. U_5(x) = 32x^5 - 32x^3 + 6x$$

$$31.25. U_2(x) = 4x^2 - 1$$

$$31.29. U_6(x) = 64x^6 - 80x^4 + 24x^2 - 1$$

$$31.26. U_3(x) = 8x^3 - 4x$$

$$31.30. U_7(x) = 128x^7 - 192x^5 + 80x^3 - 8x$$

### Generating Function for $U_n(x)$

---

$$31.31. \frac{1}{1-2tx+t^2} = \sum_{n=0}^{\infty} U_n(x)t^n$$

### Special Values

---

$$31.32. U_n(-x) = (-1)^n U_n(x)$$

$$31.34. U_n(-1) = (-1)^n (n+1)$$

$$31.36. U_{2n+1}(0) = 0$$

$$31.33. U_n(1) = n+1$$

$$31.35. U_{2n}(0) = (-1)^n$$

### Recursion Formula for $U_n(x)$

---

31.37.  $U_{n+1}(x) - 2xU_n(x) + U_{n-1}(x) = 0$

### Orthogonality

---

31.38.  $\int_{-1}^1 \sqrt{1-x^2} U_m(x) U_n(x) dx = 0 \quad m \neq n$

31.39.  $\int_{-1}^1 \sqrt{1-x^2} \{U_n(x)\}^2 dx = \frac{\pi}{2}$

### Orthogonal Series

---

31.40.  $f(x) = A_0 U_0(x) + A_1 U_1(x) + A_2 U_2(x) + \dots$

where

31.41.  $A_k = \frac{2}{\pi} \int_{-1}^1 \sqrt{1-x^2} f(x) U_k(x) dx$

### Relationships Between $T_n(x)$ and $U_n(x)$

---

31.42.  $T_n(x) = U_n(x) - xU_{n-1}(x)$

31.43.  $(1-x^2)U_{n-1}(x) = xT_n(x) - T_{n+1}(x)$

31.44.  $U_n(x) = \frac{1}{\pi} \int_{-1}^1 \frac{T_{n+1}(v) dv}{(v-x)\sqrt{1-v^2}}$

31.45.  $T_n(x) = \frac{1}{\pi} \int_{-1}^1 \frac{\sqrt{1-v^2} U_{n-1}(v) dv}{x-v}$

### General Solution of Chebyshev's Differential Equation

---

31.46.  $y = \begin{cases} AT_n(x) + B\sqrt{1-x^2}U_{n-1}(x) & \text{if } n = 1, 2, 3, \dots \\ A + B\sin^{-1} x & \text{if } n = 0 \end{cases}$

# 32 HYPERGEOMETRIC FUNCTIONS

## Hypergeometric Differential Equation

---

$$32.1. \quad x(1-x)y'' + \{c - (a+b+1)x\}y' - aby = 0$$

## Hypergeometric Functions

---

A solution of 32.1 is given by

$$32.2. \quad F(a, b; c; x) = 1 + \frac{a \cdot b}{1 \cdot c}x + \frac{a(a+1)b(b+1)}{1 \cdot 2 \cdot c(c+1)}x^2 + \frac{a(a+1)(a+2)b(b+1)(b+2)}{1 \cdot 2 \cdot 3 \cdot c(c+1)(c+2)}x^3 + \dots$$

If  $a, b, c$  are real, then the series converges for  $-1 < x < 1$  provided that  $c - (a+b) > -1$ .

## Special Cases

---

$$32.3. \quad F(-p, 1; 1; -x) = (1+x)^p$$

$$32.8. \quad F(\frac{1}{2}, \frac{1}{2}; \frac{3}{2}; x^2) = (\sin^{-1} x)/x$$

$$32.4. \quad F(1, 1; 2; -x) = [\ln(1+x)]/x$$

$$32.9. \quad F(\frac{1}{2}, 1; \frac{3}{2}; -x^2) = (\tan^{-1} x)/x$$

$$32.5. \quad \lim_{n \rightarrow \infty} F(1, n; 1; x/n) = e^x$$

$$32.10. \quad F(1, p; p; x) = 1/(1-x)$$

$$32.6. \quad F(\frac{1}{2}, -\frac{1}{2}; \frac{1}{2}; \sin^2 x) = \cos x$$

$$32.11. \quad F(n+1, -n; 1; (1-x)/2) = P_n(x)$$

$$32.7. \quad F(\frac{1}{2}, 1; 1; \sin^2 x) = \sec x$$

$$32.12. \quad F(n, -n; \frac{1}{2}; (1-x)/2) = T_n(x)$$

## General Solution of The Hypergeometric Equation

---

If  $c, a-b$ , and  $c-a-b$  are all nonintegers, then the general solution valid for  $|x| < 1$  is

$$32.13. \quad y = AF(a, b; c; x) + Bx^{1-c}F(a-c+1, b-c+1; 2-c; x)$$

**Miscellaneous Properties**

---

$$32.14. \quad F(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}$$

$$32.15. \quad \frac{d}{dx} F(a, b; c; x) = \frac{ab}{c} F(a+1, b+1; c+1; x)$$

$$32.16. \quad F(a, b; c; x) = \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 u^{b-1} (1-u)^{c-b-1} (1-ux)^{-a} du$$

$$32.17. \quad F(a, b; c; x) = (1-x)^{c-a-b} F(c-a, c-b; c; x)$$

## Section VIII: Laplace and Fourier Transforms

# 33 LAPLACE TRANSFORMS

### Definition of the Laplace Transform of $F(t)$

$$33.1. \quad \mathcal{L}\{F(t)\} = \int_0^{\infty} e^{-st} F(t) dt = f(s)$$

In general  $f(s)$  will exist for  $s > \alpha$  where  $\alpha$  is some constant.  $\mathcal{L}$  is called the *Laplace transform operator*.

### Definition of the Inverse Laplace Transform of $f(s)$

If  $\mathcal{L}\{F(t)\} = f(s)$ , then we say that  $F(t) = \mathcal{L}^{-1}\{f(s)\}$  is the *inverse Laplace transform* of  $f(s)$ .  $\mathcal{L}^{-1}$  is called the *inverse Laplace transform operator*.

### Complex Inversion Formula

The inverse Laplace transform of  $f(s)$  can be found directly by methods of complex variable theory. The result is

$$33.2. \quad F(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{st} f(s) ds = \frac{1}{2\pi i} \lim_{T \rightarrow \infty} \int_{c-iT}^{c+iT} e^{st} f(s) ds$$

where  $c$  is chosen so that all the singular points of  $f(s)$  lie to the left of the line  $\operatorname{Re}\{s\} = c$  in the complex  $s$  plane.

**Table of General Properties of Laplace Transforms**

	$f(s)$	$F(t)$
33.3.	$af_1(s) + bf_2(s)$	$aF_1(t) + bF_2(t)$
33.4.	$f(s/a)$	$aF(at)$
33.5.	$f(s - a)$	$e^{at}F(t)$
33.6.	$e^{-as}f(s)$	$\mathcal{U}(t-a) = \begin{cases} F(t-a) & t > a \\ 0 & t < a \end{cases}$
33.7.	$sf(s) - F(0)$	$F'(t)$
33.8.	$s^2f(s) - sF(0) - F'(0)$	$F''(t)$
33.9.	$s^n f(s) - s^{n-1}F(0) - s^{n-2}F'(0) - \dots - F^{(n-1)}(0)$	$F^{(n)}(t)$
33.10.	$f'(s)$	$-tF(t)$
33.11.	$f''(s)$	$t^2F(t)$
33.12.	$f^{(n)}(s)$	$(-1)^n t^n F(t)$
33.13.	$\frac{f(s)}{s}$	$\int_0^t F(u) du$
33.14.	$\frac{f(s)}{s^n}$	$\int_0^t \dots \int_0^t F(u) du^n = \int_0^t \frac{(t-u)^{n-1}}{(n-1)!} F(u) du$
33.15.	$f(s)g(s)$	$\int_0^t F(u)G(t-u) du$

	$f(s)$	$F(t)$
33.16.	$\int_s^\infty f(u) du$	$\frac{F(t)}{t}$
33.17.	$\frac{1}{1-e^{-sT}} \int_0^T e^{-su} F(u) du$	$F(t) = F(t+T)$
33.18.	$\frac{f(\sqrt{s})}{\sqrt{s}}$	$\frac{1}{\sqrt{\pi t}} \int_0^\infty e^{-u^2/4t} F(u) du$
33.19.	$\frac{1}{s} f\left(\frac{1}{s}\right)$	$\int_0^\infty J_0(2\sqrt{ut}) F(u) du$
33.20.	$\frac{1}{s^{n+1}} f\left(\frac{1}{s}\right)$	$t^{n/2} \int_0^\infty u^{-n/2} J_n(2\sqrt{ut}) F(u) du$
33.21.	$\frac{f(s+1/s)}{s^2 + 1}$	$\int_0^t J_0(2\sqrt{u(t-u)}) F(u) du$
33.22.	$\frac{1}{2\sqrt{\pi}} \int_0^\infty u^{-3/2} e^{-s^2/4u} f(u) du$	$F(t^2)$
33.23.	$\frac{f(\ln s)}{s \ln s}$	$\int_0^\infty \frac{t^u F(u)}{\Gamma(u+1)} du$
33.24.	$\frac{P(s)}{Q(s)}$  where $P(s)$ = polynomial of degree less than $n$ , $Q(s) = (s - \alpha_1)(s - \alpha_2) \dots (s - \alpha_n)$ where $\alpha_1, \alpha_2, \dots, \alpha_n$ are all distinct.	$\sum_{k=1}^n \frac{P(\alpha_k)}{Q'(\alpha_k)} e^{\alpha_k t}$

**Table of Special Laplace Transforms**

	$f(s)$	$F(t)$
33.25.	$\frac{1}{s}$	1
33.26.	$\frac{1}{s^2}$	$t$
33.27.	$\frac{1}{s^n} \quad n = 1, 2, 3, \dots$	$\frac{t^{n-1}}{(n-1)!}, \quad 0! = 1$
33.28.	$\frac{1}{s^n} \quad n > 0$	$\frac{t^{n-1}}{\Gamma(n)}$
33.29.	$\frac{1}{s-a}$	$e^{at}$
33.30.	$\frac{1}{(s-a)^n} \quad n = 1, 2, 3, \dots$	$\frac{t^{n-1}e^{at}}{(n-1)!}, \quad 0! = 1$
33.31.	$\frac{1}{(s-a)^n} \quad n > 0$	$\frac{t^{n-1}e^{at}}{\Gamma(n)}$
33.32.	$\frac{1}{s^2 + a^2}$	$\frac{\sin at}{a}$
33.33.	$\frac{s}{s^2 + a^2}$	$\cos at$
33.34.	$\frac{1}{(s-b)^2 + a^2}$	$\frac{e^{bt} \sin at}{a}$
33.35.	$\frac{s-b}{(s-b)^2 + a^2}$	$e^{bt} \cos at$
33.36.	$\frac{1}{s^2 - a^2}$	$\frac{\sinh at}{a}$
33.37.	$\frac{s}{s^2 - a^2}$	$\cosh at$
33.38.	$\frac{1}{(s-b)^2 - a^2}$	$\frac{e^{bt} \sinh at}{a}$

	$f(s)$	$F(t)$
33.39.	$\frac{s-b}{(s-b)^2 - a^2}$	$e^{bt} \cosh at$
33.40.	$\frac{1}{(s-a)(s-b)} \quad a \neq b$	$\frac{e^{bt} - e^{at}}{b-a}$
33.41.	$\frac{s}{(s-a)(s-b)} \quad a \neq b$	$\frac{be^{bt} - ae^{at}}{b-a}$
33.42.	$\frac{1}{(s^2 + a^2)^2}$	$\frac{\sin at - at \cos at}{2a^3}$
33.43.	$\frac{s}{(s^2 + a^2)^2}$	$\frac{t \sin at}{2a}$
33.44.	$\frac{s^2}{(s^2 + a^2)^2}$	$\frac{\sin at + at \cos at}{2a}$
33.45.	$\frac{s^3}{(s^2 + a^2)^2}$	$\cos at - \frac{1}{2}at \sin at$
33.46.	$\frac{s^2 - a^2}{(s^2 + a^2)^2}$	$t \cos at$
33.47.	$\frac{1}{(s^2 - a^2)^2}$	$\frac{at \cosh at - \sinh at}{2a^3}$
33.48.	$\frac{s}{(s^2 - a^2)^2}$	$\frac{t \sinh at}{2a}$
33.49.	$\frac{s^2}{(s^2 - a^2)^2}$	$\frac{\sinh at + at \cosh at}{2a}$
33.50.	$\frac{s^3}{(s^2 - a^2)^2}$	$\cosh at + \frac{1}{2}at \sinh at$
33.51.	$\frac{s^2}{(s^2 - a^2)^{3/2}}$	$t \cosh at$
33.52.	$\frac{1}{(s^2 + a^2)^3}$	$\frac{(3 - a^2 t^2) \sin at - 3at \cos at}{8a^5}$
33.53.	$\frac{s}{(s^2 + a^2)^3}$	$\frac{t \sin at - at^2 \cos at}{8a^3}$
33.54.	$\frac{s^2}{(s^2 + a^2)^3}$	$\frac{(1 + a^2 t^2) \sin at - at \cos at}{8a^3}$
33.55.	$\frac{s^3}{(s^2 + a^2)^3}$	$\frac{3t \sin at + at^2 \cos at}{8a}$

	$f(s)$	$F(t)$
33.56.	$\frac{s^4}{(s^2 + a^2)^3}$	$\frac{(3 - a^2 t^2) \sin at + 5at \cos at}{8a}$
33.57.	$\frac{s^5}{(s^2 + a^2)^3}$	$\frac{(8 - a^2 t^2) \cos at - 7at \sin at}{8}$
33.58.	$\frac{3s^2 - a^2}{(s^2 + a^2)^3}$	$\frac{t^2 \sin at}{2a}$
33.59.	$\frac{s^3 - 3a^2 s}{(s^2 + a^2)^3}$	$\frac{1}{2} t^2 \cos at$
33.60.	$\frac{s^4 - 6a^2 s^2 + a^4}{(s^2 + a^2)^4}$	$\frac{1}{6} t^3 \cos at$
33.61.	$\frac{s^3 - a^2 s}{(s^2 + a^2)^4}$	$\frac{t^3 \sin at}{24a}$
33.62.	$\frac{1}{(s^2 - a^2)^3}$	$\frac{(3 + a^2 t^2) \sinh at - 3at \cosh at}{8a^5}$
33.63.	$\frac{s}{(s^2 - a^2)^3}$	$\frac{at^2 \cosh at - t \sinh at}{8a^3}$
33.64.	$\frac{s^2}{(s^2 - a^2)^3}$	$\frac{at \cosh at + (a^2 t^2 - 1) \sinh at}{8a^3}$
33.65.	$\frac{s^3}{(s^2 - a^2)^3}$	$\frac{3t \sinh at + at^2 \cosh at}{8a}$
33.66.	$\frac{s^4}{(s^2 - a^2)^3}$	$\frac{(3 + a^2 t^2) \sinh at + 5at \cosh at}{8a}$
33.67.	$\frac{s^5}{(s^2 - a^2)^3}$	$\frac{(8 + a^2 t^2) \cosh at + 7at \sinh at}{8}$
33.68.	$\frac{3s^2 + a^2}{(s^2 - a^2)^3}$	$\frac{t^2 \sinh at}{2a}$
33.69.	$\frac{s^3 + 3a^2 s}{(s^2 - a^2)^3}$	$\frac{1}{2} t^2 \cosh at$
33.70.	$\frac{s^4 + 6a^2 s^2 + a^4}{(s^2 - a^2)^4}$	$\frac{1}{6} t^3 \cosh at$
33.71.	$\frac{s^3 + a^2 s}{(s^2 - a^2)^4}$	$\frac{t^3 \sinh at}{24a}$
33.72.	$\frac{1}{s^3 + a^3}$	$\frac{e^{at/2}}{3a^2} \left\{ \sqrt{3} \sin \frac{\sqrt{3}at}{2} - \cos \frac{\sqrt{3}at}{2} + e^{-3at/2} \right\}$






	$f(s)$	$F(t)$
33.73.	$\frac{s}{s^3 + a^3}$	$\frac{e^{at/2}}{3a} \left\{ \cos \frac{\sqrt{3}at}{2} + \sqrt{3} \sin \frac{\sqrt{3}at}{2} - e^{-3at/2} \right\}$
33.74.	$\frac{s^2}{s^3 + a^3}$	$\frac{1}{3} \left( e^{-at} + 2e^{at/2} \cos \frac{\sqrt{3}at}{2} \right)$
33.75.	$\frac{1}{s^3 - a^3}$	$\frac{e^{-at/2}}{3a^2} \left\{ e^{3at/2} - \cos \frac{\sqrt{3}at}{2} - \sqrt{3} \sin \frac{\sqrt{3}at}{2} \right\}$
33.76.	$\frac{s}{s^3 - a^3}$	$\frac{e^{-at/2}}{3a} \left\{ \sqrt{3} \sin \frac{\sqrt{3}at}{2} - \cos \frac{\sqrt{3}at}{2} + e^{3at/2} \right\}$
33.77.	$\frac{s^2}{s^3 - a^3}$	$\frac{1}{3} \left( e^{at} + 2e^{-at/2} \cos \frac{\sqrt{3}at}{2} \right)$
33.78.	$\frac{1}{s^4 + 4a^4}$	$\frac{1}{4a^3} (\sin at \cosh at - \cos at \sinh at)$
33.79.	$\frac{s}{s^4 + 4a^4}$	$\frac{\sin at \sinh at}{2a^2}$
33.80.	$\frac{s^2}{s^4 + 4a^4}$	$\frac{1}{2a} (\sin at \cosh at + \cos at \sinh at)$
33.81.	$\frac{s^3}{s^4 + 4a^4}$	$\cos at \cosh at$
33.82.	$\frac{1}{s^4 - a^4}$	$\frac{1}{2a^3} (\sinh at - \sin at)$
33.83.	$\frac{s}{s^4 - a^4}$	$\frac{1}{2a^2} (\cosh at - \cos at)$
33.84.	$\frac{s^2}{s^4 - a^4}$	$\frac{1}{2a} (\sinh at + \sin at)$
33.85.	$\frac{s^3}{s^4 - a^4}$	$\frac{1}{2} (\cosh at + \cos at)$
33.86.	$\frac{1}{\sqrt{s+a} + \sqrt{s+b}}$	$\frac{e^{-bt} - e^{-at}}{2(b-a)\sqrt{\pi t^3}}$
33.87.	$\frac{1}{s\sqrt{s+a}}$	$\frac{\operatorname{erf}\sqrt{at}}{\sqrt{a}}$
33.88.	$\frac{1}{\sqrt{s}(s-a)}$	$\frac{e^{at} \operatorname{erf}\sqrt{at}}{\sqrt{a}}$
33.89.	$\frac{1}{\sqrt{s-a} + b}$	$e^{at} \left\{ \frac{1}{\sqrt{\pi t}} - b e^{b^2 t} \operatorname{erfc}(b\sqrt{t}) \right\}$







	$f(s)$	$F(t)$
33.90.	$\frac{1}{\sqrt{s^2 + a^2}}$	$J_0(at)$
33.91.	$\frac{1}{\sqrt{s^2 - a^2}}$	$I_0(at)$
33.92.	$\frac{(\sqrt{s^2 + a^2} - s)^n}{\sqrt{s^2 + a^2}} \quad n > -1$	$a^n J_n(at)$
33.93.	$\frac{(s - \sqrt{s^2 - a^2})^n}{\sqrt{s^2 - a^2}} \quad n > -1$	$a^n I_n(at)$
33.94.	$\frac{e^{b(s-\sqrt{s^2+a^2})}}{\sqrt{s^2+a^2}}$	$J_0(a\sqrt{t(t+2b)})$
33.95.	$\frac{e^{-b\sqrt{s^2+a^2}}}{\sqrt{s^2+a^2}}$	$\begin{cases} J_0(a\sqrt{t^2-b^2}) & t > b \\ 0 & t < b \end{cases}$
33.96.	$\frac{1}{(s^2 + a^2)^{3/2}}$	$\frac{tJ_1(at)}{a}$
33.97.	$\frac{s}{(s^2 + a^2)^{3/2}}$	$tJ_0(at)$
33.98.	$\frac{s^2}{(s^2 + a^2)^{3/2}}$	$J_0(at) - atJ_1(at)$
33.99.	$\frac{1}{(s^2 - a^2)^{3/2}}$	$\frac{tI_1(at)}{a}$
33.100.	$\frac{s}{(s^2 - a^2)^{3/2}}$	$tI_0(at)$
33.101.	$\frac{s^2}{(s^2 - a^2)^{3/2}}$	$I_0(at) + atI_1(at)$
33.102.	$\frac{1}{s(e^s - 1)} = \frac{e^{-s}}{s(1 - e^{-s})}$ See also entry 33.165.	$F(t) = n, n \leq t < n+1, n = 0, 1, 2, \dots$
33.103.	$\frac{1}{s(e^s - r)} = \frac{e^{-s}}{s(1 - re^{-s})}$	$F(t) = \sum_{k=1}^{[t]} r^k$ where $[t] = \text{greatest integer } \leq t$
33.104.	$\frac{e^s - 1}{s(e^s - r)} = \frac{1 - e^{-s}}{s(1 - re^{-s})}$ See also entry 33.167.	$F(t) = r^n, n \leq t < n+1, n = 0, 1, 2, \dots$
33.105.	$\frac{e^{-als}}{\sqrt{s}}$	$\frac{\cos 2\sqrt{at}}{\sqrt{\pi t}}$

	$f(s)$	$F(t)$
33.106.	$\frac{e^{-as}}{s^{3/2}}$	$\frac{\sin 2\sqrt{at}}{\sqrt{\pi a}}$
33.107.	$\frac{e^{-as}}{s^{n+1}} \quad n > -1$	$\left(\frac{t}{a}\right)^{n/2} J_n(2\sqrt{at})$
33.108.	$\frac{e^{-a\sqrt{s}}}{\sqrt{s}}$	$\frac{e^{-a^2/4t}}{\sqrt{\pi t}}$
33.109.	$e^{-a\sqrt{s}}$	$\frac{a}{2\sqrt{\pi t^3}} e^{-a^2/4t}$
33.110.	$\frac{1-e^{-a\sqrt{s}}}{s}$	$\text{erf}(a/2\sqrt{t})$
33.111.	$\frac{e^{-a\sqrt{s}}}{s}$	$\text{erfc}(a/2\sqrt{t})$
33.112.	$\frac{e^{-a\sqrt{s}}}{\sqrt{s}(\sqrt{s} + b)}$	$e^{b(bt+a)} \text{erfc}\left(b\sqrt{t} + \frac{a}{2\sqrt{t}}\right)$
33.113.	$\frac{e^{-a/\sqrt{s}}}{s^{n+1}} \quad n > -1$	$\frac{1}{\sqrt{\pi t} a^{2n+1}} \int_0^\infty u^n e^{-u^2/4a^2t} J_{2n}(2\sqrt{u}) du$
33.114.	$\ln\left(\frac{s+a}{s+b}\right)$	$\frac{e^{-bt} - e^{-at}}{t}$
33.115.	$\frac{\ln[(s^2 + a^2)/a^2]}{2s}$	$Ci(at)$
33.116.	$\frac{\ln[(s+a)/a]}{s}$	$Ei(at)$
33.117.	$-\frac{(\gamma + \ln s)}{s}$ $\gamma = \text{Euler's constant} = .5772156 \dots$	$\ln t$
33.118.	$\ln\left(\frac{s^2 + a^2}{s^2 + b^2}\right)$	$\frac{2(\cos at - \cos bt)}{t}$
33.119.	$\frac{\pi^2}{6s} + \frac{(\gamma + \ln s)^2}{s}$ $\gamma = \text{Euler's constant} = .5772156 \dots$	$\ln^2 t$
33.120.	$\frac{\ln s}{s}$ $\gamma = \text{Euler's constant} = .5772156 \dots$	$-(\ln t + \gamma)$
33.121.	$\frac{\ln^2 s}{s}$ $\gamma = \text{Euler's constant} = .5772156 \dots$	$(\ln t + \gamma)^2 - \frac{1}{6}\pi^2$

	$f(s)$	$F(t)$
33.122.	$\frac{\Gamma'(n+1) - \Gamma(n+1)\ln s}{s^{n+1}} \quad n > -1$	$t^n \ln t$
33.123.	$\tan^{-1}(a/s)$	$\frac{\sin at}{t}$
33.124.	$\frac{\tan^{-1}(a/s)}{s}$	$Si(at)$
33.125.	$\frac{e^{als}}{\sqrt{s}} \operatorname{erfc}(\sqrt{a/s})$	$\frac{e^{-2\sqrt{at}}}{\sqrt{\pi t}}$
33.126.	$e^{s^2/4a^2} \operatorname{erfc}(s/2a)$	$\frac{2a}{\sqrt{\pi}} e^{-a^2 t^2}$
33.127.	$\frac{e^{s^2/4a^2} \operatorname{erfc}(s/2a)}{s}$	$\operatorname{erf}(at)$
33.128.	$\frac{e^{as} \operatorname{erfc}\sqrt{as}}{\sqrt{s}}$	$\frac{1}{\sqrt{\pi(t+a)}}$
33.129.	$e^{as} Ei(as)$	$\frac{1}{t+a}$
33.130.	$\frac{1}{a} \left[ \cos as \left\{ \frac{\pi}{2} - Si(as) \right\} - \sin as Ci(as) \right]$	$\frac{1}{t^2 + a^2}$
33.131.	$\sin as \left\{ \frac{\pi}{2} - Si(as) \right\} + \cos as Ci(as)$	$\frac{t}{t^2 + a^2}$
33.132.	$\frac{\cos as \left\{ \frac{\pi}{2} - Si(as) \right\} - \sin as Ci(as)}{s}$	$\tan^{-1}(t/a)$
33.133.	$\frac{\sin as \left\{ \frac{\pi}{2} - Si(as) \right\} - \cos as Ci(as)}{s}$	$\frac{1}{2} \ln \left( \frac{t^2 + a^2}{a^2} \right)$
33.134.	$\left[ \frac{\pi}{2} - Si(as) \right]^2 + Ci^2(as)$	$\frac{1}{t} \ln \left( \frac{t^2 + a^2}{a^2} \right)$
33.135.	0	$\mathcal{N}(t) = \text{null function}$
33.136.	1	$\delta(t) = \text{delta function}$
33.137.	$e^{-as}$	$\delta(t-a)$
33.138.	$\frac{e^{-as}}{s}$ See also entry 33.163.	$\mathcal{U}(t-a)$

	$f(s)$	$F(t)$
33.139.	$\frac{\sinh sx}{s \sinh sa}$	$\frac{x}{a} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \sin \frac{n\pi x}{a} \cos \frac{n\pi t}{a}$
33.140.	$\frac{\sinh sx}{s \cosh sa}$	$\frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} \sin \frac{(2n-1)\pi x}{2a} \sin \frac{(2n-1)\pi t}{2a}$
33.141.	$\frac{\cosh sx}{s \sinh as}$	$\frac{t}{a} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \cos \frac{n\pi x}{a} \sin \frac{n\pi t}{a}$
33.142.	$\frac{\cosh sx}{s \cosh sa}$	$1 + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} \cos \frac{(2n-1)\pi x}{2a} \cos \frac{(2n-1)\pi t}{2a}$
33.143.	$\frac{\sinh sx}{s^2 \sinh sa}$	$\frac{xt}{a} + \frac{2a}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \sin \frac{n\pi x}{a} \sin \frac{n\pi t}{a}$
33.144.	$\frac{\sinh sx}{s^2 \cosh sa}$	$x + \frac{8a}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)^2} \sin \frac{(2n-1)\pi x}{2a} \cos \frac{(2n-1)\pi t}{2a}$
33.145.	$\frac{\cosh sx}{s^2 \sinh sa}$	$\frac{t^2}{2a} + \frac{2a}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^2} \cos \frac{n\pi x}{a} \left(1 - \cos \frac{n\pi t}{a}\right)$
33.146.	$\frac{\cosh sx}{s^2 \cosh sa}$	$t + \frac{8a}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)^2} \cos \frac{(2n-1)\pi x}{2a} \sin \frac{(2n-1)\pi t}{2a}$
33.147.	$\frac{\cosh sx}{s^3 \cosh sa}$	$\frac{1}{2}(t^2 + x^2 - a^2) - \frac{16a^2}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)^3} \cos \frac{(2n-1)\pi x}{2a} \cos \frac{(2n-1)\pi t}{2a}$
33.148.	$\frac{\sinh x\sqrt{s}}{\sinh a\sqrt{s}}$	$\frac{2\pi}{a^2} \sum_{n=1}^{\infty} (-1)^n n e^{-n^2 \pi^2 t/a^2} \sin \frac{n\pi x}{a}$
33.149.	$\frac{\cosh x\sqrt{s}}{\cosh a\sqrt{s}}$	$\frac{\pi}{a^2} \sum_{n=1}^{\infty} (-1)^{n-1} (2n-1) e^{-(2n-1)^2 \pi^2 t/4a^2} \cos \frac{(2n-1)\pi x}{2a}$
33.150.	$\frac{\sinh x\sqrt{s}}{\sqrt{s} \cosh a\sqrt{s}}$	$\frac{2}{a} \sum_{n=1}^{\infty} (-1)^{n-1} e^{-(2n-1)^2 \pi^2 t/4a^2} \sin \frac{(2n-1)\pi x}{2a}$
33.151.	$\frac{\cosh x\sqrt{s}}{\sqrt{s} \sinh a\sqrt{s}}$	$\frac{1}{a} + \frac{2}{a} \sum_{n=1}^{\infty} (-1)^n e^{-n^2 \pi^2 t/a^2} \cos \frac{n\pi x}{a}$
33.152.	$\frac{\sinh x\sqrt{s}}{s \sinh a\sqrt{s}}$	$\frac{x}{a} + \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{n} e^{-n^2 \pi^2 t/a^2} \sin \frac{n\pi x}{a}$
33.153.	$\frac{\cosh x\sqrt{s}}{s \cosh a\sqrt{s}}$	$1 + \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} e^{-(2n-1)^2 \pi^2 t/4a^2} \cos \frac{(2n-1)\pi x}{2a}$
33.154.	$\frac{\sinh x\sqrt{s}}{s^2 \sinh a\sqrt{s}}$	$\frac{xt}{a} + \frac{2a^2}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^3} (1 - e^{-n^2 \pi^2 t/a^2}) \sin \frac{n\pi x}{a}$
33.155.	$\frac{\cosh x\sqrt{s}}{s^2 \cosh a\sqrt{s}}$	$\frac{1}{2}(x^2 - a^2) + t - \frac{16a^2}{\pi^3} \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n-1)^3} e^{-(2n-1)^2 \pi^2 t/4a^2} \cos \frac{(2n-1)\pi x}{2a}$

	$f(s)$	$F(t)$
33.156.	$\frac{J_0(ix\sqrt{s})}{sJ_0(ia\sqrt{s})}$	$1 - 2 \sum_{n=1}^{\infty} \frac{e^{-\lambda_n^2 t/a^2} J_0(\lambda_n x/a)}{\lambda_n J_1(\lambda_n)}$ <p>where <math>\lambda_1, \lambda_2, \dots</math> are the positive roots of <math>J_0(\lambda) = 0</math></p>
33.157.	$\frac{J_0(ix\sqrt{s})}{s^2 J_0(ia\sqrt{s})}$	$\frac{1}{4}(x^2 - a^2) + t + 2a^2 \sum_{n=1}^{\infty} \frac{e^{-\lambda_n^2 t/a^2} J_0(\lambda_n x/a)}{\lambda_n^3 J_1(\lambda_n)}$ <p>where <math>\lambda_1, \lambda_2, \dots</math> are the positive roots of <math>J_0(\lambda) = 0</math></p>
33.158.	$\frac{1}{as^2} \tanh\left(\frac{as}{2}\right)$	Triangular wave function  <p>Fig. 33-1</p>
33.159.	$\frac{1}{s} \tanh\left(\frac{as}{2}\right)$	Square wave function  <p>Fig. 33-2</p>
33.160.	$\frac{\pi a}{a^2 s^2 + \pi^2} \coth\left(\frac{as}{2}\right)$	Rectified sine wave function  <p>Fig. 33-3</p>
33.161.	$\frac{\pi a}{(a^2 s^2 + \pi^2)(1 - e^{-as})}$	Half-rectified sine wave function  <p>Fig. 33-4</p>
33.162.	$\frac{1}{as^2} - \frac{e^{-as}}{s(1 - e^{-as})}$	Sawtooth wave function  <p>Fig. 33-5</p>

	$f(s)$	$F(t)$
33.163.	$\frac{e^{-as}}{s}$ See also entry 33.138.	Heaviside's unit function $\mathcal{U}(t - a)$  Fig. 33-6
33.164.	$\frac{e^{-as}(1 - e^{-\epsilon s})}{s}$	Pulse function  Fig. 33-7
33.165.	$\frac{1}{s(1 - e^{-as})}$ See also entry 33.102.	Step function  Fig. 33-8
33.166.	$\frac{e^{-s} + e^{-2s}}{s(1 - e^{-s})^2}$	$F(t) = n^2, n \leq t < n + 1, n = 0, 1, 2, \dots$  Fig. 33-9
33.167.	$\frac{1 - e^{-s}}{s(1 - re^{-s})}$ See also entry 33.104.	$F(t) = r^n, n \leq t < n + 1, n = 0, 1, 2, \dots$  Fig. 33-10
33.168.	$\frac{\pi a(1 + e^{-as})}{a^2 s^2 + \pi^2}$	$F(t) = \begin{cases} \sin(\pi t/a) & 0 \leq t \leq a \\ 0 & t > a \end{cases}$  Fig. 33-11

# 34 FOURIER TRANSFORMS

## Fourier's Integral Theorem

$$34.1. \quad f(x) = \int_0^\infty \{A(\alpha)\cos \alpha x + B(\alpha)\sin \alpha x\}d\alpha$$

where

$$34.2. \quad \begin{cases} A(\alpha) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \cos \alpha x dx \\ B(\alpha) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(x) \sin \alpha x dx \end{cases}$$

Sufficient conditions under which this theorem holds are:

- (i)  $f(x)$  and  $f'(x)$  are piecewise continuous in every finite interval  $-L < x < L$ ;
- (ii)  $\int_{-\infty}^{\infty} |f(x)|dx$  converges;
- (iii)  $f(x)$  is replaced by  $\frac{1}{2}\{f(x+0) + f(x-0)\}$  if  $x$  is a point of discontinuity.

## Equivalent Forms of Fourier's Integral Theorem

$$34.3. \quad f(x) = \frac{1}{2\pi} \int_{\alpha=-\infty}^{\infty} \int_{u=-\infty}^{\infty} f(u) \cos \alpha(x-u) du d\alpha$$

$$34.4. \quad \begin{aligned} f(x) &= \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{i\alpha x} d\alpha \int_{-\infty}^{\infty} f(u) e^{-i\alpha u} du \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(u) e^{i\alpha(x-u)} du d\alpha \end{aligned}$$

$$34.5. \quad f(x) = \frac{2}{\pi} \int_0^\infty \sin \alpha x d\alpha \int_0^\infty f(u) \sin \alpha u du$$

where  $f(x)$  is an *odd function* [ $f(-x) = -f(x)$ ].

$$34.6. \quad f(x) = \frac{2}{\pi} \int_0^\infty \cos \alpha x d\alpha \int_0^\infty f(u) \cos \alpha u du$$

where  $f(x)$  is an *even function* [ $f(-x) = f(x)$ ].

## Fourier Transforms

---

The Fourier transform of  $f(x)$  is defined as

$$34.7. \quad \mathcal{F}\{f(x)\} = F(\alpha) = \int_{-\infty}^{\infty} f(x)e^{-i\alpha x} dx$$

Then from 34.7 the inverse Fourier transform of  $F(\alpha)$  is

$$34.8. \quad \mathcal{F}^{-1}\{F(\alpha)\} = f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\alpha)e^{i\alpha x} d\alpha$$

We call  $f(x)$  and  $F(\alpha)$  *Fourier transform pairs*.

## Convolution Theorem for Fourier Transforms

---

If  $F(\alpha) = \mathcal{F}\{f(x)\}$  and  $G(\alpha) = \mathcal{F}\{g(x)\}$ , then

$$34.9. \quad \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\alpha)G(\alpha)e^{i\alpha x} d\alpha = \int_{-\infty}^{\infty} f(u)g(x-u) du = f^* g$$

where  $f^* g$  is called the *convolution* of  $f$  and  $g$ . Thus,

$$34.10. \quad \mathcal{F}\{f^* g\} = \mathcal{F}\{f\} \mathcal{F}\{g\}$$

## Parseval's Identity

---

If  $F(\alpha) = \mathcal{F}\{f(x)\}$ , then

$$34.11. \quad \int_{-\infty}^{\infty} |f(x)|^2 dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} |F(\alpha)|^2 d\alpha$$

More generally if  $F(\alpha) = \mathcal{F}\{f(x)\}$  and  $G(\alpha) = \mathcal{F}\{g(x)\}$ , then

$$34.12. \quad \int_{-\infty}^{\infty} f(x)\overline{g(x)} dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\alpha)\overline{G(\alpha)} d\alpha$$

where the bar denotes complex conjugate.

## Fourier Sine Transforms

---

The Fourier sine transform of  $f(x)$  is defined as

$$34.13. \quad F_s(\alpha) = \mathcal{F}_s\{f(x)\} = \int_0^{\infty} f(x)\sin \alpha x dx$$

Then from 34.13 the inverse Fourier sine transform of  $F_s(\alpha)$  is

$$34.14. \quad f(x) = \mathcal{F}_s^{-1}\{F_s(\alpha)\} = \frac{2}{\pi} \int_0^{\infty} F_s(\alpha)\sin \alpha x d\alpha$$

### Fourier Cosine Transforms

---

The Fourier cosine transform of  $f(x)$  is defined as

$$34.15. \quad F_C(\alpha) = \mathcal{F}_C\{f(x)\} = \int_0^\infty f(x) \cos \alpha x \, dx$$

Then from 34.15 the inverse Fourier cosine transform of  $F_C(\alpha)$  is

$$34.16. \quad f(x) = \mathcal{F}_C^{-1}\{F_C(\alpha)\} = \frac{2}{\pi} \int_0^\infty F_C(\alpha) \cos \alpha x \, d\alpha$$

### Special Fourier Transform Pairs

---

	$f(x)$	$F(\alpha)$
34.17.	$\begin{cases} 1 &  x  < b \\ 0 &  x  > b \end{cases}$	$\frac{2 \sin b\alpha}{\alpha}$
34.18.	$\frac{1}{x^2 + b^2}$	$\frac{\pi e^{-b\alpha}}{b}$
34.19.	$\frac{x}{x^2 + b^2}$	$-i\pi e^{-b\alpha}$
34.20.	$f^{(n)}(x)$	$i^n \alpha^n F(\alpha)$
34.21.	$x^n f(x)$	$i^n \frac{d^n F}{d\alpha^n}$
34.22.	$f(bx) e^{itx}$	$\frac{1}{b} F\left(\frac{\alpha - t}{b}\right)$

### Special Fourier Sine Transforms

---

	$f(x)$	$F_c(\alpha)$
34.23.	$\begin{cases} 1 & 0 < x < b \\ 0 & x > b \end{cases}$	$\frac{1 - \cos b\alpha}{\alpha}$
34.24.	$x^{-1}$	$\frac{\pi}{2}$
34.25.	$\frac{x}{x^2 + b^2}$	$\frac{\pi}{2} e^{-b\alpha}$
34.26.	$e^{-bx}$	$\frac{\alpha}{\alpha^2 + b^2}$
34.27.	$x^{n-1} e^{-bx}$	$\frac{\Gamma(n) \sin(n \tan^{-1} \alpha/b)}{(\alpha^2 + b^2)^{n/2}}$
34.28.	$x e^{-bx^2}$	$\frac{\sqrt{\pi}}{4b^{3/2}} \alpha e^{-\alpha^2/4b}$
34.29.	$x^{-1/2}$	$\sqrt{\frac{\pi}{2\alpha}}$
34.30.	$x^{-n}$	$\frac{\pi \alpha^{n-1} \csc(n\pi/2)}{2\Gamma(n)} \quad 0 < n < 2$
34.31.	$\frac{\sin bx}{x}$	$\frac{1}{2} \ln\left(\frac{\alpha+b}{\alpha-b}\right)$
34.32.	$\frac{\sin bx}{x^2}$	$\begin{cases} \pi\alpha/2 & \alpha < b \\ \pi b/2 & \alpha > b \end{cases}$
34.33.	$\frac{\cos bx}{x}$	$\begin{cases} 0 & \alpha < b \\ \pi/4 & \alpha = b \\ \pi/2 & \alpha > b \end{cases}$
34.34.	$\tan^{-1}(x/b)$	$\frac{\pi}{2\alpha} e^{-b\alpha}$
34.35.	$\csc bx$	$\frac{\pi}{2b} \tanh \frac{\pi\alpha}{2b}$
34.36.	$\frac{1}{e^{2x}-1}$	$\frac{\pi}{4} \coth\left(\frac{\pi\alpha}{2}\right) - \frac{1}{2\alpha}$

### Special Fourier Cosine Transforms

---

	$f(x)$	$F_c(\alpha)$
34.37.	$\begin{cases} 1 & 0 < x < b \\ 0 & x > b \end{cases}$	$\frac{\sin b\alpha}{\alpha}$
34.38.	$\frac{1}{x^2 + b^2}$	$\frac{\pi e^{-b\alpha}}{2b}$
34.39.	$e^{-bx}$	$\frac{b}{\alpha^2 + b^2}$
34.40.	$x^{n-1}e^{-bx}$	$\frac{\Gamma(n)\cos(n\tan^{-1}\alpha/b)}{(\alpha^2 + b^2)^{n/2}}$
34.41.	$e^{-bx^2}$	$\frac{1}{2}\sqrt{\frac{\pi}{b}}e^{-\alpha^2/4b}$
34.42.	$x^{-1/2}$	$\sqrt{\frac{\pi}{2\alpha}}$
34.43.	$x^{-n}$	$\frac{\pi\alpha^{n-1}\sec(n\pi/2)}{2\Gamma(n)}, \quad 0 < n < 1$
34.44.	$\ln\left(\frac{x^2 + b^2}{x^2 + c^2}\right)$	$\frac{e^{-c\alpha} - e^{-b\alpha}}{\pi\alpha}$
34.45.	$\frac{\sin bx}{x}$	$\begin{cases} \pi/2 & \alpha < b \\ \pi/4 & \alpha = b \\ 0 & \alpha > b \end{cases}$
34.46.	$\sin bx^2$	$\sqrt{\frac{\pi}{8b}}\left(\cos\frac{\alpha^2}{4b} - \sin\frac{\alpha^2}{4b}\right)$
34.47.	$\cos bx^2$	$\sqrt{\frac{\pi}{8b}}\left(\cos\frac{\alpha^2}{4b} + \sin\frac{\alpha^2}{4b}\right)$
34.48.	$\operatorname{sech} bx$	$\frac{\pi}{2b} \operatorname{sech}\frac{\pi\alpha}{2b}$
34.49.	$\frac{\cosh(\sqrt{\pi}x/2)}{\cosh(\sqrt{\pi}x)}$	$\sqrt{\frac{\pi}{2}} \frac{\cosh(\sqrt{\pi}\alpha/2)}{\cosh(\sqrt{\pi}\alpha)}$
34.50.	$\frac{e^{-b\sqrt{x}}}{\sqrt{x}}$	$\sqrt{\frac{\pi}{2\alpha}}\{\cos(2b\sqrt{\alpha}) - \sin(2b\sqrt{\alpha})\}$

## Section IX: Elliptic and Miscellaneous Special Functions

# 35 ELLIPTIC FUNCTIONS

### Incomplete Elliptic Integral of the First Kind

$$35.1. \quad u = F(k, \phi) = \int_0^\phi \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} = \int_0^x \frac{dv}{\sqrt{(1 - v^2)(1 - k^2 v^2)}}$$

where  $\phi = \operatorname{am} u$  is called the *amplitude* of  $u$  and  $x = \sin \phi$ , and where here and below  $0 < k < 1$ .

### Complete Elliptic Integral of the First Kind

$$35.2. \quad K = F(k, \pi/2) = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} = \int_0^1 \frac{dv}{\sqrt{(1 - v^2)(1 - k^2 v^2)}} \\ = \frac{\pi}{2} \left\{ 1 + \left( \frac{1}{2} \right)^2 k^2 + \left( \frac{1 \cdot 3}{2 \cdot 4} \right)^2 k^4 + \left( \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^2 k^6 + \dots \right\}$$

### Incomplete Elliptic Integral of the Second Kind

$$35.3. \quad E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \theta} d\theta = \int_0^x \frac{\sqrt{1 - k^2 v^2}}{\sqrt{1 - v^2}} dv$$

### Complete Elliptic Integral of the Second Kind

$$35.4. \quad E = E(k, \pi/2) = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta = \int_0^1 \frac{\sqrt{1 - k^2 v^2}}{\sqrt{1 - v^2}} dv \\ = \frac{\pi}{2} \left\{ 1 - \left( \frac{1}{2} \right)^2 k^2 - \left( \frac{1 \cdot 3}{2 \cdot 4} \right)^2 \frac{k^4}{3} - \left( \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^2 \frac{k^6}{5} - \dots \right\}$$

### Incomplete Elliptic Integral of the Third Kind

$$35.5. \quad \Pi(k, n, \phi) = \int_0^\phi \frac{d\theta}{(1 + n \sin^2 \theta) \sqrt{1 - k^2 \sin^2 \theta}} = \int_0^x \frac{dv}{(1 + nv^2) \sqrt{(1 - v^2)(1 - k^2 v^2)}}$$

### Complete Elliptic Integral of the Third Kind

---

$$35.6. \quad \Pi(k, n, \pi/2) = \int_0^{\pi/2} \frac{d\theta}{(1+n \sin^2 \theta) \sqrt{1-k^2 \sin^2 \theta}} = \int_0^1 \frac{dv}{(1+nv^2) \sqrt{(1-v^2)(1-k^2 v^2)}}$$

### Landen's Transformation

---

$$35.7. \quad \tan \phi = \frac{\sin 2\phi_1}{k + \cos 2\phi_1} \quad \text{or} \quad k \sin \phi = \sin(2\phi_1 - \phi)$$

This yields

$$35.8. \quad F(k, \phi) = \int_0^\phi \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}} = \frac{2}{1+k} \int_0^{\phi_1} \frac{d\theta_1}{\sqrt{1-k_1^2 \sin^2 \theta_1}}$$

where  $k_1 = 2\sqrt{k}/(1+k)$ . By successive applications, sequences  $k_1, k_2, k_3, \dots$  and  $\phi_1, \phi_2, \phi_3, \dots$  are obtained such that  $k < k_1 < k_2 < k_3 < \dots < 1$  where  $\lim_{n \rightarrow \infty} k_n = 1$ . It follows that

$$35.9. \quad F(k, \Phi) = \sqrt{\frac{k_1 k_2 k_3 \dots}{k}} \int_0^\Phi \frac{d\theta}{\sqrt{1-\sin^2 \theta}} = \sqrt{\frac{k_1 k_2 k_3 \dots}{k}} \ln \tan \left( \frac{\pi}{4} + \frac{\Phi}{2} \right)$$

where

$$35.10. \quad k_1 = \frac{2\sqrt{k}}{1+k}, \quad k_2 = \frac{2\sqrt{k_1}}{1+k_1}, \dots \quad \text{and} \quad \Phi \lim_{n \rightarrow \infty} \phi_n$$

The result is used in the approximate evaluation of  $F(k, \phi)$ .

### Jacobi's Elliptic Functions

---

From 35.1 we define the following elliptic functions:

$$35.11. \quad x = \sin(\operatorname{am} u) = \operatorname{sn} u$$

$$35.12. \quad \sqrt{1-x^2} = \cos(\operatorname{am} u) = \operatorname{cn} u$$

$$35.13. \quad \sqrt{1-k^2 x^2} = \sqrt{1-k^2 \operatorname{sn}^2 u} = \operatorname{dn} u$$

We can also define the inverse functions  $\operatorname{sn}^{-1} x, \operatorname{cn}^{-1} x, \operatorname{dn}^{-1} x$  and the following:

$$35.14. \quad \operatorname{ns} u = \frac{1}{\operatorname{sn} u}$$

$$35.17. \quad \operatorname{sc} u = \frac{\operatorname{sn} u}{\operatorname{cn} u}$$

$$35.20. \quad \operatorname{cs} u = \frac{\operatorname{cn} u}{\operatorname{sn} u}$$

$$35.15. \quad \operatorname{nc} u = \frac{1}{\operatorname{cn} u}$$

$$35.18. \quad \operatorname{sd} u = \frac{\operatorname{sn} u}{\operatorname{dn} u}$$

$$35.21. \quad \operatorname{dc} u = \frac{\operatorname{dn} u}{\operatorname{cn} u}$$

$$35.16. \quad \operatorname{nd} u = \frac{1}{\operatorname{dn} u}$$

$$35.19. \quad \operatorname{cd} u = \frac{\operatorname{cn} u}{\operatorname{dn} u}$$

$$35.22. \quad \operatorname{ds} u = \frac{\operatorname{dn} u}{\operatorname{sn} u}$$

### Addition Formulas

---

$$35.23. \quad \operatorname{sn}(u+v) = \frac{\operatorname{sn} u \operatorname{cn} v \operatorname{dn} v + \operatorname{cn} u \operatorname{sn} v \operatorname{dn} u}{1-k^2 \operatorname{sn}^2 u \operatorname{sn}^2 v}$$

$$35.24. \quad \text{cn}(u+v) = \frac{\text{cn } u \text{ cn } v - \text{sn } u \text{ sn } v \text{ dn } u \text{ dn } v}{1 - k^2 \text{sn}^2 u \text{ sn}^2 v}$$

$$35.25. \quad \text{dn}(u+v) = \frac{\text{dn } u \text{ dn } v - k^2 \text{sn } u \text{ sn } v \text{ cn } u \text{ cn } v}{1 - k^2 \text{sn}^2 u \text{ sn}^2 v}$$

## Derivatives

---

$$35.26. \quad \frac{d}{du} \text{sn } u = \text{cn } u \text{ dn } u$$

$$35.28. \quad \frac{d}{du} \text{dn } u = -k^2 \text{sn } u \text{ cn } u$$

$$35.27. \quad \frac{d}{du} \text{cn } u = -\text{sn } u \text{ dn } u$$

$$35.29. \quad \frac{d}{du} \text{sc } u = \text{dc } u \text{ nc } u$$

## Series Expansions

---

$$35.30. \quad \text{sn } u = u - (1+k^2) \frac{u^3}{3!} + (1+14k^2+k^4) \frac{u^5}{5!} - (1+135k^2+135k^4+k^6) \frac{u^7}{7!} + \dots$$

$$35.31. \quad \text{cn } u = 1 - \frac{u^2}{2!} + (1+4k^2) \frac{u^4}{4!} - (1+44k^2+16k^4) \frac{u^6}{6!} + \dots$$

$$35.32. \quad \text{dn } u = 1 - k^2 \frac{u^2}{2!} + k^2(4+k^2) \frac{u^4}{4!} - k^2(16+44k^2+k^4) \frac{u^6}{6!} + \dots$$

## Catalan's Constant

---

$$35.33. \quad \frac{1}{2} \int_0^1 K dk = \frac{1}{2} \int_{k=0}^1 \int_{\theta=0}^{\pi/2} \frac{d\theta dk}{\sqrt{1-k^2 \sin^2 \theta}} = \frac{1}{1^2} - \frac{1}{3^2} + \frac{1}{5^2} - \dots = .915965594\dots$$

## Periods of Elliptic Functions

---

Let

$$35.34. \quad K = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k^2 \sin^2 \theta}}, \quad K' = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1-k'^2 \sin^2 \theta}} \quad \text{where } k' = \sqrt{1-k^2}$$

Then

$$35.35. \quad \text{sn } u \text{ has periods } 4K \text{ and } 2iK'$$

$$35.36. \quad \text{cn } u \text{ has periods } 4K \text{ and } 2K + 2iK'$$

$$35.37. \quad \text{dn } u \text{ has periods } 2K \text{ and } 4iK'$$

**Identities Involving Elliptic Functions**

35.38.  $\operatorname{sn}^2 u + \operatorname{cn}^2 u = 1$

35.42.  $\operatorname{cn}^2 u = \frac{\operatorname{dn} 2u + \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}$

35.39.  $\operatorname{dn}^2 u + k^2 \operatorname{sn}^2 u = 1$

35.43.  $\operatorname{dn}^2 u = \frac{1 - k^2 + \operatorname{dn} 2u + k^2 \operatorname{cn} u}{1 + \operatorname{dn} 2u}$

35.40.  $\operatorname{dn}^2 u - k^2 \operatorname{cn}^2 u = k'^2$  where  $k' = \sqrt{1 - k^2}$

35.44.  $\sqrt{\frac{1 - \operatorname{cn} 2u}{1 + \operatorname{cn} 2u}} = \frac{\operatorname{sn} u \operatorname{dn} u}{\operatorname{cn} u}$

35.41.  $\operatorname{sn}^2 u = \frac{1 - \operatorname{cn} 2u}{1 + \operatorname{dn} 2u}$

35.45.  $\sqrt{\frac{1 - \operatorname{dn} 2u}{1 + \operatorname{dn} 2u}} = \frac{k \operatorname{sn} u \operatorname{cn} u}{\operatorname{dn} u}$

**Special Values**

35.46.  $\operatorname{sn} 0 = 0$

35.47.  $\operatorname{cn} 0 = 1$

35.48.  $\operatorname{dn} 0 = 1$

35.49.  $\operatorname{sc} 0 = 0$

35.50.  $\operatorname{am} 0 = 0$

**Integrals**

35.51.  $\int \operatorname{sn} u \, du = \frac{1}{k} \ln(\operatorname{dn} u - k \operatorname{cn} u)$

35.52.  $\int \operatorname{cn} u \, du = \frac{1}{k} \cos^{-1}(\operatorname{dn} u)$

35.53.  $\int \operatorname{dn} u \, du = \sin^{-1}(\operatorname{sn} u)$

35.54.  $\int \operatorname{sc} u \, du = \frac{1}{\sqrt{1 - k^2}} \ln(\operatorname{dc} u + \sqrt{1 - k^2} \operatorname{nc} u)$

35.55.  $\int \operatorname{cs} u \, du = \ln(\operatorname{ns} u - \operatorname{ds} u)$

35.56.  $\int \operatorname{cd} u \, du = \frac{1}{k} \ln(\operatorname{nd} u + k \operatorname{sd} u)$

35.57.  $\int \operatorname{dc} u \, du = \ln(\operatorname{nc} u + \operatorname{sc} u)$

35.58.  $\int \operatorname{sd} u \, du = \frac{-1}{k \sqrt{1 - k^2}} \sin^{-1}(k \operatorname{cd} u)$

35.59.  $\int \operatorname{ds} u \, du = \ln(\operatorname{ns} u - \operatorname{cs} u)$

35.60.  $\int \operatorname{ns} u \, du = \ln(\operatorname{ds} u - \operatorname{cs} u)$

35.61.  $\int \operatorname{nc} u \, du = \frac{1}{\sqrt{1 - k^2}} \ln\left(\operatorname{dc} u + \frac{\operatorname{sc} u}{\sqrt{1 - k^2}}\right)$

35.62.  $\int \operatorname{nd} u \, du = \frac{1}{\sqrt{1 - k^2}} \cos^{-1}(\operatorname{cd} u)$

### Legendre's Relation

---

35.63.  $EK' + E'K - KK' = \pi/2$

where

35.64.  $E = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta$        $K = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}$

35.65.  $E' = \int_0^{\pi/2} \sqrt{1 - k'^2 \sin^2 \theta} d\theta$        $K' = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k'^2 \sin^2 \theta}}$

# 36 MISCELLANEOUS and RIEMANN ZETA FUNCTIONS

**Error Function**  $\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du$

$$36.1. \quad \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \left( x - \frac{x^3}{3 \cdot 1!} + \frac{x^5}{5 \cdot 2!} - \frac{x^7}{7 \cdot 3!} + \dots \right)$$

$$36.2. \quad \operatorname{erf}(x) \sim 1 - \frac{e^{-x^2}}{\sqrt{\pi}x} \left( 1 - \frac{1}{2x^2} + \frac{1 \cdot 3}{(2x^2)^2} - \frac{1 \cdot 3 \cdot 5}{(2x^2)^3} + \dots \right)$$

$$36.3. \quad \operatorname{erf}(-x) = -\operatorname{erf}(x), \quad \operatorname{erf}(0) = 0, \quad \operatorname{erf}(\infty) = 1$$

**Complementary Error Function**  $\operatorname{erfc}(x) = 1 - \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_x^\infty e^{-u^2} du$

$$36.4. \quad \operatorname{erfc}(x) = 1 - \frac{2}{\sqrt{\pi}} \left( x - \frac{x^3}{3 \cdot 1!} + \frac{x^5}{5 \cdot 2!} - \frac{x^7}{7 \cdot 3!} + \dots \right)$$

$$36.5. \quad \operatorname{erfc}(x) \sim \frac{e^{-x^2}}{\sqrt{\pi}x} \left( 1 - \frac{1}{2x^2} + \frac{1 \cdot 3}{(2x^2)^2} - \frac{1 \cdot 3 \cdot 5}{(2x^2)^3} + \dots \right)$$

$$36.6. \quad \operatorname{erfc}(0) = 1, \quad \operatorname{erfc}(\infty) = 0$$

**Exponential Integral**  $\operatorname{Ei}(x) = \int_x^\infty \frac{e^{-u}}{u} du$

$$36.7. \quad \operatorname{Ei}(x) = -\gamma - \ln x + \int_0^x \frac{1 - e^{-u}}{u} du$$

$$36.8. \quad \operatorname{Ei}(x) = -\gamma - \ln x + \left( \frac{x}{1 \cdot 1!} - \frac{x^2}{2 \cdot 2!} + \frac{x^3}{3 \cdot 3!} - \dots \right)$$

$$36.9. \quad \operatorname{Ei}(x) \sim \frac{e^{-x}}{x} \left( 1 - \frac{1!}{x} + \frac{2!}{x^2} - \frac{3!}{x^3} + \dots \right)$$

$$36.10. \quad \operatorname{Ei}(\infty) = 0$$

**Sine Integral**  $\operatorname{Si}(x) = \int_0^x \frac{\sin u}{u} du$

$$36.11. \quad \operatorname{Si}(x) = \frac{x}{1 \cdot 1!} - \frac{x^3}{3 \cdot 3!} + \frac{x^5}{5 \cdot 5!} - \frac{x^7}{7 \cdot 7!} + \dots$$

$$36.12. \quad \operatorname{Si}(x) \sim \frac{\pi}{2} - \frac{\sin x}{x} \left( 1 - \frac{3!}{x^3} + \frac{5!}{x^5} - \dots \right) - \frac{\cos x}{x} \left( 1 - \frac{2!}{x^2} + \frac{4!}{x^4} - \dots \right)$$

$$36.13. \quad \operatorname{Si}(-x) = -\operatorname{Si}(x), \quad \operatorname{Si}(0) = 0, \quad \operatorname{Si}(\infty) = \pi/2$$

**Cosine Integral  $\text{Ci}(x) = \int_x^{\infty} \frac{\cos u}{u} du$**

---

36.14.  $\text{Ci}(x) = -\gamma - \ln x + \int_0^x \frac{1 - \cos u}{u} du$

36.15.  $\text{Ci}(x) = -\gamma - \ln x + \frac{x^2}{2 \cdot 2!} - \frac{x^4}{4 \cdot 4!} + \frac{x^6}{6 \cdot 6!} - \frac{x^8}{8 \cdot 8!} + \dots$

36.16.  $\text{Ci}(x) \sim \frac{\cos x}{x} \left( \frac{1}{x} - \frac{3!}{x^3} + \frac{5!}{x^5} - \dots \right) - \frac{\sin x}{x} \left( 1 - \frac{2!}{x^2} + \frac{4!}{x^4} - \dots \right)$

36.17.  $\text{Ci}(\infty) = 0$

**Fresnel Sine Integral  $S(x) = \sqrt{\frac{2}{\pi}} \int_0^x \sin u^2 du$**

---

36.18.  $S(x) = \sqrt{\frac{2}{\pi}} \left( \frac{x^3}{3 \cdot 1!} - \frac{x^7}{7 \cdot 3!} + \frac{x^{11}}{11 \cdot 5!} - \frac{x^{15}}{15 \cdot 7!} + \dots \right)$

36.19.  $S(x) \sim \frac{1}{2} - \frac{1}{\sqrt{2\pi}} \left\{ (\cos x^2) \left( \frac{1}{x} - \frac{1 \cdot 3}{2^2 x^5} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{2^4 x^9} - \dots \right) + (\sin x^2) \left( \frac{1}{2x^3} - \frac{1 \cdot 3 \cdot 5}{2^3 x^7} + \dots \right) \right\}$

36.20.  $S(-x) = -S(x), \quad S(0) = 0, \quad S(\infty) = \frac{1}{2}$

**Fresnel Cosine Integral  $C(x) = \sqrt{\frac{2}{\pi}} \int_0^x \cos u^2 du$**

---

36.21.  $C(x) = \sqrt{\frac{2}{\pi}} \left( \frac{x}{1!} - \frac{x^5}{5 \cdot 2!} + \frac{x^9}{9 \cdot 4!} - \frac{x^{13}}{13 \cdot 6!} + \dots \right)$

36.22.  $C(x) \sim \frac{1}{2} + \frac{1}{\sqrt{2\pi}} \left\{ (\sin x^2) \left( \frac{1}{x} - \frac{1 \cdot 3}{2^2 x^5} + \frac{1 \cdot 3 \cdot 5 \cdot 7}{2^4 x^9} - \dots \right) - (\cos x^2) \left( \frac{1}{2x^3} - \frac{1 \cdot 3 \cdot 5}{2^3 x^7} + \dots \right) \right\}$

36.23.  $C(-x) = -C(x), \quad C(0) = 0, \quad C(\infty) = \frac{1}{2}$

**Riemann Zeta Function  $\zeta(x) = \frac{1}{1^x} + \frac{1}{2^x} + \frac{1}{3^x} + \dots$**

---

36.24.  $\zeta(x) = \frac{1}{\Gamma(x)} \int_0^{\infty} \frac{u^{x-1}}{e^{u-1}} du, \quad x > 1$

36.25.  $\zeta(1-x) = 2^{1-x} \pi^{-x} \Gamma(x) \cos(\pi x/2) \zeta(x)$  (extension to other values)

36.26.  $\zeta(2k) = \frac{2^{2k-1} \pi^{2k} B_k}{(2k)!} \quad k = 1, 2, 3, \dots$

## Section X: Inequalities and Infinite Products

# 37 INEQUALITIES

### Triangle Inequality

$$37.1. |a_1| - |a_2| \leq |a_1 + a_2| \leq |a_1| + |a_2|$$

$$37.2. |a_1 + a_2 + \dots + a_n| \leq |a_1| + |a_2| + \dots + |a_n|$$

### Cauchy-Schwarz Inequality

$$37.3. (a_1 b_1 + a_2 b_2 + \dots + a_n b_n)^2 \leq (a_1^2 + a_2^2 + \dots + a_n^2)(b_1^2 + b_2^2 + \dots + b_n^2)$$

The equality holds if, and only if,  $a_1/b_1 = a_2/b_2 = \dots = a_n/b_n$ .

### Inequalities Involving Arithmetic, Geometric, and Harmonic Means

If  $A$ ,  $G$ , and  $H$  are the arithmetic, geometric, and harmonic means of the positive numbers  $a_1, a_2, \dots, a_n$ , then

$$37.4. H \leq G \leq A$$

where

$$37.5. A = \frac{a_1 + a_2 + \dots + a_n}{n}$$

$$37.6. G = \sqrt[n]{a_1 a_2 \dots a_n}$$

$$37.7. \frac{1}{H} = \frac{1}{n} \left( \frac{1}{a_1} + \frac{1}{a_2} + \dots + \frac{1}{a_n} \right)$$

The equality holds if, and only if,  $a_1 = a_2 = \dots = a_n$ .

### Holder's Inequality

$$37.8. |a_1 b_1 + a_2 b_2 + \dots + a_n b_n| \leq (|a_1|^p + |a_2|^p + \dots + |a_n|^p)^{1/p} (|b_1|^q + |b_2|^q + \dots + |b_n|^q)^{1/q}$$

where

$$37.9. \frac{1}{p} + \frac{1}{q} = 1 \quad p > 1, q > 1$$

The equality holds if, and only if,  $|a_1|^{p-1}/|b_1| = |a_2|^{p-1}/|b_2| = \dots = |a_n|^{p-1}/|b_n|$ . For  $p = q = 2$  it reduces to 37.3.

### Chebyshev's Inequality

---

If  $a_1 \geq a_2 \geq \dots \geq a_n$  and  $b_1 \geq b_2 \geq \dots \geq b_n$ , then

$$37.10. \quad \left( \frac{a_1 + a_2 + \dots + a_n}{n} \right) \left( \frac{b_1 + b_2 + \dots + b_n}{n} \right) \leq \frac{a_1 b_1 + a_2 b_2 + \dots + a_n b_n}{n}$$

or

$$37.11. \quad (a_1 + a_2 + \dots + a_n)(b_1 + b_2 + \dots + b_n) \leq n(a_1 b_1 + a_2 b_2 + \dots + a_n b_n)$$

### Minkowski's Inequality

---

If  $a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n$  are all positive and  $p > 1$ , then

$$37.12. \quad \{(a_1 + b_1)^p + (a_2 + b_2)^p + \dots + (a_n + b_n)^p\}^{1/p} \leq (a_1^p + a_2^p + \dots + a_n^p)^{1/p} + (b_1^p + b_2^p + \dots + b_n^p)^{1/p}$$

The equality holds if, and only if,  $a_1/b_1 = a_2/b_2 = \dots = a_n/b_n$ .

### Cauchy-Schwarz Inequality for Integrals

---

$$37.13. \quad \left[ \int_a^b f(x)g(x) dx \right]^2 \leq \left\{ \int_a^b [f(x)]^2 dx \right\} \left\{ \int_a^b [g(x)]^2 dx \right\}$$

The equality holds if, and only if,  $f(x)/g(x)$  is a constant.

### Hölder's Inequality for Integrals

---

$$37.14. \quad \int_a^b |f(x)g(x)| dx \leq \left\{ \int_a^b |f(x)|^p dx \right\}^{1/p} \left\{ \int_a^b |g(x)|^q dx \right\}^{1/q}$$

where  $1/p + 1/q = 1$ ,  $p > 1$ ,  $q > 1$ . If  $p = q = 2$ , this reduces to 37.13.

The equality holds if, and only if,  $|f(x)|^{p-1}/|g(x)|$  is a constant.

### Minkowski's Inequality for Integrals

---

If  $p > 1$ ,

$$37.15. \quad \left\{ \int_a^b |f(x) + g(x)|^p dx \right\}^{1/p} \leq \left\{ \int_a^b |f(x)|^p dx \right\}^{1/p} + \left\{ \int_a^b |g(x)|^p dx \right\}^{1/p}$$

The equality holds if, and only if,  $f(x)/g(x)$  is a constant.

# 38 INFINITE PRODUCTS

$$38.1. \quad \sin x = x \left(1 - \frac{x^2}{x^2}\right) \left(1 - \frac{x^2}{4\pi^2}\right) \left(1 - \frac{x^2}{9\pi^2}\right) \dots$$

$$38.2. \quad \cos x = \left(1 - \frac{4x^2}{\pi^2}\right) \left(1 - \frac{4x^2}{9\pi^2}\right) \left(1 - \frac{4x^2}{25\pi^2}\right) \dots$$

$$38.3. \quad \sinh x = x \left(1 + \frac{x^2}{\pi^2}\right) \left(1 + \frac{x^2}{4\pi^2}\right) \left(1 + \frac{x^2}{9\pi^2}\right) \dots$$

$$38.4. \quad \cosh x = \left(1 + \frac{4x^2}{\pi^2}\right) \left(1 + \frac{4x^2}{9\pi^2}\right) \left(1 + \frac{4x^2}{25\pi^2}\right) \dots$$

$$38.5. \quad \frac{1}{\Gamma(x)} = xe^{\gamma x} \left\{ \left(1 + \frac{x}{1}\right) e^{-x} \right\} \left\{ \left(1 + \frac{x}{2}\right) e^{-x/2} \right\} \left\{ \left(1 + \frac{x}{3}\right) e^{-x/3} \right\} \dots$$

See also 25.11.

$$38.6. \quad J_0(x) = \left(1 - \frac{x^2}{\lambda_1^2}\right) \left(1 - \frac{x^2}{\lambda_2^2}\right) \left(1 - \frac{x^2}{\lambda_3^2}\right) \dots$$

where  $\lambda_1, \lambda_2, \lambda_3, \dots$  are the positive roots of  $J_0(x) = 0$ .

$$38.7. \quad J_1(x) = x \left(1 - \frac{x^2}{\lambda_1^2}\right) \left(1 - \frac{x^2}{\lambda_2^2}\right) \left(1 - \frac{x^2}{\lambda_3^2}\right) \dots$$

where  $\lambda_1, \lambda_2, \lambda_3, \dots$  are the positive roots of  $J_1(x) = 0$ .

$$38.8. \quad \frac{\sin x}{x} = \cos \frac{x}{2} \cos \frac{x}{4} \cos \frac{x}{8} \cos \frac{x}{16} \dots$$

$$38.9. \quad \frac{\pi}{2} = \frac{2}{1} \cdot \frac{2}{3} \cdot \frac{4}{3} \cdot \frac{4}{5} \cdot \frac{6}{5} \cdot \frac{6}{7} \cdot \dots$$

This is called Wallis' product.

## Section XI: Probability and Statistics

# 39 DESCRIPTIVE STATISTICS

The numerical data  $x_1, x_2, \dots$  will either come from a random sample of a larger population or from the larger population itself. We distinguish these two cases using different notation as follows:

$n$  = number of items in a sample,  
 $N$  = number of items in the population,

$\bar{x}$  = (read:  $x$ -bar) = sample mean,  
 $s^2$  = sample variance,  
 $s$  = sample standard deviation,

$\mu$  (read: mu) = population mean,  
 $\sigma^2$  = population variance,  
 $\sigma$  = population standard deviation

Note that Greek letters are used with the population and are called *parameters*, whereas Latin letters are used with the samples and are called *statistics*. First we give formulas for the data coming from a sample. This is followed by formulas for the population.

### Grouped Data

Frequently, the sample data are collected into groups (grouped data). A group refers to a set of numbers all with the same value  $x_i$ , or a set (class) of numbers in a given interval with class value  $x_i$ . In such a case, we assume there are  $k$  groups with  $f_i$  denoting the number of elements in the group with value or class value  $x_i$ .

Thus, the total number of data items is

$$39.1. \quad n = \sum f_i$$

As usual,  $\Sigma$  will denote a summation over all the values of the index, unless otherwise specified.

Accordingly, some of the formulas will be designated as (a) or as (b), where (a) indicates ungrouped data and (b) indicates grouped data.

### Measures of Central Tendency

#### Mean (Arithmetic Mean)

The *arithmetic mean* or simply *mean* of a sample  $x_1, x_2, \dots, x_n$ , frequently called the “average value,” is the sum of the values divided by the number of values. That is:

$$39.2(a). \quad \text{Sample mean:} \quad \bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum x_i}{n}$$

$$39.2(b). \quad \text{Sample mean:} \quad \bar{x} = \frac{f_1 x_1 + f_2 x_2 + \dots + f_k x_k}{f_1 + f_2 + \dots + f_k} = \frac{\sum f_i x_i}{\sum f_i}$$

#### Median

Suppose that the data  $x_1, x_2, \dots, x_n$  are now sorted in increasing order. The *median* of the data, denoted by

$M$  or Median

is defined to be the “middle value.” That is:

$$39.3(a). \quad \text{Median} = \begin{cases} x_{k+1} & \text{when } n \text{ is odd and } n = 2k+1, \\ \frac{x_k + x_{k+1}}{2} & \text{when } n \text{ is even and } n = 2k. \end{cases}$$

The median of grouped data is obtained by first finding the *cumulative frequency* function  $F_s$ . Specifically, we define

$$F_s = f_1 + f_2 + \cdots + f_s$$

that is,  $F_s$  is the sum of the frequencies up to  $f_s$ . Then:

$$39.3(b.1). \quad \text{Median} = \begin{cases} x_{j+1} & \text{when } n = 2k+1 \text{ (odd) and } F_j < k+1 \leq F_{j+1} \\ \frac{x_j + x_{j+1}}{2} & \text{when } n = 2k \text{ (even), and } F_j = k. \end{cases}$$

Finding the median of data arranged in classes is more complicated. First one finds the median class  $m$ , the class with the median value, and then one linearly interpolates in the class using the formula

$$39.3(b.2). \quad \text{Median} = L_m + c \frac{(n/2) - F_{m-1}}{f_m}$$

where  $L_m$  denotes the lower class boundary of the median class and  $c$  denotes its class width (length of the class interval).

### Mode

The mode is the value or values which occur most often. Namely:

**39.4.** Mode  $x_m$  = numerical value that occurs the most number of times

The mode is not defined if every  $x_m$  occurs the same number of times, and when the mode is defined it may not be unique.

### Weighted and grand means

Suppose that each  $x_i$  is assigned a weight  $w_i \geq 0$ . Then:

$$39.5. \quad \text{Weighted Mean } \bar{x}_w = \frac{w_1 x_1 + w_2 x_2 + \cdots + w_k x_k}{w_1 + w_2 + \cdots + w_k} = \frac{\sum w_i x_i}{\sum w_i}$$

Note that 39.2(b.1) is a special case of 39.4 where the weight  $w_i$  of  $x_i$  is its frequency.

Suppose that there are  $k$  sample sets and that each sample set has  $n_i$  elements and a mean  $\bar{x}_i$ . Then the *grand mean*, denoted by  $\bar{\bar{x}}$ , is the “mean of the means” where each mean is weighted by the number of elements in its sample. Specifically:

$$39.6. \quad \text{Grand Mean } \bar{\bar{x}} = \frac{n_1 \bar{x}_1 + n_2 \bar{x}_2 + \cdots + n_k \bar{x}_k}{n_1 + n_2 + \cdots + n_k} = \frac{\sum n_i \bar{x}_i}{\sum n_i}$$

### Geometric and Harmonic Means

The *geometric mean* (G.M.) and *harmonic mean* (H.M.) are defined as follows:

$$39.7(a). \quad \text{G.M.} = \sqrt[n]{x_1 x_2 \cdots x_n}$$

$$39.7(b). \quad \text{G.M.} = \sqrt[n]{x_1^{f_1} x_2^{f_2} \cdots x_k^{f_k}}$$

**39.8(a).** H.M. =  $\frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}} = \frac{n}{\sum (1/x_i)}$

**39.8(b).** H.M. =  $\frac{n}{f_1/x_1 + f_2/x_2 + \dots + f_k/x_k} = \frac{n}{\sum (f_i/x_i)}$

### Relation Between Arithmetic, Geometric, and Harmonic Means

**39.9.** H.M.  $\leq$  G.M.  $\leq$   $\bar{x}$

The equality sign holds only when all the sample values are equal.

### Midrange

The *midrange* is the average of the smallest value  $x_1$  and the largest value  $x_n$ . That is:

**39.10.** midrange: mid =  $\frac{x_1 + x_n}{2}$

### Population Mean

The formula for the population mean  $\mu$  follows:

**39.11(a).** Population mean:  $\mu = \frac{x_1 + x_2 + \dots + x_N}{N} = \frac{\sum x_i}{N}$

**39.11(b).** Population mean:  $\mu = \frac{f_1 x_1 + f_2 x_2 + \dots + f_k x_k}{f_1 + f_2 + \dots + f_k} = \frac{\sum f_i x_i}{\sum f_i}$

(Recall that  $N$  denotes the number of elements in a population.)

Observe that the formula for the population mean  $\mu$  is the same as the formula for the sample mean  $\bar{x}$ . On the other hand, the formula for the population standard deviation  $\sigma$  is not the same as the formula for the sample standard deviation  $s$ . (This is the main reason we give separate formulas for  $\mu$  and  $\bar{x}$ .)

---

## Measures of Dispersion

### Sample Variance and Standard Deviation

Here the sample set has  $n$  elements with mean  $\bar{x}$ .

**39.12(a).** Sample variance:  $s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1} = \frac{\sum x_i^2 - (\sum x_i)^2/n}{n-1}$

**39.12(b).** Sample variance:  $s^2 = \frac{\sum f_i (x_i - \bar{x})^2}{(\sum f_i) - 1} = \frac{\sum f_i x_i^2 - (\sum f_i x_i)^2 / \sum f_i}{(\sum f_i) - 1}$

**39.13.** Sample standard deviation:  $s = \sqrt{\text{Variance}} = \sqrt{s^2}$

**EXAMPLE 39.1:** Consider the following frequency distribution:

$x_i$	1	2	3	4	5	6
$f_i$	8	14	7	12	3	1

Then  $n = \sum f_i = 45$  and  $\sum f_i x_i = 126$ . Hence, by 39.2(b),

$$\text{Mean } \bar{x} = \frac{\sum x_i f_i}{\sum f_i} = \frac{126}{45} = 2.8$$

Also,  $n - 1 = 44$  and  $\sum f_i x_i^2 = 430$ . Hence, by 39.12(b) and 39.13,

$$s^2 = \frac{430 - (126)^2/45}{44} \approx 1.75 \quad \text{and} \quad s = 1.32$$

We find the median M, first finding the cumulative frequencies:

$$F_1 = 8, \quad F_2 = 22, \quad F_3 = 29, \quad F_4 = 41, \quad F_5 = 44, \quad F_6 = 45 = n$$

Here  $n$  is odd, and  $(n + 1)/2 = 23$ . Hence,

$$\text{Median } M = 23\text{rd value} = 3$$

The value 2 occurs most often, hence

$$\text{Mode} = 2$$

### M.D. and R.M.S.

Here M.D. stands for *mean deviation* and R.M.S. stands for *root mean square*. As previously,  $\bar{x}$  is the mean of the data and, for grouped data,  $n = \sum f_i$ .

$$\mathbf{39.14(a).} \quad \text{M.D.} = \frac{1}{n} \left| x_i - \bar{x} \right|$$

$$\mathbf{39.14(b).} \quad \text{M.D.} = \frac{1}{n} \left| f_i x_i - \bar{x} \right|$$

$$\mathbf{39.15(a).} \quad \text{R.M.S.} = \sqrt{\frac{1}{n} (\sum x_i^2)}$$

$$\mathbf{39.15(b).} \quad \text{R.M.S.} = \sqrt{\frac{1}{n} (\sum f_i x_i^2)}$$

### Measures of Position (Quartiles and Percentiles)

Now we assume that the data  $x_1, x_2, \dots, x_n$  are arranged in increasing order.

**39.16.** Sample range:  $x_n - x_1$ .

There are three quartiles: the first or lower quartile, denoted by  $Q_1$  or  $Q_L$ ; the second quartile or median, denoted by  $Q_2$  or  $M$ ; and the third or upper quartile, denoted by  $Q_3$  or  $Q_U$ . These quartiles (which essentially divide the data into “quarters”) are defined as follows, where “half” means  $n/2$  when  $n$  is even and  $(n-1)/2$  when  $n$  is odd:

**39.17.**  $Q_L (= Q_1)$  = median of the first half of the values.

$M (= Q_2)$  = median of the values.

$Q_U (= Q_3)$  = median of the second half of the values.

**39.18.** Five-number summary:  $[L, Q_L, M, Q_U, H]$  where  $L = x_1$  (lowest value) and  $H = x_n$  (highest value).

**39.19.** Innerquartile range:  $Q_U - Q_L$

**39.20.** Semi-innerquartile range:  $Q = \frac{Q_U - Q_L}{2}$

The  $k$ th percentile, denoted by  $P_k$ , is the number for which  $k$  percent of the values are at most  $P_k$  and  $(100-k)$  percent of the values are greater than  $P_k$ . Specifically:

**39.21.**  $P_k$  = largest  $x_s$  such that  $F_s \leq k/100$ . Thus,  $Q_L$  = 25th percentile,  $M$  = 50th percentile,  $Q_U$  = 75th percentile.

### Higher-Order Statistics

**39.22.** The  $r$ th moment: (a)  $m_r = \frac{1}{n} \sum x_i^r$ , (b)  $m_r = \frac{1}{n} \sum f_i x_i^r$

**39.23.** The  $r$ th moment about the mean  $\bar{x}$ :

$$(a) \mu_r = \frac{1}{n} \sum (x_i - \bar{x})^r, \quad (b) \mu_r = \frac{1}{n} \sum (f_i x_i - \bar{x})^r$$

**39.24.** The  $r$ th absolute moment about mean  $\bar{x}$ :

$$(a) \mu_r = \frac{1}{n} \sum |x_i - \bar{x}|^r, \quad (b) \mu_r = \frac{1}{n} \sum |f_i x_i - \bar{x}|^r$$

**39.25.** The  $r$ th moment in standard  $z$  units about  $z = 0$ :

$$(a) \alpha_r = \frac{1}{n} \sum z_i^r, \quad (b) \alpha_r = \frac{1}{n} \sum f_i z_i^r \text{ where } z_i = \frac{x_i - \bar{x}}{\sigma}$$

### Measures of Skewness and Kurtosis

**39.26.** Coefficient of skewness:  $\gamma_1 = \frac{\mu_3}{\sigma^3} = \alpha_3$

**39.27.** Momental skewness:  $\frac{\mu_3}{2\sigma^3}$

**39.28.** Coefficient of kurtosis:  $\alpha_4 = \frac{\mu_4}{\sigma^4}$

**39.29.** Coefficient of excess (kurtosis):  $\alpha_4 - 3 = \frac{\mu_4}{\sigma^4} - 3$

**39.30.** Quartile coefficient of skewness:  $\frac{Q_U - 2\hat{x} + Q_L}{Q_U - Q_L} = \frac{Q_3 - 2Q_2 + Q_1}{Q_3 - Q_1}$

### Population Variance and Standard Deviation

Recall that  $N$  denotes the number of values in the population.

**39.31.** Population variance:  $\sigma^2 = \frac{\sum (x_i - \bar{x})^2}{N} = \frac{\sum x_i^2 - (\sum x_i)^2/N}{N}$

**39.32.** Population standard deviation:  $\sigma = \sqrt{\text{Variance}} = \sqrt{\sigma^2}$

---

### Bivariate Data

The following formulas apply to a list of pairs of numerical values:

$$(x_1, y_1), (x_2, y_2), (x_3, y_3), \dots, (x_n, y_n)$$

where the first values correspond to a variable  $x$  and the second to a variable  $y$ . The primary objective is to determine whether there is a mathematical relationship, such as a linear relationship, between the data.

The *scatterplot* of the data is simply a picture of the pairs of values as points in a coordinate plane.

### Correlation Coefficient

A numerical indicator of a linear relationship between variables  $x$  and  $y$  is the *sample correlation coefficient*  $r$  of  $x$  and  $y$ , defined as follows:

**39.33.** Sample correlation coefficient: 
$$r = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum(x_i - \bar{x})^2} \sqrt{\sum(y_i - \bar{y})^2}}$$

We assume that the denominator in Formula 39.33 is not zero. An alternative formula for computing  $r$  follows:

**39.34.** 
$$r = \frac{\sum x_i y_i - (\sum x_i)(\sum y_i)/n}{\sqrt{\sum x_i^2 - (\sum x_i)^2/n} \sqrt{\sum y_i^2 - (\sum y_i)^2/n}}$$

Properties of the correlation coefficient  $r$  follow:

- 39.35.** (1)  $-1 \leq r \leq 1$  or, equivalently,  $|r| \leq 1$ .  
 (2)  $r$  is positive or negative according as  $y$  tends to increase or decrease as  $x$  increases.  
 (3) The closer  $|r|$  is to 1, the stronger the linear relationship between  $x$  and  $y$ .

The *sample covariance* of  $x$  and  $y$  is denoted and defined as follows:

**39.36.** Sample covariance: 
$$s_{xy} = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

Using the sample covariance, Formula 39.33 can be written in the compact form:

**39.37.** 
$$r = \frac{s_{xy}}{s_x s_y}$$

where  $s_x$  and  $s_y$  are the sample standard deviations of  $x$  and  $y$ , respectively.


**EXAMPLE 39.2:** Consider the following data:

$x$	50	45	40	38	32	40	55
$y$	2.5	5.0	6.2	7.4	8.3	4.7	1.8

The scatterplot of the data appears in Fig. 39-1. The correlation coefficient  $r$  for the data may be obtained by first constructing the table in Fig. 39-2. Then, by Formula 39.34 with  $n = 7$ ,

$$r = \frac{1431.8 - (300)(35.9)/7}{\sqrt{13,218 - (300)^2/7} \sqrt{218.67 + (35.9)^2/7}} \approx -0.9562$$

Here  $r$  is close to  $-1$ , and the scatterplot in Fig. 39-1 does indicate a strong negative linear relationship between  $x$  and  $y$ .



$x_i$	$y_i$	$x_i^2$	$y_i^2$	$x_i y_i$
50	2.5	2,500	6.25	125.0
45	5.0	2,025	25.00	225.0
40	6.2	1,600	38.44	248.0
38	7.4	1,444	54.76	281.2
32	8.3	1,024	68.89	265.6
40	4.7	1,600	22.09	188.0
55	1.8	3,025	3.24	99.0
Sums				
300	35.9	13,218	218.67	1431.8

Fig. 39-2

### Regression Line

Consider a given set of  $n$  data points  $P_i(x_i, y_i)$ . Any (nonvertical) line  $L$  may be defined by an equation of the form

$$y = a + bx$$

Let  $y_i^*$  denote the  $y$  value of the point on  $L$  corresponding to  $x_i$ ; that is, let  $y_i^* = a + bx_i$ . Now let

$$d_i = y_i - y_i^* = y_i - (a + bx_i)$$

that is,  $d_i$  is the vertical (directed) distance between the point  $P_i$  and the line  $L$ . The *squares error* between the line  $L$  and the data points is defined by

$$39.38. \quad \sum d_i^2 = d_1^2 + d_2^2 + \cdots + d_n^2$$

The *least-squares line* or the *line of best fit* or the *regression line* of  $y$  on  $x$  is, by definition, the line  $L$  whose squares error is as small as possible. It can be shown that such a line  $L$  exists and is unique.

The constants  $a$  and  $b$  in the equation  $y = a + bx$  of the line  $L$  of best fit can be obtained from the following two *normal equations*, where  $a$  and  $b$  are the unknowns and  $n$  is the number of points:

$$39.39. \quad \begin{cases} na + (\sum x_i)b = \sum y_i \\ (\sum x_i)a + (\sum x_i^2)b = \sum x_i y_i \end{cases}$$

The solution of the above normal equations follows:

$$39.40. \quad b = \frac{n \sum x_i y_i - (\sum x_i)(\sum y_i)}{n \sum x_i^2 - (\sum x_i)^2} = \frac{rs_y}{s_x}; \quad a = \frac{\sum y_i}{n} - b \frac{\sum x_i}{n} = \bar{y} - b\bar{x}$$

The second equation tells us that the point  $(\bar{x}, \bar{y})$  lies on  $L$ , and the first equation tells us that the point  $(\bar{x} + s_x, \bar{y} + rs_y)$  also lies on  $L$ .

**EXAMPLE 39.3:** Suppose we want the line  $L$  of best fit for the data in Example 39.2. Using the table in Fig. 39-2 and  $n = 7$ , we obtain the normal equations

$$\begin{aligned} 7a + 300b &= 35.9 \\ 300a + 13,218b &= 1431 \end{aligned}$$

Substitution in 39.40 yields

$$\begin{aligned} b &= \frac{7(1431.8) - (300)(35.9)}{7(13,218) - (300)^2} = -0.2959 \\ a &= \frac{35.9}{7} - (-0.2959) \frac{300}{7} = 17.8100 \end{aligned}$$

Thus, the line  $L$  of best fit is

$$y = 17.8100 - 0.2959x$$

The graph of  $L$  appears in Fig. 39-3.




Fig. 39-3

### Curve Fitting

Suppose that  $n$  data points  $P_i(x_i, y_i)$  are given, and that the data (using the scatterplot or the correlation coefficient  $r$ ) do not indicate a linear relationship between the variables  $x$  and  $y$ , but do indicate that some other standard (well-known) type of curve  $y = f(x)$  approximates the data. Then the particular curve  $C$  that one uses to approximate that data, called the *best-fitting* or *least-squares* curve, is the curve in the collection which minimizes the squares error sum

$$\sum d_i^2 = d_1^2 + d_2^2 + \cdots + d_n^2$$

where  $d_i = y_i - f(x_i)$ . Three such types of curve are discussed as follows.

*Polynomial function of degree  $m$ :*  $y = a_0 + a_1x + a_2x^2 + \cdots + a_mx^m$

The coefficients  $a_0, a_1, a_2, \dots, a_m$  of the best-fitting polynomial can be obtained by solving the following system of  $m+1$  normal equations:

$$39.41. \quad \begin{aligned} na_0 + a_1 \sum x_i + a_2 \sum x_i^2 + \cdots + a_m \sum x_i^m &= \sum y_i \\ a_0 \sum x_i + a_1 \sum x_i^2 + a_2 \sum x_i^3 + \cdots + a_m \sum x_i^{m+1} &= \sum x_i y_i \\ a_0 \sum x_i^m + a_1 \sum x_i^{m+1} + a_2 \sum x_i^{m+2} + \cdots + a_m \sum x_i^{2m} &= \sum x_i^m y_i \end{aligned}$$

*Exponential curve:*  $y = ab^x$  or  $\log y = \log a + (\log b)x$

The exponential curve is used if the scatterplot of  $\log y$  versus  $x$  indicates a linear relationship. Then  $\log a$  and  $\log b$  are obtained from transformed data points. Namely, the best-fit line  $L$  for data points  $P'(x_i, \log y_i)$  is

$$39.42. \quad \begin{cases} na' + (\sum x_i)b' = \sum(\log y_i) \\ (\sum x_i)a' + (\sum x_i^2)b' = \sum(x_i \log y_i) \end{cases}$$

Then  $a = \text{antilog } a'$ ,  $b = \text{antilog } b'$ .

**EXAMPLE 39.4:** Consider the following data which indicates exponential growth:

$x$	1	2	3	4	5	6
$y$	6	18	55	160	485	1460

Thus, we seek the least-squares line  $L$  for the following data:

$x$	1	2	3	4	5	6
$\log y$	0.7782	1.2553	1.7404	2.2041	2.6857	3.1644

Using the normal equation 39.42 for  $L$ , we get

$$a' = 0.3028, \quad b' = 0.4767$$

The antiderivatives of  $a'$  and  $b'$  yield, approximately,

$$a = 2.0, \quad b = 3.0$$

Hence,  $y = 2(3^x)$  is the required exponential curve  $C$ . The data points and  $C$  are depicted in Fig. 39-4.




Fig. 39-4

*Power function:*  $y = ax^b$  or  $\log y = \log a + b \log x$

The power curve is used if the scatterplot of  $\log y$  versus  $\log x$  indicates a linear relationship. The  $\log a$  and  $b$  are obtained from transformed data points. Namely, the best-fit line  $L$  for transformed data points  $P'(\log x_i, \log y_i)$  is

$$39.43. \quad \begin{cases} na' + \sum(\log x_i)b = \sum(\log y_i) \\ \sum(\log x_i)a' + \sum(\log x_i)^2b = \sum(\log x_i \log y_i) \end{cases}$$

Then  $a = \text{antilog } a'$ .

# 40 PROBABILITY

## Sample Spaces and Events

Let  $S$  be a sample space which consists of the possible outcomes of an experiment where the events are subsets of  $S$ . The sample space  $S$  itself is called the *certain event*, and the null set  $\emptyset$  is called the *impossible event*.

It would be convenient if all subsets of  $S$  could be events. Unfortunately, this may lead to contradictions when a probability function is defined on the events. Thus, the events are defined to be a limited collection  $C$  of subsets of  $S$  as follows.

**DEFINITION 40.1:** The class  $C$  of events of a sample space  $S$  form a  $\sigma$ -field. That is,  $C$  has the following three properties:

- (i)  $S \in C$ .
- (ii) If  $A_1, A_2, \dots$  belong to  $C$ , then their union  $A_1 \cup A_2 \cup A_3 \cup \dots$  belongs to  $C$ .
- (iii) If  $A \in C$ , then its complement  $A^c \in C$ .

Although the above definition does not mention intersections, DeMorgan's law (40.3) tells us that the complement of a union is the intersection of the complements. Thus, the events form a collection that is closed under unions, intersections, and complements of denumerable sequences.

If  $S$  is finite, then the class of all subsets of  $S$  form a  $\sigma$ -field. However, if  $S$  is nondenumerable, then only certain subsets of  $S$  can be the events. In fact, if  $B$  is the collection of all open intervals on the real line  $\mathbf{R}$ , then the smallest  $\sigma$ -field containing  $B$  is the collection of Borel sets in  $\mathbf{R}$ .

If Condition (ii) in Definition 40.1 of a  $\sigma$ -field is replaced by finite unions, then the class of subsets of  $S$  is called a *field*. Thus a  $\sigma$ -field is a field, but not visa versa.

First, for completeness, we list basic properties of the set operations of union, intersection, and complement.

**40.1.** Sets satisfy the properties in Table 40-1.

**TABLE 40-1 Laws of the Algebra of Sets**

Idempotent laws:	(1a) $A \cup A = A$	(1b) $A \cap A = A$
Associative laws:	(2a) $(A \cup B) \cup C = A \cup (B \cup C)$	(2b) $(A \cap B) \cap C = A \cap (B \cap C)$
Commutative laws:	(3a) $A \cup B = B \cup A$	(3b) $A \cap B = B \cap A$
Distributive laws:	(4a) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$	(4b) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
Identity laws:	(5a) $A \cup \emptyset = A$	(5b) $A \cap U = A$
	(6a) $A \cup U = U$	(6b) $A \cap \emptyset = \emptyset$
Involution law:	(7) $(A^c)^c = A$	
Complement laws:	(8a) $A \cup A^c = U$	(8b) $A \cap A^c = \emptyset$
	(9a) $U^c = \emptyset$	(9b) $\emptyset^c = U$
DeMorgan's laws:	(10a) $(A \cup B)^c = A^c \cap B^c$	(10b) $(A \cap B)^c = A^c \cup B^c$

- 40.2.** The following are equivalent: (i)  $A \subseteq B$ , (ii)  $A \cap B = A$ , (iii)  $A \cup B = B$ .

Recall that the union and intersection of any collection of sets is defined as follows:

$$\bigcup_j A_j = \{x \mid \text{there exists } j \text{ such that } x \in A_j\} \quad \text{and} \quad \bigcap_j A_j = \{x \mid \text{for every } j \text{ we have } x \in A_j\}$$

- 40.3.** (Generalized DeMorgan's Law) (10a)'  $(\bigcup_j A_j)^c = \bigcap_j A_j^c$ ; (10b)'  $(\bigcap_j A_j)^c = \bigcup_j A_j^c$

## Probability Spaces and Probability Functions

---

**DEFINITION 40.2:** Let  $P$  be a real-valued function defined on the class  $C$  of events of a sample space  $S$ . Then  $P$  is called a *probability function*, and  $P(A)$  is called the *probability* of an event  $A$ , when the following axioms hold:

**Axiom [P<sub>1</sub>]** For every event  $A$ ,  $P(A) \geq 0$ .

**Axiom [P<sub>2</sub>]** For the certain event  $S$ ,  $P(S) = 1$ .

**Axiom [P<sub>3</sub>]** For any sequence of mutually exclusive (disjoint) events  $A_1, A_2, \dots$ ,

$$P(A_1 \cup A_2 \cup \dots) = P(A_1) + P(A_2) + \dots$$

The triple  $(S, C, P)$ , or simply  $S$  when  $C$  and  $P$  are understood, is called a *probability space*.

**Axiom [P<sub>3'</sub>]** implies an analogous axiom for any finite number of sets. That is:

**Axiom [P<sub>3''</sub>]** For any finite collection of mutually exclusive events  $A_1, A_2, \dots, A_n$ ,

$$P(A_1 \cup A_2 \cup \dots \cup A_n) = P(A_1) + P(A_2) + \dots + P(A_n)$$

In particular, for two disjoint events  $A$  and  $B$ , we have  $P(A \cup B) = P(A) + P(B)$ .

The following properties follow directly from the above axioms.

- 40.4.** (Complement rule)  $P(A^c) = 1 - P(A)$ . Thus,  $P(\emptyset) = 0$ .

- 40.5.** (Difference Rule)  $P(A \setminus B) = P(A) - P(A \cap B)$ .

- 40.6.** (Addition Rule)  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ .

- 40.7.** For  $n \geq 2$ ,  $P\left(\bigcup_{j=1}^n A_j\right) \leq \sum_{j=1}^n P(A_j)$

- 40.8.** (Monotonicity Rule) If  $A \subseteq B$ , then  $P(A) \leq P(B)$ .

## Limits of Sequences of Events

---

- 40.9.** (Continuity) Suppose  $A_1, A_2, \dots$  form a monotonic increasing (decreasing) sequence of events; that is,  $A_j \subseteq A_{j+1}$  ( $A_j \supseteq A_{j+1}$ ). Let  $A = \bigcup_j A_j$  ( $A = \bigcap_j A_j$ ). Then  $\lim P(A_n)$  exists and

$$\lim P(A_n) = P(A)$$

For any sequence of events  $A_1, A_2, \dots$ , we define

$$\liminf A_n = \bigcup_{k=1}^{+\infty} \bigcap_{j=k}^{+\infty} A_j \quad \text{and} \quad \limsup A_n = \bigcap_{k=1}^{+\infty} \bigcup_{j=k}^{+\infty} A_j$$

If  $\liminf A_n = \limsup A_n$ , then we call this set  $\lim A_n$ . Note  $\lim A_n$  exists when the sequence is monotonic.

- 40.10.** For any sequence  $A_j$  of events in a probability space,

$$P(\liminf A_n) \leq \liminf P(A_n) \leq \limsup P(A_n) \leq P(\limsup A_n)$$

Thus, if  $\lim A_n$  exists, then  $P(\lim A_n) = \lim P(A_n)$ .

- 40.11.** For any sequence  $A_j$  of events in a probability space,  $P(\cup_j A_j) \leq \sum_j P(A_j)$ .

- 40.12.** (Borel-Cantelli Lemma) Suppose  $A_j$  is any sequence of events in a probability space. Furthermore, suppose  $\sum_{n=1}^{+\infty} P(A_n) < +\infty$ . Then  $P(\limsup A_n) = 0$ .

- 40.13.** (Extension Theorem) Let  $F$  be a field of subsets of  $S$ . Let  $P$  be a function on  $F$  satisfying Axioms  $P_1$ ,  $P_2$ , and  $P_3'$ . Then there exists a unique probability function  $P^*$  on the smallest  $\sigma$ -field containing  $F$  such that  $P^*$  is equal to  $P$  on  $F$ .

## Conditional Probability

---

**DEFINITION 40.3:** Let  $E$  be an event with  $P(E) > 0$ . The conditional probability of an event  $A$  given  $E$  is denoted and defined as follows:

$$P(A|E) = \frac{P(A \cap E)}{P(E)}$$

- 40.14.** (Multiplication Theorem for Conditional Probability)  $P(A \cap B) = P(A)P(B|A)$ . This theorem can be generalized as follows:

- 40.15.**  $P(A_1 \cap \dots \cap A_n) = P(A_1)P(A_2|A_1)P(A_3|A_1 \cap A_2) \dots P(A_n|A_1 \cap \dots \cap A_{n-1})$

**EXAMPLE 40.1:** A lot contains 12 items of which 4 are defective. Three items are drawn at random from the lot one after the other. Find the probability that all three are nondefective.

The probability that the first item is nondefective is  $8/12$ . Assuming the first item is nondefective, the probability that the second item is nondefective is  $7/11$ . Assuming the first and second items are nondefective, the probability that the third item is nondefective is  $6/10$ . Thus,

$$p = \frac{8}{12} \cdot \frac{7}{11} \cdot \frac{6}{10} = \frac{14}{55}$$

## Stochastic Processes and Probability Tree Diagrams

---

A (finite) stochastic process is a finite sequence of experiments where each experiment has a finite number of outcomes with given probabilities. A convenient way of describing such a process is by means of a probability tree diagram, illustrated below, where the multiplication theorem (40.14) is used to compute the probability of an event which is represented by a given path of the tree.

**EXAMPLE 40.2:** Let  $X, Y, Z$  be three coins in a box where  $X$  is a fair coin,  $Y$  is two-headed, and  $Z$  is weighted so the probability of heads is  $1/3$ . A coin is selected at random and is tossed. (a) Find  $P(H)$ , the probability that heads appears. (b) Find  $P(X|H)$ , the probability that the fair coin  $X$  was picked if heads appears.

The probability tree diagram corresponding to the two-step stochastic process appears in Fig. 40-1a.

- (a) Heads appears on three of the paths (from left to right); hence,

$$P(H) = \frac{1}{3} \cdot \frac{1}{2} + \frac{1}{3} \cdot 1 + \frac{1}{3} \cdot \frac{1}{3} = \frac{11}{18}$$

- (b) X and heads H appear only along the top path; hence

$$P(X \cap H) = \frac{1}{3} \cdot \frac{1}{2} = \frac{1}{6} \text{ and so } P(X|H) = \frac{P(X \cap H)}{P(H)} = \frac{1/6}{11/18} = \frac{3}{11}$$




Fig. 40-1

### Law of Total Probability and Bayes' Theorem

Here we assume E is an event in a sample space S, and  $A_1, A_2, \dots, A_n$  are mutually disjoint events whose union is S; that is, the events  $A_1, A_2, \dots, A_n$  form a partition of S.

**40.16.** (Law of Total Probability)  $P(E) = P(A_1)P(E|A_1) + P(A_2)P(E|A_2) + \dots + P(A_n)P(E|A_n)$

**40.17.** (Bayes' Formula) For  $k = 1, 2, \dots, n$ ,

$$P(A_k|E) = \frac{P(A_k)P(E|A_k)}{P(E)} = \frac{P(A_k)P(E|A_k)}{P(A_1)P(E|A_1) + P(A_2)P(E|A_2) + \dots + P(A_n)P(E|A_n)}$$

**EXAMPLE 40.3:** Three machines, A, B, C, produce, respectively, 50%, 30%, and 20% of the total number of items in a factory. The percentages of defective output of these machines are, respectively, 3%, 4%, and 5%. An item is randomly selected.

(a) Find  $P(D)$ , the probability the item is defective.

(b) If the item is defective, find the probability it came from machine: (i) A, (ii) B, (iii) C.

(a) By **40.16** (Total Probability Law),

$$\begin{aligned} P(D) &= P(A)P(D|A) + P(B)P(D|B) + P(C)P(D|C) \\ &= (0.50)(0.03) + (0.30)(0.04) + (0.20)(0.05) = 3.7\% \end{aligned}$$

(b) By **40.17** (Bayes' rule), (i)  $P(A|D) = \frac{P(A)P(D|A)}{P(D)} = \frac{(0.50)(0.03)}{0.037} = 40.5\%$ . Similarly,

$$(ii) P(B|D) = \frac{P(B)P(D|B)}{P(D)} = 32.5\%; (iii) P(C|D) = \frac{P(C)P(D|C)}{P(D)} = 27.0\%$$

Alternately, we may consider this problem as a two-step stochastic process with a probability tree diagram, as in Fig. 40-1(b). We find  $P(D)$  by adding the three probability paths to D:

$$(0.50)(0.03) + (0.30)(0.04) + (0.20)(0.05) = 3.7\%$$

We find  $P(A|D)$  by dividing the top path to A and D by the sum of the three paths to D.

$$(0.50)(0.03)/0.037 = 40.5\%$$

Similarly, we find  $P(B|D) = 32.5\%$  and  $P(C|D) = 27.0\%$ .

## Independent Events

---

**DEFINITION 40.4:** Events A and B are independent if  $P(A \cap B) = P(A)P(B)$ .

**40.18.** The following are equivalent:

- (i)  $P(A \cap B) = P(A)P(B)$ , (ii)  $P(A|B) = P(A)$ , (iii)  $P(B|A) = P(B)$ .

That is, events A and B are independent if the occurrence of one of them does not influence the occurrence of the other.

**EXAMPLE 40.4:** Consider the following events for a family with children where we assume the sample space S is an equiprobable space:

$$E = \{\text{children of both sexes}\}, \quad F = \{\text{at most one boy}\}$$

- (a) Show that E and F are independent events if a family has three children.
- (b) Show that E and F are dependent events if a family has two children.
- (a) Here  $S = \{\text{bbb, bbg, bgb, bgg, gbb, gbg, ggb, ggg}\}$ . So:

$$E = \{\text{bbg, bgb, bgg, gbb, gbg, ggb}\}, P(E) = 6/8 = 3/4,$$

$$F = \{\text{bgg, gbg, ggb, ggg}\}, P(F) = 4/8 = 1/2$$

$$E \cap F = \{\text{bgg, gbg, ggb}\}, P(E \cap F) = 3/8$$

Therefore,  $P(E)P(F) = (3/4)(1/2) = 3/8 = P(E \cap F)$ . Hence, E and F are independent.

- (b) Here  $S = \{\text{bb, bg, gb, gg}\}$ . So:

$$E = \{\text{bg, gb}\}, P(E) = 2/4 = 1/2,$$

$$F = \{\text{bg, gb, gg}\}, P(F) = 3/4$$

$$E \cap F = \{\text{bg, gb}\}, P(E \cap F) = 2/4 = 1/2$$

Therefore,  $P(E)P(F) = (1/2)(3/4) = 3/8 \neq P(E \cap F)$ . Hence, E and F are dependent.

**DEFINITION 40.5:** For  $n > 2$ , the events  $A_1, A_2, \dots, A_n$  are independent if any proper subset of them is independent and

$$P(A_1 \cap A_2 \cap \dots \cap A_n) = P(A_1)P(A_2) \dots P(A_n)$$

Observe that induction is used in this definition.

**DEFINITION 40.6:** A collection  $\{A_j \mid j \in J\}$  of events is independent if, for any  $n > 0$ , the sets  $A_{j_1}, A_{j_2}, \dots, A_{j_n}$  are independent.

The concept of independent repeated trials, when S is a finite set, is formalized as follows.

**DEFINITION 40.7:** Let  $S$  be a finite probability space. The probability space of  $n$  independent trials or repeated trials, denoted by  $S_n$ , consists of ordered  $n$ -tuples  $(s_1, s_2, \dots, s_n)$  of elements of  $S$  with the probability of an  $n$ -tuple defined by

$$P((s_1, s_2, \dots, s_n)) = P(s_1)P(s_2) \dots P(s_n)$$

**EXAMPLE 40.5:** Suppose whenever horses  $a, b, c$  race together, their respective probabilities of winning are 20%, 30%, and 50%. That is,  $S = \{a, b, c\}$  with  $P(a) = 0.2$ ,  $P(b) = 0.3$ , and  $P(c) = 0.5$ .

They race three times. Find the probability that

- (a) the same horse wins all three times
  - (b) each horse wins once
- (a) Writing  $xyz$  for  $(x, y, z)$ , we seek the probability of the event  $A = \{aaa, bbb, ccc\}$ . Here,

$$P(aaa) = (0.2)^3 = 0.008, P(bbb) = (0.3)^3 = 0.027, P(ccc) = (0.5)^3 = 0.125$$

Thus,  $P(A) = 0.008 + 0.027 + 0.125 = 0.160$ .

- (b) We seek the probability of the event  $B = \{abc, acb, bac, bca, cab, cba\}$ . Each element in  $B$  has the same probability  $(0.2)(0.3)(0.5) = 0.03$ . Thus,  $P(B) = 6(0.03) = 0.18$ .

# 41 RANDOM VARIABLES

Consider a probability space  $(S, C, P)$ .

**DEFINITION 41.1.** A random variable  $X$  on the sample space  $S$  is a function from  $S$  into the set  $\mathbf{R}$  of real numbers such that the preimage of every interval of  $\mathbf{R}$  is an event of  $S$ .

If  $S$  is a discrete sample space in which every subset of  $S$  is an event, then every real-valued function on  $S$  is a random variable. On the other hand, if  $S$  is uncountable, then certain real-valued functions on  $S$  may not be random variables.

Let  $X$  be a random variable on  $S$ , where we let  $R_X$  denote the range of  $X$ ; that is,

$$R_X = \{x \mid \text{there exists } s \in S \text{ for which } X(s) = x\}$$

There are two cases that we treat separately. (i)  $X$  is a discrete random variable; that is,  $R_X$  is finite or countable. (ii)  $X$  is a continuous random variable; that is,  $R_X$  is a continuum of numbers such as an interval or a union of intervals.

Let  $X$  and  $Y$  be random variables on the same sample space  $S$ . Then, as usual,  $X + Y$ ,  $X + k$ ,  $kX$ , and  $XY$  (where  $k$  is a real number) are the functions on  $S$  defined as follows (where  $s$  is any point in  $S$ ):

$$\begin{aligned} (X + Y)(s) &= X(s) + Y(s), & (kX)(s) &= kX(s), \\ (X + k)(s) &= X(s) + k, & (XY)(s) &= X(s)Y(s). \end{aligned}$$

More generally, for any polynomial, exponential, or continuous function  $h(t)$ , we define  $h(X)$  to be the function on  $S$  defined by

$$[h(X)](s) = h[X(s)]$$

One can show that these are also random variables on  $S$ .

The following short notation is used:

$P(X = x_i)$	denotes the probability that $X = x_i$ .
$P(a \leq X \leq b)$	denotes the probability that $X$ lies in the closed interval $[a, b]$ .
$\mu_X$ or $E(X)$ or simply $\mu$	denotes the mean or expectation of $X$ .
$\sigma_{X^2}$ or $\text{Var}(X)$ or simply $\sigma^2$	denotes the variance of $X$ .
$\sigma_X$ or simply $\sigma$	denotes the standard deviation of $X$ .

Sometimes we let  $Y$  be a random variable such that  $Y = g(X)$ , that is, where  $Y$  is some function of  $X$ .

## Discrete Random Variables

Here  $X$  is a random variable with only a finite or countable number of values, say

$R_X = \{x_1, x_2, x_3, \dots\}$  where, say,  $x_1 < x_2 < x_3 < \dots$ . Then  $X$  induces a function  $f(x)$  on  $R_X$  as follows:

$$f(x_i) = P(X = x_i) = P(\{s \in S \mid X(s) = x_i\})$$

The function  $f(x)$  has the following properties:

$$(i) f(x_i) \geq 0 \quad \text{and} \quad (ii) \sum_i f(x_i) = 1$$

Thus,  $f$  defines a probability function on the range  $R_X$  of  $X$ . The pair  $(x_i, f(x_i))$ , usually given by a table, is called the *probability distribution* or *probability mass function* of  $X$ .

## Mean

---

**41.1.**  $\mu_x = E(X) = \sum x_i f(x_i)$

Here,  $Y = g(X)$ .

**41.2.**  $\mu_y = E(Y) = \sum g(x_i) f(x_i)$

Here,  $Y = g(X)$ .

## Variance and Standard Deviation

---

**41.3.**  $\sigma_x^2 = \text{Var}(X) = \sum (x_i - \mu)^2 f(x_i) = E((X - \mu)^2)$

Alternately,  $\text{Var}(X) = \sigma^2$  may be obtained as follows:

**41.4.**  $\text{Var}(X) = \sum x_i^2 f(x_i) - \mu^2 = E(X^2) - \mu^2$

**41.5.**  $\sigma_x = \sqrt{\text{Var}(X)} = \sqrt{E(X^2) - \mu^2}$

**REMARK:** Both the variance  $\text{Var}(X) = \sigma^2$  and the standard deviation  $\sigma$  measure the weighted spread of the values  $x_i$  about the mean  $\mu$ ; however, the standard deviation has the same units as  $\mu$ .

**EXAMPLE 41.1:** Suppose X has the following probability distribution:

x	2	4	6	8
f(x)	0.1	0.2	0.3	0.4

Then:

$$\mu = E(X) = \sum x_i f(x_i) = 2(0.1) + 4(0.2) + 6(0.3) + 8(0.4) = 6$$

$$E(X^2) = \sum x_i^2 f(x_i) = 2^2(0.1) + 4^2(0.2) + 6^2(0.3) + 8^2(0.4) = 40$$

$$\sigma^2 = \text{Var}(X) = E(X^2) - \mu^2 = 40 - 36 = 4$$

$$\sigma = \sqrt{\text{Var}(X)} = \sqrt{4} = 2$$

## Continuous Random Variable

---

Here X is a random variable with a continuum number of values. Then X determines a function  $f(x)$ , called the *density function* of X, such that

$$(i) f(x) \geq 0 \quad \text{and} \quad (ii) \int_{-\infty}^{\infty} f(x) dx = \int_R f(x) dx = 1$$

Furthermore,

$$P(a \leq X \leq b) = \int_a^b f(x) dx$$

## Mean

---

**41.6.**  $\mu_x = E(X) = \int_{-\infty}^{\infty} x f(x) dx$

Here,  $Y = g(X)$ .

**41.7.**  $\mu_y = E(Y) = \int_{-\infty}^{\infty} g(x) f(x) dx$

Here,  $Y = g(X)$ .

## Variance and Standard Deviation

---

**41.8.**  $\sigma_x^2 = \text{Var}(X) = \int_{-\infty}^{\infty} (x - \mu)^2 f(x) dx = E((X - \mu)^2)$

Alternately,  $\text{Var}(X) = \sigma^2$  may be obtained as follows:

**41.9.**  $\text{Var}(X) = \int_{-\infty}^{\infty} x^2 f(x) dx - \mu^2 = E(X^2) - \mu^2$

**41.10.**  $\sigma_x = \sqrt{\text{Var}(X)} = \sqrt{E(X^2) - \mu^2}$

**EXAMPLE 41.2:** Let X be the continuous random variable with the following density function:

$$f(x) = \begin{cases} (1/2)x & \text{if } 0 \leq x \leq 2 \\ 0 & \text{elsewhere} \end{cases}$$

Then:

$$E(X) = \int_{-\infty}^{\infty} x f(x) dx = \int_0^2 \frac{1}{2} x^2 dx = \left[ \frac{x^3}{6} \right]_0^2 = \frac{4}{3}$$

$$E(X^2) = \int_{-\infty}^{\infty} x^2 f(x) dx = \int_0^2 \frac{1}{2} x^3 dx = \left[ \frac{x^4}{8} \right]_0^2 = 2$$

$$\sigma^2 = \text{Var}(X) = E(X^2) - \mu^2 = 2 - \frac{16}{9} = \frac{2}{9}$$

$$\sigma = \sqrt{\text{Var}(X)} = \sqrt{\frac{2}{9}} = \frac{1}{3}\sqrt{2}$$

## Cumulative Distribution Function

---

The *cumulative distribution function* F(x) of a random variable X is the function  $F: \mathbf{R} \rightarrow \mathbf{R}$  defined by

**41.11.**  $F(a) = P(X \leq a)$

The function F is well-defined since the inverse of the interval  $(-\infty, a]$  is an event.

The function F(x) has the following properties:

**41.12.**  $F(a) \leq F(b)$  whenever  $a \leq b$ .

**41.13.**  $\lim_{x \rightarrow -\infty} F(x) = 0$  and  $\lim_{x \rightarrow +\infty} F(x) = 1$

That is, F(x) is monotonic, and the limit of F to the left is 0 and to the right is 1.

If X is the discrete random variable with distribution f(x), then F(x) is the following step function:

**41.14.**  $F(x) = \sum_{x_i \leq x} f(x_i)$

If X is a continuous random variable, then the density function f(x) of X can be obtained from the cumulative distribution function F(x) by differentiation. That is,

**41.15.**  $f(x) = \frac{d}{dx} F(x) = F'(x)$

Accordingly, for a continuous random variable X,

**41.16.**  $F(x) = \int_{-\infty}^x f(t) dt$

## Binomial Distribution

Consider an experiment with two outcomes, one called success (S) and the other called failure (F). A fixed number of independent repeated trials of such an experiment is called a binomial experiment. The notation

$$B(n, p)$$

denotes a binomial experiment with  $n$  independent trials and probability  $p$  of success. [We let  $q = 1 - p$  denote the probability of failure.]

**41.17.**  $P(k) = \binom{n}{k} p^k q^{n-k}$  is the probability of  $k$  successes in  $B(n, p)$  where  $\binom{n}{k}$  is the binomial coefficient.

**41.18.** The probability of at least one success in  $B(n, p)$  is  $1 - q^n$ .

The number  $X$  of  $k$  successes in  $B(n, p)$  is a random variable with the following distribution called the binomial distribution:

$k$	0	1	2	...	$n - 1$	$n$
$P(k)$	$q^n$	$\binom{n}{1} p q^{n-1}$	$\binom{n}{2} p^2 q^{n-2}$	...	$\binom{n}{n-1} p^{n-1} q$	$p^n$

**41.19.** Mean  $B(n, p) = \mu = np$ .

**41.20.** Variance  $B(n, p) = \sigma^2 = npq$ , and standard deviation of  $B(n, p) = \sigma = \sqrt{npq}$

### EXAMPLE 41.3:

(a) Jane hits a target with probability  $p = 1/3$ . So  $q = 2/3$ . She fires  $n = 6$  times. The probability of hitting the target exactly 2 times follows:

$$P(2) = \binom{6}{2} (1/3)^2 (2/3)^4 = 15(1/9)(16/81) = 80/243 = 0.329$$

(b) John hits a target with probability  $p = 1/4$ . So  $q = 3/4$ . He fires  $n = 100$  times. The expected number  $\mu$  of times he will hit the target and the standard deviation  $\sigma$  follows:

$$\mu = np = 100(1/4) = 25, \sigma^2 = npq = 100(1/4)(3/4) = 75/4 \text{ so } \sigma = 2.5$$

## Normal Distribution

The *normal random variable*  $X$ , whose density function  $f$  has the familiar bell-shaped curve, pictured in Fig. 41-1 below, is defined by:

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left[-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2\right]$$

This distribution is denoted by  $N(\mu, \sigma^2)$ .




Fig. 41-1

41.21. Mean  $N(\mu, \sigma^2) = \mu$ .

41.22. Variance  $N(\mu, \sigma^2) = \sigma^2$ , and standard deviation of  $N(\mu, \sigma^2) = \sigma$ .

### 68-95-99.7 Rule

The normal distribution  $N(\mu, \sigma^2)$  follows the “**68-95-99.7 Rule**” which is illustrated in Fig. 41-1. That is:

- (a) About 68% (more precisely 68.3%) or just over two-thirds of the data points (population) fall within 1 standard deviation from the mean.
- (b) About 95% (more precisely 95.4%) of the data points (population) fall within 2 standard deviations from the mean.
- (c) About 99.7% of the data points (population) fall within 3 standard deviations from the mean.

### Standardized Normal Distribution

Corresponding to  $X = N(\mu, \sigma^2)$  there is the standardized random variable

$$Z = \frac{X - \mu}{\sigma}$$

which is also normal with mean  $\mu = 0$  and standard deviation  $\sigma = 1$ . Thus  $Z = N(0, 1)$ . The density function  $\phi$  for  $Z$  follows:

$$\phi(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}$$

Any value of  $x$  in a normal random variable  $X$  can be changed into a  $z$ -value by the formula:  $z = (x - \mu)/\sigma$ .

### Evaluating Standard Normal Probabilities

Table 36 gives the area under the standard normal curve  $\phi(z)$  between 0 and  $z$ , as indicated by the picture in the table. This area is denoted by  $\Phi(z)$ .

Using Table 36 and the symmetry of the curve, we can find  $P(z_1 \leq Z \leq z_2)$ , the area under the curve between any two values  $z_1$  and  $z_2$ , as pictured in Fig. 41-2 below:

- (a)  $z_1 < 0 < z_2$
- (b)  $0 < z_1 < z_2$
- (c)  $z_1 < z_2 < 0$




Fig. 41-2

**EXAMPLE 41.4:** Find the following probabilities:

(a)  $P(-0.5 \leq Z \leq 1.3)$ , (b)  $P(1.4 \leq Z \leq 2.0)$ , (c)  $P(-1.5 \leq Z \leq -0.8)$

(a)  $P(-0.5 \leq Z \leq 1.3) = \Phi(1.3) + \Phi(0.5) = 0.4032 + 0.1915 = 0.5947 = 59.47\%$ .

(b)  $P(1.4 \leq Z \leq 2.0) = \Phi(2.0) - \Phi(1.4) = 0.4772 - 0.4192 = 0.0580 = 5.80\%$

(c)  $P(-1.5 \leq Z \leq -0.8) = \Phi(1.5) - \Phi(0.8) = 0.4332 - 0.2881 = 0.1451 = 14.51\%$

Also, using the fact that the total area under the standard normal curve is 1 and that half the area is  $1/2 = 0.5000$ , we can also find the “tail-end” of a one-sided probability of  $Z$  as shown in Fig. 41-3 below:




Fig. 41-3

(a)  $P(Z < z_1)$       (b)  $P(Z > z_1)$

**EXAMPLE 41.5:** Find the following probabilities:

(a)  $P(Z \leq 1.3)$  (b)  $P(Z \geq 1.5)$ , (c)  $P(Z \geq -1.4)$ ,

(a)  $P(Z \leq 1.3) = 0.5 + \Phi(1.3) = 0.5000 + 0.4032 = 0.9032$ .

(b)  $P(Z \geq 1.5) = 0.5 - \Phi(1.5) = 0.5000 - 0.4332 = 0.0668 = 6.68\%$

(c)  $P(Z \leq -1.6) = 0.5 - \Phi(1.6) = 0.5000 - 0.4452 = 0.0548 = 5.48\%$

## Evaluating Arbitrary Normal Probabilities

Suppose  $X$  is the normal distribution  $N(\mu, \sigma^2)$ . To evaluate  $P(a \leq X \leq b)$  we first change  $a$  and  $b$  into standard units using:

$$z_1 = \frac{a - \mu}{\sigma} \text{ and } z_2 = \frac{b - \mu}{\sigma}$$

Then we can rewrite  $P(a \leq X \leq b)$  as a  $z$ -equation:

$$P(a \leq X \leq b) = P(z_1 \leq Z \leq z_2)$$

which is the area under the standard normal curve between  $z_1$  and  $z_2$ .

**EXAMPLE 41.6:** Suppose the heights of American men (in inches) are (approximately) normally distributed with mean  $\mu = 68$  and standard deviation  $\sigma = 2$ . Find the percentage P of American men who are:

(a) between  $a = 67$  and  $b = 71$  inches tall, (b) at least 6 feet (72 inches) tall.

(a) Transform a and b into standard units obtaining

$$z_1 = (67 - 68)/2 = -0.5 \text{ and } z_2 = (71 - 68)/2 = 1.5$$

Here  $z_1 < 0 < z_2$ . Hence

$$\begin{aligned} P &= P(67 \leq X \leq 71) = P(-0.5 \leq Z \leq 1.5) \\ &= \Phi(1.5) + \Phi(0.5) = 0.4332 + 0.1915 = 0.6247 \end{aligned}$$

That is, approximately 62.5 percent of American men are between 67 and 71 inches tall.

(b) Transform  $a = 72$  into standard units obtaining  $z = (72 - 68)/2 = 2.0$ . Here  $0 < z$ . Therefore,

$$P = P(X \geq 72) = P(Z \geq 2) = 0.5 - \Phi(2.0) = 0.5000 - 0.4772 = 0.0228$$

That is, approximately 2.3% of American men are at least 6 feet tall.

## Central Limit Theorem

---

Roughly speaking, the Central Limit Theorem (CLT) says that in any sequence of repeated trials the distribution of the standard sample mean approaches the standard normal distribution as the number of trials increases.

**41.23.** Central Limit Theorem: Let  $X_1, X_2, X_3, \dots$  be a sequence of independent random variables with the same distribution with mean  $\mu$  and variance  $\sigma^2$ . Let

$$\bar{X}_n = (X_1 + X_2 + \dots + X_n)/n \text{ and } Z_n = \frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}}$$

Then for large n and any interval  $\{a \leq x \leq b\}$ ,

$$P(a \leq Z_n \leq b) \approx P(a \leq \phi \leq b)$$

where  $\phi$  is the standard normal distribution.

## Additional Probability Distributions

---

**41.24.** Poisson Distribution:  $\Phi(x) = \sum_{t \leq x} \frac{\lambda^t e^{-\lambda}}{t!}$

**41.25.** Hypergeometric Distribution:  $\Phi(x) = \sum_{t \leq x} z \frac{\binom{r}{t} \binom{s}{n-t}}{\binom{r+s}{n}}$

**41.26.** Student's t Distribution:  $\Phi(x) = \frac{1}{\sqrt{n}\pi} \frac{\Gamma(\frac{n+1}{2})}{\Gamma(n/2)} \int_{-\infty}^x \left(1 + \frac{t^2}{n}\right)^{-(n+1)/2} dt$

$$41.27. \quad \chi^2 \text{ (Chi Square) Distribution: } \Phi(x) = \frac{1}{2^{n/2} \Gamma(n/2)} \int_0^x t^{(n-2)/2} e^{-t/2} dt$$

$$41.28. \quad F \text{ Distribution: } \Phi(x) = \frac{\Gamma\left(\frac{n_1+n_2}{2}\right) n_1^{n_1/2} n_2^{n_2/2}}{\Gamma(n_1/2) \Gamma(n_2/2)} \int_0^x t^{(n_1/2)-1} (n_2 + n_1 t)^{-(n_1+n_2)/2} dt$$

## Section XII: Numerical Methods

# 42 INTERPOLATION

### Lagrange Interpolation

#### Two-point formula

$$42.1. \quad p(x) = f(x_0) \frac{x - x_1}{x_0 - x_1} + f(x_1) \frac{x - x_0}{x_1 - x_0}$$

where  $p(x)$  is a linear polynomial interpolating two points

$$(x_0, f(x_0)), \quad (x_1, f(x_1)), \quad x_0 \neq x_1$$

#### General formula

$$42.2. \quad p(x) = f(x_0)L_{n,0}(x) + f(x_1)L_{n,1}(x) + \cdots + f(x_n)L_{n,n}(x)$$

where

$$L_{n,k} = \prod_{i=0, i \neq k}^n \frac{x - x_i}{x_k - x_i}$$

and where  $p(x)$  is an  $n$ th-order polynomial interpolating  $n + 1$  points

$$(x_k, f(x_k)), \quad k = 0, 1, \dots, n; \quad \text{and} \quad x_i \neq x_j \text{ for } i \neq j$$

#### Remainder formula

Suppose  $f(x) \in C^{n+1}[a, b]$ . Then there is a  $\xi(x) \in (a, b)$  such that:

$$42.3. \quad f(x) = p(x) + \frac{f^{n+1}(\xi(x))}{(n+1)!} (x - x_0)(x - x_1) \cdots (x - x_n)$$

### Newton's Interpolation

#### First-order divided-difference formula

$$42.4. \quad f[x_0, x_1] = \frac{f(x_1) - f(x_0)}{x_1 - x_0}$$

#### Two-point interpolatory formula

$$42.5. \quad p(x) = f(x_0) + f[x_0, x_1](x - x_0)$$

where  $p(x)$  is a linear polynomial interpolating two points

$$(x_0, f(x_0)), \quad (x_1, f(x_1)), \quad x_0 \neq x_1$$

**Second-order divided-difference formula**

$$42.6. \quad f[x_0, x_1, x_2] = \frac{f[x_1, x_2] - f[x_0, x_1]}{x_2 - x_0}$$

**Three-point interpolatory formula**

$$42.7. \quad p(x) = f(x_0) + f[x_0, x_1](x - x_0) + f[x_0, x_1, x_2](x - x_0)(x - x_1)$$

where  $p(x)$  is a quadratic polynomial interpolating three points

$$(x_0, f(x_0)), \quad (x_1, f(x_1)), \quad (x_2, f(x_2))$$

**General  $k$ th-order divided-difference formula**

$$42.8. \quad f[x_0, x_1, \dots, x_k] = \frac{f[x_1, x_2, \dots, x_k] - f[x_0, x_1, \dots, x_{k-1}]}{x_k - x_0}$$

**General interpolatory formula**

$$42.9. \quad p(x) = f(x_0) + f[x_0, x_1](x - x_0) + \dots + f[x_0, x_1, \dots, x_n](x - x_0)(x - x_1)\cdots(x - x_{n-1})$$

where  $p(x)$  is an  $n$ th-order polynomial interpolating  $n + 1$  points

$$(x_k, f(x_k)), \quad k = 0, 1, \dots, n; \quad \text{and} \quad x_i \neq x_j \text{ for } i \neq j$$

**Remainder formula**

Suppose  $f(x) \in C^{n+1}[a, b]$ . Then there is a  $\xi(x) \in (a, b)$  such that

$$42.10. \quad f(x) = p(x) + \frac{f^{n+1}(\xi(x))}{(n+1)!}(x - x_0)(x - x_1)\cdots(x - x_n)$$

---

**Newton's Forward-Difference Formula****First-order forward-difference at  $x_0$** 

$$42.11. \quad \Delta f(x_0) = f(x_1) - f(x_0)$$

**Second-order forward difference at  $x_0$** 

$$42.12. \quad \Delta^2 f(x_0) = \Delta f(x_1) - \Delta f(x_0)$$

**General  $k$ th-order forward difference at  $x_0$** 

$$42.13. \quad \Delta^k f(x_0) = \Delta^{k-1} f(x_1) - \Delta^{k-1} f(x_0)$$

**Binomial coefficient**

$$42.14. \quad \binom{s}{k} = \frac{s(s-1)\cdots(s-k+1)}{k!}$$

**Newton's forward-difference formula**

$$42.15. \quad p(x) = \sum_{k=0}^n \binom{n}{k} \Delta^k f(x_0)$$

where  $p(x)$  is an  $n$ th-order polynomial interpolating  $n + 1$  equal spaced points

$$(x_k, f(x_k)), \quad x_k = x_0 + kh \quad k = 0, 1, \dots, n$$

## Newton's Backward-Difference Formula

---

First-order backward difference at  $x_n$

$$42.16. \quad \nabla f(x_n) = f(x_n) - f(x_{n-1})$$

Second-order backward difference at  $x_n$

$$42.17. \quad \nabla^2 f(x_n) = \nabla f(x_n) - \nabla f(x_{n-1})$$

General  $k$ th-order backward difference at  $x_n$

$$42.18. \quad \nabla^k f(x_n) = \nabla^{k-1} f(x_n) - \nabla^{k-1} f(x_{n-1})$$

**Newton's backward-difference formula**

$$42.19. \quad p(x) = \sum_{k=0}^n (-1)^k \binom{-n}{k} \nabla^k f(x_n)$$

where  $p(x)$  is an  $n$ th-order polynomial interpolating  $n + 1$  equal spaced points

$$(x_k, f(x_k)), \quad x_k = x_0 + kh \quad k = 0, 1, \dots, n$$

## Hermite Interpolation

---

Two-point basis polynomials

$$42.20. \quad H_{1,0} = \left(1 - 2 \frac{x - x_0}{x_0 - x_1}\right) \frac{(x - x_1)^2}{(x_0 - x_1)^2}, \quad H_{1,1} = \left(1 - 2 \frac{x - x_1}{x_1 - x_0}\right) \frac{(x - x_0)^2}{(x_1 - x_0)^2}$$

$$\hat{H}_{1,0} = (x - x_0) \frac{(x - x_1)^2}{(x_0 - x_1)^2}, \quad \hat{H}_{1,1} = (x - x_1) \frac{(x - x_0)^2}{(x_1 - x_0)^2}$$

Two-point interpolatory formula

$$42.21. \quad H_3(x) = f(x_0)H_{1,0} + f(x_1)H_{1,1} + f'(x_0)\hat{H}_{1,0} + f'(x_1)\hat{H}_{1,1}$$

where  $H_3(x)$  is a third-order polynomial, agrees with  $f(x)$  and its first-order derivatives at two points, i.e.,

$$H_3(x_0) = f(x_0), \quad H'_3(x_0) = f'(x_0), \quad H_3(x_1) = f(x_1), \quad H'_3(x_1) = f'(x_1)$$

General basis polynomials

$$42.22. \quad H_{n,j} = \left(1 - 2 \frac{x - x_j}{L'_{n,j}(x_j)}\right) L_{n,j}^2(x), \quad \hat{H}_{n,j} = (x - x_j) L_{n,j}^2(x)$$

where

$$L_{n,j} = \prod_{i=0, i \neq j}^n \frac{x - x_i}{x_j - x_i}$$

### General interpolatory formula

$$42.23. \quad H_{2n+1}(x) = \sum_{j=0}^n f(x_j) H_{n,j}(x) + \sum_{j=0}^n f'(x_j) \hat{H}_{n,j}(x)$$

where  $H_{2n+1}(x)$  is a  $(2n + 1)$ th-order polynomial, agrees with  $f(x)$  and its first order derivatives at  $n + 1$  points, i.e.,

$$H_{2n+1}(x_k) = f(x_k), \quad H'_{2n+1}(x_k) = f'(x_k) \quad k = 0, 1, \dots, n$$

### Remainder formula

Suppose  $f(x) \in C^{2n+2}[a, b]$ . Then there is a  $\xi(x) \in (a, b)$  such that

$$42.24. \quad f(x) = H_{2n+1}(x) + \frac{f^{2n+2}(\xi(x))}{(2n+2)!} (x - x_0)^2 (x - x_1)^2 \cdots (x - x_n)^2$$

# 43 QUADRATURE

## Trapezoidal Rule

---

Trapezoidal rule

$$43.1. \int_a^b f(x) dx \sim \frac{b-a}{2} [f(a) + f(b)]$$

Composite trapezoidal rule

$$43.2. \int_a^b f(x) dx \sim \frac{h}{2} \left( f(a) + 2 \sum_{i=1}^{n-1} f(a + ih) + f(b) \right)$$

where  $h = (b - a)/n$  is the grid size.

## Simpson's Rule

---

Simpson's rule

$$43.3. \int_a^b f(x) dx \sim \frac{b-a}{6} \left[ f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right]$$

Composite Simpson's rule

$$43.4. \int_a^b f(x) dx \sim \frac{h}{3} \left( f(x_0) + 2 \sum_{i=2}^{n/2} f(x_{2i-2}) + 4 \sum_{i=1}^{n/2} f(x_{2i-1}) + f(x_n) \right)$$

where  $n$  even,  $h = (b - a)/n$ ,  $x_i = a + ih$ ,  $i = 0, 1, \dots, n$ .

## Midpoint Rule

---

Midpoint rule

$$43.5. \int_a^b f(x) dx \sim (b-a)f\left(\frac{a+b}{2}\right)$$

Composite midpoint rule

$$43.6. \int_a^b f(x) dx \sim 2h \sum_{i=0}^{n/2} f(x_{2i})$$

where  $n$  even,  $h = (b - a)/(n+2)$ ,  $x_i = a + (i-1)h$ ,  $i = -1, 0, \dots, n+1$ .

## Gaussian Quadrature Formula

### Legendre polynomial

$$43.7. \quad P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} [(x^2 - 1)^n]$$

### Abscissa points and weight formulas

The abscissa points  $x_k^{(n)}$  and weight coefficient  $\omega_k^{(n)}$  are defined as follows:

$$43.8. \quad x_k^{(n)} = \text{the } k\text{th zero of the Legendre polynomial } P_n(x)$$

$$43.9. \quad \omega_k^{(n)} = \frac{2P'_n(x_k^{(n)})^2}{1 - x_k^{(n)2}}$$

Tables for Gauss-Legendre abscissas and weights appear in Fig. 43-1.

### Gauss-Legendre formula in interval $(-1, 1)$

$$43.10. \quad \int_{-1}^1 f(x) dx = \sum_{k=1}^n \omega_k^{(n)} f(x_k^{(n)}) + R_n$$

### Gauss-Legendre formula in general interval $(a, b)$

$$43.11. \quad \int_a^b f(x) dx = \frac{b-a}{2} \sum_{k=1}^n \omega_k^{(n)} f\left(\frac{a+b}{2} + x_k^{(n)} \frac{b-a}{2}\right) + R_n$$

### Remainder formula

$$43.12. \quad R_n = \frac{(b-a)^{2n+1} (n!)^4}{(2n+1)[(2n)!]^3} f^{(2n)}(\xi)$$

for some  $a < \xi < b$ .

$n$	$x_k^{(n)}$	$\omega_k^{(n)}$
2	0.5773502692	1.0000000000
	-0.5773502692	1.0000000000
3	0.7745966692	0.5555555556
	0.0000000000	0.8888888889
	-0.7745966692	0.5555555556
4	0.8611363116	0.3478548451
	0.3399810436	0.6521451549
	-0.3399810436	0.6521451549
	-0.8611363116	0.3478548451
5	0.9061798459	0.2369268850
	0.5384693101	0.4786286705
	-0.0000000000	0.5688888889
	-0.5384693101	0.4786286705
	-0.9061798459	0.2369268850

Fig. 43-1

# 44 SOLUTION of NONLINEAR EQUATIONS

Here we give methods to solve nonlinear equations which come in two forms:

**44.1.** Nonlinear equation:  $f(x) = 0$

**44.2.** Fixed point nonlinear equation:  $x = g(x)$

One can change from 44.1 to 44.2 or from 44.2 to 44.1 by setting:

$$g(x) = f(x) + x \quad \text{or} \quad f(x) = g(x) - x$$

Since the methods are iterative, there are two types of error estimates:

**44.3.**  $|f(x_n)| < \epsilon$  or  $|x_{n+1} - x_n| < \epsilon$

for some preassigned  $\epsilon > 0$ .

## Bisection Method

The following theorem applies:

*Intermediate Value Theorem:* Suppose  $f$  is continuous on an interval  $[a, b]$  and  $f(a)f(b) < 0$ . Then there is a root  $x^*$  to  $f(x) = 0$  in  $(a, b)$ .

The bisection method approximates one such solution  $x^*$ .

**44.4.** Bisection method:

*Initial step:* Set  $a_0 = a$  and  $b_0 = b$ .

*Repetitive step:*

(a) Set  $c_n = (a_n + b_n)/2$ .

(b) If  $f(a_n)f(c_n) < 0$ , then set  $a_{n+1} = a_n$  and  $b_{n+1} = c_n$ ; else set  $a_{n+1} = c_n$  and  $b_{n+1} = b_n$ .

## Newton's Method

### Newton method

**44.5.**  $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$

### Quadratic convergence

**44.6.**  $\lim_{n \rightarrow \infty} \frac{|x_{n+1} - x^*|}{|x_n - x^*|^2} = \frac{f''(x^*)}{2(f'(x^*))^2}$

where  $x^*$  is a root of the nonlinear equation 44.1.

## Secant Method

---

### Secant method

$$44.7. \quad x_{n+1} = x_n - \frac{(x_n - x_{n-1})f(x_n)}{f(x_n) - f(x_{n-1})}$$

### Rate of convergence

$$44.8. \quad \lim_{n \rightarrow \infty} \frac{|x_{n+1} - x^*|}{|x_n - x^*| |x_{n-1} - x^*|} = \frac{f''(x^*)}{2(f'(x^*))^2}$$

where  $x^*$  is a root of the nonlinear equation 44.1.

## Fixed-Point Iteration

---

The following definition and theorem apply:

*Definition:* A function  $g$  from  $(a, b)$  to  $(a, b)$  is called a *contraction mapping* if

$$|g(x) - g(y)| \leq L|x - y| \quad \text{for any } x, y \in (a, b)$$

where  $L < 1$  is a positive constant.

*Fixed-point theorem:* Suppose that  $g$  is a contraction mapping on  $(a, b)$ . Then  $g$  has a unique fixed point in  $(a, b)$ .

Given such a contraction mapping  $g$ , the following method may be used.

### Fixed-point iteration

$$44.9. \quad x_{n+1} = g(x_n)$$

# 45

## NUMERICAL METHODS for ORDINARY DIFFERENTIAL EQUATIONS

Here we give methods to solve the following initial-value problem of an ordinary differential equation:

$$45.1. \quad \begin{cases} \frac{dx}{dt} = f(x, t) \\ x(t_0) = x_0 \end{cases}$$

The methods will use a computational grid:

$$45.2. \quad t_n = t_0 + nh$$

where  $h$  is the grid size.

### First-Order Methods

---

**Forward Euler method (first-order explicit method)**

$$45.3. \quad x(t+h) = x(t) + hf(x(t), t)$$

**Backward Euler method (first-order implicit method)**

$$45.4. \quad x(t+h) = x(t) + hf(x(t+h), t+h)$$

---

### Second-Order Methods

**Mid-point rule (second-order explicit method)**

$$45.5. \quad \begin{cases} x^* = x(t) + \frac{h}{2}f(x(t), t) \\ x(t+h) = x(t) + hf\left(x^*, t + \frac{h}{2}\right) \end{cases}$$

**Trapezoidal rule (second-order implicit method)**

$$45.6. \quad x(t+h) = x(t) + \frac{h}{2}\{f(x(t), t) + f(x(t+h), t+h)\}$$

**Heun's method (second-order explicit method)**

$$45.7. \quad \begin{cases} x^* = x(t) + hf(x(t), t) \\ x(t+h) = x(t) + \frac{h}{2}\{f(x(t), t) + f(x^*, t+h)\} \end{cases}$$

## Single-Stage High-Order Methods

---

### Fourth-order Runge–Kutta method (fourth-order explicit method)

$$45.8. \quad x(t+h) = x(t) + \frac{1}{6}(F_1 + 2F_2 + 2F_3 + F_4)$$

where

$$F_1 = hf(x, t), \quad F_2 = hf\left(x + \frac{F_1}{2}, t + \frac{h}{2}\right), \quad F_3 = hf\left(x + \frac{F_2}{2}, t + \frac{h}{2}\right), \quad F_4 = hf(x + F_3, t + h)$$

## Multi-Step High-Order Methods

---

### Adams-Basforth two-step method

$$45.9. \quad x(t+h) = x(t) + h\left(\frac{3}{2}f(x(t), t) - \frac{1}{2}f(x(t-h), t-h)\right)$$

### Adams-Basforth three-step method

$$45.10. \quad x(t+h) = x(t) + h\left(\frac{23}{12}f(x(t), t) - \frac{4}{3}f(x(t-h), t-h) + \frac{5}{12}f(x(t-2h), t-2h)\right)$$

### Adams-Basforth four-step method

$$45.11. \quad x(t+h) = x(t) + h\left(\frac{55}{24}f(x(t), t) - \frac{59}{24}f(x(t-h), t-h) + \frac{37}{24}f(x(t-2h), t-2h) - \frac{9}{24}f(x(t-3h), t-3h)\right)$$

### Milne's method

$$45.12. \quad x(t+h) = x(t-3h) + h\left(\frac{8}{3}f(x(t), t) - \frac{4}{3}f(x(t-h), t-h) + \frac{8}{3}f(x(t-2h), t-2h)\right)$$

### Adams-Moulton two-step method

$$45.13. \quad x(t+h) = x(t) + h\left(\frac{5}{12}f(x(t+h), t+h) + \frac{2}{3}f(x(t), t) - \frac{1}{12}f(x(t-h), t-h)\right)$$

### Adams-Moulton three-step method

$$45.14. \quad x(t+h) = x(t) + h\left(\frac{3}{8}f(x(t+h), t+h) + \frac{19}{24}f(x(t), t) - \frac{5}{24}f(x(t-h), t-h) + \frac{1}{24}f(x(t-2h), t-3h)\right)$$

# 46 NUMERICAL METHODS for PARTIAL DIFFERENTIAL EQUATIONS

## Finite-Difference Method for Poisson Equation

The following is the Poisson equation in a domain  $(a, b) \times (c, d)$ :

$$46.1. \quad \nabla^2 u = f, \quad \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$$

Boundary condition:

$$46.2. \quad u(x, y) = g(x, y) \quad \text{for } x = a, b \quad \text{or} \quad y = c, d$$

Computation grid:

$$46.3. \quad \begin{aligned} x_i &= a + i\Delta x && \text{for } i = 0, 1, \dots, n \\ y_j &= c + j\Delta y && \text{for } j = 0, 1, \dots, m \end{aligned}$$

where  $\Delta x = (b - a)/n$  and  $\Delta y = (d - c)/m$  are grid sizes for  $x$  and  $y$  variables, respectively.

### Second-order difference approximation

$$46.4. \quad (D_x^2 + D_y^2)u(x_i, y_j) = f(x_i, y_j)$$

where

$$\begin{aligned} D_x^2 u(x_i, y_j) &= \frac{u(x_{i+1}, y_j) - 2u(x_i, y_j) + u(x_{i-1}, y_j)}{\Delta x^2} \\ D_y^2 u(x_i, y_j) &= \frac{u(x_i, y_{j+1}) - 2u(x_i, y_j) + u(x_i, y_{j-1})}{\Delta y^2} \end{aligned}$$

### Computational boundary condition

$$46.5. \quad \begin{aligned} u(x_0, y_j) &= g(a, y_j), & u(x_n, y_j) &= g(b, y_j) && \text{for } j = 1, 2, \dots, m \\ u(x_i, y_0) &= g(x_i, c), & u(x_i, y_m) &= g(x_i, d) && \text{for } i = 1, 2, \dots, n \end{aligned}$$

## Finite-Difference Method for Heat Equation

The following is the heat equation in a domain  $(a, b) \times (c, d) \times (0, T)$ :

$$46.6. \quad \frac{\partial u}{\partial t} = \nabla^2 u$$

Boundary condition:

$$46.7. \quad u(x, y, t) = g(x, y) \quad \text{for } x = a, b \quad \text{or} \quad y = c, d$$

Initial condition:

$$46.8. \quad u(x, y, 0) = u_0(x, y)$$

Computational grid:

$$\begin{aligned} 46.9. \quad x_i &= a + i\Delta x && \text{for } i = 0, 1, \dots, n \\ y_j &= c + j\Delta y && \text{for } j = 0, 1, \dots, m \\ t_k &= k\Delta t && \text{for } k = 0, 1, \dots, \end{aligned}$$

where  $\Delta x = (b - a)/n$ ,  $\Delta y = (d - c)/m$ , and  $\Delta t$  are grid sizes for  $x$ ,  $y$  and  $t$  variables, respectively.

### Computational boundary condition

$$\begin{aligned} 46.10. \quad u(x_0, y_j) &= g(a, y_j), u(x_n, y_j) = g(b, y_j) && \text{for } j = 1, 2, \dots, m \\ u(x_i, y_0) &= g(x_i, c), u(x_i, y_m) = g(x_i, d) && \text{for } i = 1, 2, \dots, n \end{aligned}$$

### Computational initial condition

$$46.11. \quad u(x_i, y_j, 0) = u_0(x_i, y_j) \quad \text{for } i = 1, 2, \dots, n; j = 0, 1, \dots, m$$

### Forward Euler method with stability condition

$$46.12. \quad u(x_i, y_j, t_{k+1}) = u(x_i, y_j, t_k) + \Delta t(D_x^2 + D_y^2)u(x_i, y_j, t_k)$$

$$46.13. \quad \frac{2\Delta t}{\Delta x^2} + \frac{2\Delta t}{\Delta y^2} \leq 1$$

### Backward Euler method (unconditional stable)

$$46.14. \quad u(x_i, y_j, t_{k+1}) = u(x_i, y_j, t_k) + \Delta t(D_x^2 + D_y^2)u(x_i, y_j, t_{k+1})$$

### Crank-Nicholson method (unconditional stable)

$$46.15. \quad u(x_i, y_j, t_{k+1}) = u(x_i, y_j, t_k) + \Delta t(D_x^2 + D_y^2)\{u(x_i, y_j, t_k) + u(x_i, y_j, t_{k+1})\}/2$$

## Finite-Difference Method for Wave Equation

---

The following is a wave equation in a domain  $(a, b) \times (c, d) \times (0, T)$ :

$$46.16. \quad \frac{\partial^2 u}{\partial t^2} = A^2 \nabla^2 u$$

where  $A$  is a constant representing the speed of the wave.

Boundary condition:

$$46.17. \quad u(x, y, t) = g(x, y) \quad \text{for } x = a, b \text{ or } y = c, d$$

Initial condition:

$$46.18. \quad u(x, y, 0) = u_0(x, y), \quad \frac{\partial u}{\partial t}(x, y, 0) = u_i(x, y)$$

Computational grids:

$$\begin{aligned} 46.19. \quad x_i &= a + i\Delta x && \text{for } i = 0, 1, \dots, n \\ y_j &= c + j\Delta y && \text{for } j = 0, 1, \dots, m \\ t_k &= k\Delta t && \text{for } k = -1, 0, 1, \dots \end{aligned}$$

where  $\Delta x = (b - a)/n$ ,  $\Delta y = (d - c)/m$ , and  $\Delta t$  are the grid sizes for  $x$ ,  $y$ , and  $t$  variables, respectively.

#### A second-order finite-difference approximation

$$46.20. \quad u(x_i, y_j, t_{k+1}) = 2u(x_i, y_j, t_k) - u(x_i, y_j, t_{k-1}) + \Delta t^2 A^2 (D_x^2 + D_y^2) u(x_i, y_j, t_k)$$

#### Computational boundary condition

$$\begin{aligned} 46.21. \quad u(x_0, y_j) &= g(a, y_j), u(x_n, y_j) = g(b, y_j) && \text{for } j = 1, 2, \dots, m \\ u(x_i, y_0) &= g(x_i, c), u(x_i, y_m) = g(x_i, d) && \text{for } i = 1, 2, \dots, n \end{aligned}$$

#### Computational initial condition

$$46.22. \quad u(x_i, y_j, t_0) = u_0(x_i, y_j) \quad \text{for } i = 1, 2, \dots, n; j = 0, 1, \dots, m$$

$$u(x_i, y_j, t_{-1}) = u_0(x_i, y_j) + \Delta t^2 u_i(x_i, y_j) \quad \text{for } i = 1, 2, \dots, n; j = 0, 1, \dots, m$$

#### Stability condition

$$46.23. \quad \Delta t \leq A \min(\Delta x, \Delta y)$$

# 47 ITERATION METHODS for LINEAR SYSTEMS

## Iteration Methods for Poisson Equation

The finite-difference approximation to the Poisson equation follows:

$$47.1. \quad \begin{cases} u_{i+1,j} + u_{i-1,j} + u_{i,j+1} + u_{i,j-1} - 4u_{i,j} = f_{i,j} & \text{for } i, j = 1, 2, \dots, n-1 \\ u_{0,j} = u_{n,j} = 0 & \text{for } j = 1, 2, \dots, n-1 \\ u_{i,0} = u_{i,n} = 0 & \text{for } i = 1, 2, \dots, n-1 \end{cases}$$

Three iteration methods for solving the system follow:

### Jacobi method

$$47.2. \quad u_{i,j}^{k+1} = \frac{1}{4}(u_{i+1,j}^k + u_{i-1,j}^k + u_{i,j+1}^k + u_{i,j-1}^k - f_{i,j})$$

### Gauss-Seidel method

$$47.3. \quad u_{i,j}^{k+1} = \frac{1}{4}(u_{i+1,j}^k + u_{i-1,j}^{k+1} + u_{i,j+1}^k + u_{i,j-1}^{k+1} - f_{i,j})$$

### Successive-overrelaxation (SOR) method

$$47.4. \quad \begin{cases} u_{i,j}^* = \frac{1}{4}(u_{i+1,j}^k + u_{i-1,j}^* + u_{i,j+1}^k + u_{i,j-1}^* - f_{i,j}) \\ u_{i,j}^{k+1} = (1 - \omega)u_{i,j}^k + \omega u_{i,j}^* \end{cases}$$

## Iteration Methods for General Linear Systems

Consider the linear system

$$47.5. \quad Ax = b$$

where  $A$  is an  $n \times n$  matrix and  $x$  and  $b$  are  $n$ -vectors. We assume the coefficient matrix  $A$  is partitioned as follows:

$$47.6. \quad A = D - L - U$$

where  $D = \text{diag}(A)$ ,  $L$  is the negative of the strictly lower triangular part of  $A$ , and  $U$  is the negative of the strictly upper triangular part of  $A$ .

Four iteration methods for solving the system follow:

**Richardson method**

$$47.7. \quad x^{k+1} = (I - A)x^k + b$$

**Jacobi method**

$$47.8. \quad Dx^{k+1} = (L + U)x^k + b$$

**Gauss-Seidel method**

$$47.9. \quad (D - L)x^{k+1} = Ux^k + b$$

**Successive-overrelaxation (SOR) method**

$$47.10. \quad (D - \omega L)x^{k+1} = \omega(Ux^k + b) + (1 - \omega)Dx^k$$

## Section XIII: Turing Machines

# 48 BASIC DEFINITIONS, EXPRESSIONS

A *Turing machine M* involves three disjoint nonempty sets:

- (1) A finite *tape* set where  $B = a_0$  is the “blank” symbol:

$$A = \{a_1, a_2, \dots, a_m\} \cup \{B\}$$

- (2) A finite *state* set where  $s_0$  is the *initial state*:

$$S = \{s_1, s_2, \dots, s_n\} \cup \{s_0\} \cup \{s_H, s_Y, s_N\}$$

Here  $s_H$  (HALT) is the halting state,  $s_Y$  (YES) is the accepting state, and  $s_N$  (NO) is the nonaccepting state.

- (3) A *direction* set where  $L$  denotes “left” and  $R$  denotes “right”:

$$d = \{L, R\}$$

**Definition 48.1:** An *expression* is a finite (possibly empty) sequence of elements from  $A \cup S \cup d$ .

In other words, an expression is a word whose letters (symbols) come from the sets  $A$ ,  $S$ , and  $d$ .

**Definition 48.2:** A *tape expression* is an expression using only elements from the tape set  $A$ .

# 49 PICTURES

The Turing machine  $M$  may be viewed as a read/write tape head that moves back and forth along an infinite tape. The tape is divided lengthwise into squares (cells), and each square may be blank or hold one tape symbol. At each step in time, the Turing machine  $M$  is in a certain internal state  $s_i$  scanning one of the tape symbols  $a_j$  on the tape. We assume that only a finite number of nonblank symbols appear on the tape.

Figure 49-1(a) is a picture of a Turing machine  $M$  in state  $s_2$  scanning the second symbol where  $a_1 a_3 B a_1 a_1$  is printed on the tape. (Note again that  $B$  is the blank symbol.) This picture may be represented by the expression  $\alpha = a_1 s_2 a_3 B a_1 a_1$ , where we write the state  $s_2$  of  $M$  before the tape symbol  $a_3$  that  $M$  is scanning. Observe that  $\alpha$  is an expression using only the tape alphabet  $A$  except for the state symbol  $s_2$ , which is not at the end of the expression since it appears before the tape symbol  $a_3$  that  $M$  is scanning. Figure 49-1 shows two other informal pictures and their corresponding picture expressions.




Fig. 49-1

We give formal definitions.

**Definition 49.1:** A *picture*  $\alpha$  is an expression as follows where  $P$  and  $Q$  are tape expressions (possibly empty):

$$\alpha = Ps_i a_k Q$$

**Definition 49.2:** Let  $\alpha = Ps_i a_k Q$  be a picture. We say that the Turing machine  $M$  is in state  $s_i$  scanning the letter  $a_k$  and that the *expression on the tape* is the expression  $Pa_k Q$ , that is, without its state symbol  $s_i$ .

# 50

## QUINTUPLE, TURING MACHINE

As mentioned above, at each step in time the Turing machine  $M$  is in a certain state  $s_i$  and is scanning a tape symbol  $a_k$ . The Turing machine  $M$  is able to do the following three things simultaneously:

- (i)  $M$  erases the scanned symbol  $a_k$  and writes a tape symbol  $a_l$  (where we permit  $a_l = a_k$ ) in its place.
- (ii)  $M$  changes its internal state  $s_i$  to a state  $s_j$  (where we permit  $s_j = s_i$ ).
- (iii)  $M$  moves one square to the left or one square to the right.

### Quintuple

The above action by  $M$  may be described by a five-letter expression called a *quintuple*, which we define below.

**Definition 50.1:** A quintuple  $q$  is a five-letter expression of the following form:

$$q = \left( s_i, a_k, a_l, s_j, \begin{cases} L \\ R \end{cases} \right)$$

### Turing Machine

That is, the first letter of  $q$  is a state symbol, the second is a tape symbol, the third is a tape symbol, the fourth is a state symbol, and the last is a direction symbol  $L$  or  $R$ .

Next we give a formal definition of a Turing machine.

**Definition 50.2:** A Turing machine  $M$  is a finite set of quintuples such that:

- (i) No two quintuples begin with the same first two letters.
- (ii) No quintuple begins with  $s_H$ ,  $s_Y$  or  $s_N$ .

Condition (i) in the definition guarantees that the machine  $M$  cannot do more than one thing at any given step, and condition (ii) guarantees that  $M$  halts in state  $s_H$ ,  $s_Y$  or  $s_N$ .

The following is an alternative equivalent definition.

**Definition 50.2':** Turing machine  $M$  is a partial function from

$$S \setminus \{s_H, s_Y \text{ or } s_N\} \times A \text{ into } A \times S \times d$$

The term *partial function* simply means that the domain of  $M$  is a subset of  $S \setminus \{s_H, s_Y \text{ or } s_N\} \times A$ .

### Action of Turing Machine

The action of the Turing machine described above can now be formally defined.

**Definition 50.3:** Let  $\alpha$  and  $\beta$  be pictures. We write

$$\alpha \rightarrow \beta$$

if one of the following holds where  $a$ ,  $b$ , and  $c$  are tape letters, and  $P$  and  $Q$  are tape expressions (possibly empty):

- (i)  $\alpha = Ps_i acQ$ ,  $\beta = Pbs_j cQ$ , and  $M$  contains the quintuple  $q = s_i abs_j R$ .
- (ii)  $\alpha = Pcs_i aQ$ ,  $\beta = Ps_j cbQ$ , and  $M$  contains the quintuple  $q = s_i abs_j L$ .
- (iii)  $\alpha = Ps_i a$ ,  $\beta = Pbs_j B$ , and  $M$  contains the quintuple  $q = s_i abs_j R$ .
- (iv)  $\alpha = s_i aQ$ ,  $\beta = s_j BbQ$ , and  $M$  contains the quintuple  $q = s_i abs_j L$ .

Observe that, in all four cases,  $M$  replaces  $a$  on the tape with  $b$  (where we permit  $b = a$ ), and  $M$  changes its state from  $s_i$  to  $s_j$  (where we permit  $s_j = s_i$ ). Furthermore:

- (i) Here  $M$  moves to the right.
- (ii) Here  $M$  moves to the left.
- (iii) Here  $M$  moves to the right; however, since  $M$  is scanning the rightmost letter, it must add the blank symbol  $B$  on the right.
- (iv) Here  $M$  moves to the left; however, since  $M$  is scanning the leftmost letter, it must add the blank symbol  $B$  on the left.

**Definition 50.4:** A picture  $\alpha$  is said to be *terminal* if there is no picture  $\beta$  such that  $\alpha \rightarrow \beta$ .

In particular, any picture  $\alpha$  in one of the three halt states must be terminal since no quintuple begins with  $s_H$ ,  $s_Y$ , or  $s_n$ .

# 51 COMPUTING WITH A TURING MACHINE

The above is a static (one-step) description of a Turing machine  $M$ . Now we discuss its dynamics.

**Definition 51.1:** A *computation* of a Turing machine  $M$  is a sequence of pictures  $\alpha_1, \alpha_2, \dots, \alpha_m$  such that  $\alpha_{i-1} \rightarrow \alpha_i$ , for  $i = 1, 2, \dots, m$ , and  $\alpha_m$  is a terminal picture.

In other words, a computation is a sequence

$$\alpha_0 \rightarrow \alpha_1 \rightarrow \alpha_2 \rightarrow \dots \rightarrow \alpha_m$$

that cannot be extended since  $\alpha_m$  is terminal. We will let  $\text{term}(\alpha)$  denote the final picture of a computation beginning with  $\alpha$ . Thus  $\text{term}(\alpha_0) = \alpha_m$  in the above computation.

## Turing Machines with Input

The following definition applies.

**Definition 51.2:** An *input* for a Turing machine  $M$  is a tape expression  $W$ . The *initial picture* for an input  $W$  is  $\alpha(W)$  where  $\alpha(W) = s_0(W)$ .

Observe that the initial picture  $\alpha(W)$  of the input  $W$  is obtained by placing the initial state  $s_0$ , in front of the input tape expression  $W$ . In other words, the Turing machine  $M$  begins in its initial state  $s_0$ , and it is scanning the first letter of  $W$ .

**Definition 51.3:** Let  $M$  be a Turing machine, and let  $W$  be an input. We say  $M$  halts on  $W$  if there is a computation beginning with the initial picture  $\alpha(W)$ .

That is, given an input  $W$ , we can form the initial picture  $\alpha(W) = s_0(W)$  and apply  $M$  to obtain the sequence

$$\alpha(W) \rightarrow \alpha_1 \rightarrow \alpha_2 \rightarrow \dots$$

Two things can happen:

- (1)  **$M$  halts on  $W$ .** That is, the sequence ends with some terminal picture  $\alpha_r$ .
- (2)  **$M$  does not halt on  $W$ .** That is, the sequence never ends.

## Computable Functions

Computable functions are defined on the set of nonnegative integers. Some texts use  $\mathbf{N}$  to denote this set. We use  $\mathbf{N}$  to denote the set of positive integers, so we will use the notation

$$\mathbf{N}_0 = \{0, 1, 2, 3, \dots\}$$

Throughout this section, the terms *number*, *integer*, and *nonnegative integer* are used synonymously. Here we show how  $M$  manipulates numerical data. First, however, we need to be able to represent our numbers with our tape set  $A$ . We will write  $1$  for the tape symbol  $a_1$  and  $1^n$  for  $111\dots 1$ , where  $1$  occurs  $n$  times.

**Definition 51.4:** Each number  $n$  will be represented by the tape expression  $\langle n \rangle$  where  $\langle n \rangle = 1^{n+1}$ . Thus,

$$\langle 4 \rangle = 11111 = 1^5, \quad \langle 0 \rangle = 1, \quad \langle 2 \rangle = 111 = 1^3.$$

**Definition 51.5:** Let  $E$  be an expression. Then  $[E]$  will denote the number of times 1 occurs in  $E$ . Thus,

$$[11Bs_2 a_3 111Ba_4] = 5, \quad [a_4s_2Ba_2] = 0, \quad [\langle n \rangle] = n + 1.$$

**Definition 51.6:** A function  $f: \mathbf{N}_0 \rightarrow \mathbf{N}_0$  is computable if there exists a Turing machine  $M$  such that, for every integer  $n$ ,  $M$  halts on  $\langle n \rangle$  and

$$f(n) = [\text{term}(\alpha(\langle n \rangle))].$$

We then say that  $M$  computes  $f$ .

That is, given a function  $f$  and an integer  $n$ , we input  $\langle n \rangle$  and apply  $M$ . If  $M$  always halts on  $\langle n \rangle$  and the number of 1s in the final picture is equal to  $f(n)$ , then  $f$  is a computable function, and we say that  $M$  computes  $f$ .

# 52 EXAMPLES

**Example 1:** Let  $M$  be a Turing machine. Determine the picture  $\alpha$  corresponding to each situation:

- (a)  $M$  is in state  $s_3$  and scanning the third letter of the tape expression  $w = aabca$ .
- (b)  $M$  is in state  $s_2$  and scanning the last letter of the tape expression  $w = abca$ .
- (c) The input is the tape expression  $w = 1^4B1^2$ .

The picture  $\alpha$  is obtained by placing the state symbol before the tape letter being scanned. Initially,  $M$  is in state  $s_0$  scanning the first letter of an input. Thus,

$$(a) \alpha = aas_3bca; \quad (b) \alpha = abcs_2a; \quad (c) \alpha = s_01111B11.$$

**Example 2:** Suppose  $\alpha = aas_2ba$  is a picture. Find  $\beta$  such that  $\alpha \rightarrow \beta$  if the Turing machine  $M$  has the quintuple  $q$  where (a)  $q = s_2bas_1L$ ; (b)  $q = s_2bbs_3R$ ; (c)  $q = s_2bas_2R$ ; and (d)  $q = s_3abs_1L$ .

- (a) Here  $M$  erases  $b$  and writes  $a$ , changes its state to  $s_1$ , and moves left. Thus,  $\beta = as_1aaa$ .
- (b) Here  $M$  does not change the scanned letter  $b$ , changes its state to  $s_3$ , and moves right. Thus,  $\beta = aabs_3a$ .
- (c) Here  $M$  erases  $b$  and writes  $a$ , keeps its state  $s_2$ , and moves right. Thus,  $\beta = aaas_2a$ .
- (d) Here  $q$  has no effect on  $\alpha$  since  $q$  does not begin with  $s_2b$ .

## Computable Functions

**Example 3:** Find  $\langle m \rangle$  if (a)  $m = 5$ ; (b)  $m = (4, 0, 3)$ ; and (c)  $m = (3, -2, 5)$ .

Recall  $\langle n \rangle = 1^{n+1} = 11^n$  and  $\langle (n_1, n_2, \dots, n_r) \rangle = \langle n_1 \rangle B \langle n_2 \rangle B \dots B \langle n_r \rangle$ . Thus,

- (a)  $\langle m \rangle = 1^6 = 111111$ .
- (b)  $\langle m \rangle = 1^5B1^1B1^4 = 11111B1B1111$ .
- (c)  $\langle m \rangle$  is not defined for negative integers.

**Example 4:** Find  $[E]$  for the expressions:

- (a)  $E = aas_2Bb111$ ; (c)  $E = \langle m \rangle$  where  $m = (4, 1, 2)$ ;
- (b)  $E = aas_3bb$ ; (d)  $E = \langle m \rangle$  where  $m = (n_1, n_2, \dots, n_r)$ .

Recall that  $[E]$  counts the number of 1s in  $E$ . Thus,

- (a)  $[E] = 5$ ; (b)  $[E] = 0$ ; (c)  $[E] = 10$  since  $E = 1^5B1^2B1^3$ ;
- (d)  $[E] = n_1 + n_2 + \dots + n_r + r$  since the number of 1s contributed by each  $n_k$  to  $E$  is  $n_k + 1$ .

**Example 5:** The function  $f(n) = n + 3$  is computable. The input is  $W = 1^{n+1}$ . Thus we need only add two 1's to the input. A Turing machine  $M$  that computes  $f$  follows

$$M = \{q_1, q_2, q_3\} = \{s_0 1s_0L, s_0B1s_1L, s_1B_1s_HL\}$$

Observe that:

- (1)  $q_1$  moves the machine  $M$  to the left.
- (2)  $q_2$  writes 1 in the blank square  $B$  and moves  $M$  to the left.
- (3)  $q_3$  writes 1 in the blank square  $B$  and halts  $M$ .

Accordingly, for any positive integer  $n$ ,

$$s_0 1^{n+1} \rightarrow s_0 B 1^{n+1} \rightarrow s_1 B 1^{n+2} \rightarrow s_H B 1^{n+3}.$$

Thus  $M$  computes  $f(n) = n + 3$ . It is clear that, for any positive integer  $k$ , the function  $f(n) = n + k$  is computable.

**Example 6:** Let  $f$  be the function  $f(n) = n - 1$ , when  $n > 0$  and  $f(0) = 0$ . Show that  $f$  is computable.

We need to find a Turing machine  $M$  that computes  $f$ . Specifically, we want  $M$  to erase two of the 1s in the input  $\langle n \rangle$  when  $n > 0$ , but only one 1 when  $n = 0$ . This is accomplished with the following quintuples:

$$q_1 = s_0 1 B s_1 R, \quad q_2 = s_1 B B s_H R, \quad q_3 = s_1 1 B s_H R$$

Here  $q_1$  erases the first 1 and moves  $M$  right. If there is only one 1, then  $M$  is now scanning a blank symbol  $B$ , and  $q_2$  tells the computer to halt. Otherwise,  $q_3$  erases the second 1 and halts  $M$ .

## Section XIV: Mathematical Finance

# 53 BASIC PROBABILITY

**53.1.** Standard normal random variable  $Z$

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$

$$\Phi(x) = P(Z \leq x) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-z^2/2} dz$$

**53.2.** Normal random variable  $X$

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$$

where  $\mu$  is the expected value, and  $\sigma^2$  is the variance.

$Z = \frac{X - \mu}{\sigma}$  is a standard normal random variable.

**53.3.** Lognormal random variable  $Y$

$$Y = e^X, X \text{ is a normal random variable}$$

$$E(Y) = e^{\mu + \sigma^2/2}$$

$$\text{Var } Y = e^{2\mu + 2\sigma^2} - e^{2\mu + \sigma^2}$$

**53.4.** Discrete-time Brownian motion with drift parameter  $\mu$  and variance parameter  $\sigma^2$ , for every  $\Delta$  time units,

$$X(t + \Delta) = x(t) + X, \quad X \text{ normal random variable with } \mu\Delta, \text{ and } \sigma^2\Delta$$

**53.5.** Binomial model for Brownian motion

$$X(n\Delta) = X(0) + \sigma\sqrt{\Delta}(X_1 + \dots + X_n)$$

$$X_i = \begin{cases} 1 & \text{with probability } p \\ -1 & \text{with probability } 1-p \end{cases}$$

where

$$p = \frac{1}{2} \left( 1 + \frac{\mu}{\sigma} \sqrt{\Delta} \right)$$

$$E(X(n\Delta) - X(0)) = \mu n\Delta$$

$$\text{Var}(X(n\Delta) - X(0)) = \sigma^2 n\Delta - \mu^2 n\Delta^2 \rightarrow \sigma^2 t, \text{ as } \Delta \rightarrow 0, n\Delta \rightarrow t$$

By central limit theorem

$$x(t) - x(0) \sim N(\mu t, \sigma^2 t),$$

$x(t)$  approximates Brownian motion with drift parameter  $\mu$  and variance parameter  $\sigma^2$ .

**53.6.** Geometric Brownian motion with drift parameter  $\mu$  and variance parameter  $\sigma^2$

$$S(t) = e^{X(t)},$$

where  $X(t)$  is a Brownian motion with drift parameter  $\mu$  and variance parameter  $\sigma^2$ .

**53.7.** Discrete-time geometric Brownian motion

$$S(t + \Delta) = S(t)e^{X(t+\Delta)-X(t)}$$


**53.8.** Binomial model for geometric Brownian motion

$$P = \frac{1}{2} \left( 1 + \frac{\mu}{\sigma} \sqrt{\Delta} \right), \quad u = e^{\sigma \sqrt{\Delta}}, \quad d = e^{-\sigma \sqrt{\Delta}}$$

with probability  $p$  up  $u$  factor and probability  $1 - p$  down  $d$  factor.

**53.9.** Multi-period binomial stock model:

The price  $S(i)$  of a stock at the end of period  $i$ ,  $i = 1, 2, \dots, n$ . The period is  $\Delta$  and hence  $t_i = i\Delta$ . The initial price  $S(0)$  is non-random and fixed. The stock price  $S(i)$  is random, going up or down probabilistic rules:



# 54

## INTEREST RATES

**54.1.** Borrow an amount  $P$  (called the principal) for a time  $T$  with simple interest rate  $r$ . The amount to be repaid at time  $T$  is  $(1 + r)P$ .

**54.2.** Borrow an amount  $P$  for  $t$  years at a nominal interest rate of  $r$  per year compounded continuously. The amount owed at time  $t$  years is  $Pe^r$ .

**54.3.** The present value of a payoff of  $v$  to be made at time period  $i$  is  $v(1 + r)^{-i}$ .

# 55

## ARBITRAGE THEOREM AND OPTIONS

**55.1.** A sure-win investment scheme is called an *arbitrage*.

**55.2.** The law of one price: two investments with the same payoff, then either there is an arbitrage or the two investments have the same costs.

**55.3.** Selling it short: you sell a stock that you do not own.

**55.4.** European-style call option: option of calling/buying for the stock at a specified price  $K$  (known as the *exercise* or *strike* price) at the expiration time  $t$ . The current price of stock  $S$  and the cost of the call option  $C$ .

**55.5.** European-style put option: option of selling/putting for the stock at a specified price  $K$  (known as the *exercise* or *strike* price) at the expiration time  $t$ . The cost of the call option  $P$ .

**55.6.** American-style option: allows the buyer to exercise the option at any time up to the expiration time  $t$ .

**55.7.** Putcall option parity formula:

$$S + P - C = Ke^{-rt}$$

No-arbitrage pricing:

Investment 1: buying one share stock, buying one share put option, and selling one share call option.  
Costs  $S + P - C$ .

Investment 2: putting  $Ke^{-rt}$  money in bank. Costs:  $Ke^{-rt}$ . No-arbitrage pricing:  $S + P - C = Ke^{-rt}$ .

**55.8.** American-style call option price = European-style call option price.

**55.9.** Forward contract  $F$  on stock: agrees at time 0 to pay the amount  $F$  at time  $t$  for one share of the stock at time  $t$  with current value  $S$ .

$$F = Se^{rt}$$

No-arbitrage pricing:

Investment 1: Put  $Fe^{-rt}$  in the bank and purchase a forward contract. Costs  $Fe^{-rt}$ .

Investment 2: Buy the stock. Costs:  $S$ .

No-arbitrage pricing:  $Fe^{-rt} = S$ .

**55.10.** Forward contract  $F$  on currency: agrees at time 0 to pay the amount  $F$  at time  $t$  for one unit of foreign currency at time  $t$  with the current exchange rate  $S$ . The interest rate in the foreign country is  $r_f$ , and the interest rate in (the) home country is  $r_h$ .

$$F = Se^{r_h - r_f t}$$

No-arbitrage pricing:

Investment 1: Put  $Fe^{-r_h t}$  in a home bank and purchase a forward contract. Costs:  $Fe^{-r_h t}$ .

Investment 2: Buy  $e^{-r_f t}$  units of foreign currency and put them in a foreign bank. Costs:  $Se^{-r_f t}$ .

No-arbitrage pricing:  $Fe^{-r_h t} = Se^{-r_f t}$ .

# 56 ARBITRAGE THEOREM

1. Let  $i = 1, 2, \dots, n$  be  $n$  investments,
2. Let  $j = 1, 2, \dots, m$  be  $m$  outcomes,
3. Let  $r_i(j)$  be the return from investment  $i$  when outcome  $j$  occurs,
4. Let  $p_j$  be the probability of outcome  $j$ ,
5. Let  $x_i$  be the allocation (wager) on investment  $i$ ,
6. Let  $r$  be the risk-free return.

*The arbitrage theorem:* Exactly one of the following is true:

1. There exists a probability vector  $p = (p_1, \dots, p_m)$  for which for every  $i$

$$\sum_{j=1}^m p_j r_i(j) = 0$$

This probability of existence is called the *risk-neutral probability*.

2. There exists an allocation  $x = (x_1, \dots, x_n)$  for which for every  $j$

$$\sum_{i=1}^n x_i r_i(j) > r$$

# 57

## BLACK-SCHOLES FORMULA

- Risk-neutral geometric Brownian motion with drift parameter  $\mu_m = r - \sigma^2/2$  and variance parameter  $\sigma^2$ :

$$S(t) = e^{X(t)},$$

where  $X(t)$  is a Brownian motion with drift parameter  $\mu_m = r - \sigma^2/2$  and variance parameter  $\sigma^2$ .

- Under the risk-neutral geometric Brownian motion,  $S(t)/S(0)$  is a log-normal random variable with mean parameter  $(r - \sigma^2/2)t$  and variance parameter  $\sigma^2 t$ .
- No-arbitrage cost of a call option to purchase the security at time  $t$  for the specified price  $K$  is

$$C = e^{-rt} E[(S(t) - K)_+] = e^{-rt} E[(S(0)e^W - K)_+]$$

where  $W$  is a normal random variable with mean  $(r - \sigma^2/2)t$  and variance  $\sigma^2 t$ .

- Black-Scholes option pricing formula:

$$\begin{aligned} C(S, t, K, \sigma, r) &= S\Phi(\omega) - Ke^{-rt}\Phi(\omega - \sigma\sqrt{t}) \\ \omega &= \frac{rt + \sigma^2/2 - \log(K/S)}{\sigma\sqrt{t}} \end{aligned}$$

where  $\Phi(x)$  is the standard normal distribution function.


- European put option formula:

$$P(S, t, K, \sigma, r) = C(S, t, K, \sigma, r) + Ke^{-rt} - S$$

- Risk-neutral multi-period binomial stock model:
- Risk-neutral probability

$$p_m = \frac{1}{2} \left( 1 + \frac{r - \sigma^2/2}{\sigma} \sqrt{\Delta} \right), \quad u = e^{\sigma\sqrt{\Delta}}, \quad d = e^{-\sigma\sqrt{\Delta}}$$


The price  $S(i)$  of a stock at the end of period  $i$ ,  $i = 1, 2, \dots, n$ . The period is  $\Delta$ , and hence  $t_i = i\Delta$ . The initial price  $S(0)$  is non-random and fixed. The stock price  $S(i)$  is random, going up or down probabilistic rules:



- Properties of the Black-Scholes option cost:
  1.  $C$  is increasing in  $S$ ,  $t$ ,  $\sigma$ , and  $r$ .
  2.  $C$  is decreasing in  $K$ .
  3.  $C$  is a convex function of  $S$  and  $K$ .

# 58 THE DELTA HEDGING ARBITRAGE STRATEGY


- A general option for a one-period binomial stock model with payoff:



The interest rate for the period is  $r$ . The price of the option is


$$x = (pa + (1-p)b)/(1+r), \quad p = \frac{1+r-d}{u-d}.$$

- Replicate the option: invest  $y$  shares of stock, and deposit  $x - ys$  money in a bank



Solve for  $y = \frac{a-b}{(u-d)s}$ .

- Consider an option for a two-period binomial stock model:



- Payoff  $x_{2,2}, x_{1,2}, x_{0,2}$  at the end of period two  $t = 2\Delta$  when the stock prices reach the value  $u^2s$ ,  $uds$ , and  $d^2s$ , respectively.
- The value of the option  $x_{1,1}, x_{0,1}$  at  $t = \Delta$  when the stock prices reach the value  $us$  and  $ds$  respectively.

$$x_{1,1} = (px_{2,2} + (1-p)x_{1,2})/(1+r)$$

$$x_{0,1} = (px_{1,2} + (1-p)x_{0,2})/(1+r)$$

The value of the option  $x_0$  at  $t = 0$

$$x_{0,0} = (px_{1,1} + (1-p)x_{0,1})/(1+r) = (p^2x_{2,2} + 2p(1-p)x_{1,2} + (1-p)^2x_{0,2})/(1+r)^2$$

- Replicate the option through holding  $y_{0,0}$  shares of stock at  $t = 0$  and holding  $y_{1,1}, y_{0,1}$  shares of stock at  $t = \Delta$  when the stock prices reach the value  $us, ds$ , respectively. The rest of the money is deposited in a bank.

At time 0: holding  $y_{0,0}$  shares of stock:

$$y_{0,0} = \frac{x_{1,1} - x_{0,1}}{(u-d)s}$$

At time 1 and the stock is up: holding  $y_{1,1}$  shares of stock

$$y_{1,1} = \frac{x_{2,2} - x_{1,2}}{(u-d)us}$$

At time 1 and the stock is down: holding  $y_{0,1}$  shares of stock

$$y_{0,1} = \frac{x_{1,2} - x_{0,2}}{(u-d)ds}$$

This investment always has the same value as that of the option. The investor is risk-free.

- General delta hedging arbitrage strategy:

Invest capital of  $C(S,t,K,\sigma,r)$  in the option and then calls for owning exactly  $\frac{\partial C}{\partial S}$  shares of the security when its current price is  $S$  and time  $t$  remains before the option expires.

$\frac{\partial C}{\partial S}$  is called *delta*.

$$\frac{\partial C}{\partial S} = \Phi(\omega)$$

- $\frac{\partial C}{\partial r}$  is called *rho*.

$$\frac{\partial C}{\partial r} = Kte^{-rt}\Phi(\omega - \sigma\sqrt{t})$$

- $\frac{\partial C}{\partial \sigma}$  is called *vega*.

$$\frac{\partial C}{\partial \sigma} = S\sqrt{t}\Phi'(\omega)$$

- $\frac{\partial C}{\partial t}$  is called *theta*.

$$\frac{\partial C}{\partial t} = \frac{\sigma}{2\sqrt{t}}S\Phi'(\omega) + Kre^{-rt}\Phi(\omega - \sigma\sqrt{t})$$

**PART B**

# **TABLES**

## Section I: Logarithmic, Trigonometric, Exponential Functions

# 1

### FOUR PLACE COMMON LOGARITHMS

$\log_{10} N$  or  $\log N$

N	N					Proportional Parts													
	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9
10	0000	0043	0086	0128	0170	0212	0253	0294	0334	0374	4	8	12	17	21	25	29	33	37
11	0414	0453	0492	0531	0569	0607	0645	0682	0719	0755	4	8	11	15	19	23	26	30	34
12	0792	0828	0864	0899	0934	0969	1004	1038	1072	1106	3	7	10	14	17	21	24	28	31
13	1139	1173	1206	1239	1271	1303	1335	1367	1399	1430	3	6	10	13	16	19	23	26	29
14	1461	1492	1523	1553	1584	1614	1644	1673	1703	1732	3	6	9	12	15	18	21	24	27
15	1761	1790	1818	1847	1875	1903	1931	1959	1987	2014	3	6	8	11	14	17	20	22	25
16	2041	2068	2095	2122	2148	2175	2201	2227	2253	2279	3	5	8	11	13	16	18	21	24
17	2304	2330	2355	2380	2405	2430	2455	2480	2504	2529	2	5	7	10	12	15	17	20	22
18	2553	2577	2601	2625	2648	2672	2695	2718	2742	2765	2	5	7	9	12	14	16	19	21
19	2788	2810	2833	2856	2878	2900	2923	2945	2967	2989	2	4	7	9	11	13	16	18	20
20	3010	3032	3054	3075	3096	3118	3139	3160	3181	3201	2	4	6	8	11	13	15	17	19
21	3222	3243	3263	3284	3304	3324	3345	3365	3385	3404	2	4	6	8	10	12	14	16	18
22	3424	3444	3464	3483	3502	3522	3541	3560	3579	3598	2	4	6	8	10	12	14	15	17
23	3617	3636	3655	3674	3692	3711	3729	3747	3766	3784	2	4	6	7	9	11	13	15	17
24	3802	3820	3838	3856	3874	3892	3909	3927	3945	3962	2	4	5	7	9	11	12	14	16
25	3979	3997	4014	4031	4048	4065	4082	4099	4116	4133	2	3	5	7	9	10	12	14	15
26	4150	4166	4183	4200	4216	4232	4249	4265	4281	4298	2	3	5	7	8	10	11	13	15
27	4314	4330	4346	4362	4378	4393	4409	4425	4440	4456	2	3	5	6	8	9	11	13	14
28	4472	4487	4502	4518	4533	4548	4564	4579	4594	4609	2	3	5	6	8	9	11	12	14
29	4624	4639	4654	4669	4683	4698	4713	4728	4742	4757	1	3	4	6	7	9	10	12	13
30	4771	4786	4800	4814	4829	4843	4857	4871	4886	4900	1	3	4	6	7	9	10	11	13
31	4914	4928	4942	4955	4969	4983	4997	5011	5024	5038	1	3	4	6	7	8	10	11	12
32	5051	5065	5079	5092	5105	5119	5132	5145	5159	5172	1	3	4	5	7	8	9	11	12
33	5185	5198	5211	5224	5237	5250	5263	5276	5289	5302	1	3	4	5	6	8	9	10	12
34	5315	5328	5340	5353	5366	5378	5391	5403	5416	5428	1	3	4	5	6	8	9	10	11
35	5441	5453	5465	5478	5490	5502	5514	5527	5539	5551	1	2	4	5	6	7	9	10	11
36	5563	5575	5587	5599	5611	5623	5635	5647	5658	5670	1	2	4	5	6	7	8	10	11
37	5682	5694	5705	5717	5729	5740	5752	5763	5775	5786	1	2	3	5	6	7	8	9	10
38	5798	5809	5821	5832	5843	5855	5866	5877	5888	5899	1	2	3	5	6	7	8	9	10
39	5911	5922	5933	5944	5955	5966	5977	5988	5999	6010	1	2	3	4	5	7	8	9	10
40	6021	6031	6042	6053	6064	6075	6085	6096	6107	6117	1	2	3	4	5	6	8	9	10
41	6128	6138	6149	6160	6170	6180	6191	6201	6212	6222	1	2	3	4	5	6	7	8	9
42	6232	6243	6253	6263	6274	6284	6294	6304	6314	6325	1	2	3	4	5	6	7	8	9
43	6335	6345	6355	6365	6375	6385	6395	6405	6415	6425	1	2	3	4	5	6	7	8	9
44	6435	6444	6454	6464	6474	6484	6493	6503	6513	6522	1	2	3	4	5	6	7	8	9
45	6532	6542	6551	6561	6571	6580	6590	6599	6609	6618	1	2	3	4	5	6	7	8	9
46	6628	6637	6646	6656	6665	6675	6684	6693	6702	6712	1	2	3	4	5	6	7	7	8
47	6721	6730	6739	6749	6758	6767	6776	6785	6794	6803	1	2	3	4	5	5	6	7	8
48	6812	6821	6830	6839	6848	6857	6866	6875	6884	6893	1	2	3	4	4	5	6	7	8
49	6902	6911	6920	6928	6937	6946	6955	6964	6972	6981	1	2	3	4	4	5	6	7	8
50	6990	6998	7007	7016	7024	7033	7042	7050	7059	7067	1	2	3	3	4	5	6	7	8
51	7076	7084	7093	7101	7110	7118	7126	7135	7143	7152	1	2	3	3	4	5	6	7	8
52	7160	7168	7177	7185	7193	7202	7210	7218	7226	7235	1	2	2	3	4	5	6	7	7
53	7243	7251	7259	7267	7275	7284	7292	7300	7308	7316	1	2	2	3	4	5	6	6	7
54	7324	7332	7340	7348	7356	7364	7372	7380	7388	7396	1	2	2	3	4	5	6	6	7

N 0 1 2 3 4 5 6 7 8 9

## 1

## FOUR PLACE COMMON LOGARITHMS

 $\log_{10} N$  or  $\log N$  (*Continued*)

N	0 1 2 3 4					5 6 7 8 9					Proportional Parts								
	1	2	3	4	5	6	7	8	9	1	2	3	4	5	6	7	8	9	
55	7404	7412	7419	7427	7435	7443	7451	7459	7466	7474	1	2	2	3	4	5	5	6	7
56	7482	7490	7497	7505	7513	7520	7528	7536	7543	7551	1	2	2	3	4	5	5	6	7
57	7559	7566	7574	7582	7589	7597	7604	7612	7619	7627	1	2	2	3	4	5	5	6	7
58	7634	7642	7649	7657	7664	7672	7679	7686	7694	7701	1	1	2	3	4	4	5	6	7
59	7709	7716	7723	7731	7738	7745	7752	7760	7767	7774	1	1	2	3	4	4	5	6	7
60	7782	7789	7796	7803	7810	7818	7825	7832	7839	7846	1	1	2	3	4	4	5	6	6
61	7853	7860	7868	7875	7882	7889	7896	7903	7910	7917	1	1	2	3	4	4	5	6	6
62	7924	7931	7938	7945	7952	7959	7966	7973	7980	7987	1	1	2	3	3	4	5	6	6
63	7993	8000	8007	8014	8021	8028	8035	8041	8048	8055	1	1	2	3	3	4	5	5	6
64	8062	8069	8075	8082	8089	8096	8102	8109	8116	8122	1	1	2	3	3	4	5	5	6
65	8129	8136	8142	8149	8156	8162	8169	8176	8182	8189	1	1	2	3	3	4	5	5	6
66	8195	8202	8209	8215	8222	8228	8235	8241	8248	8254	1	1	2	3	3	4	5	5	6
67	8261	8267	8274	8280	8287	8293	8299	8306	8312	8319	1	1	2	3	3	4	5	5	6
68	8325	8331	8338	8344	8351	8357	8363	8370	8376	8382	1	1	2	3	3	4	4	5	6
69	8388	8395	8401	8407	8414	8420	8426	8432	8439	8445	1	1	2	2	3	4	4	5	6
70	8451	8457	8463	8470	8476	8482	8488	8494	8500	8506	1	1	2	2	3	4	4	5	6
71	8513	8519	8525	8531	8537	8543	8549	8555	8561	8567	1	1	2	2	3	4	4	5	5
72	8573	8579	8585	8591	8597	8603	8609	8615	8621	8627	1	1	2	2	3	4	4	5	5
73	8633	8639	8645	8651	8657	8663	8669	8675	8681	8686	1	1	2	2	3	4	4	5	5
74	8692	8698	8704	8710	8716	8722	8727	8733	8739	8745	1	1	2	2	3	4	4	5	5
75	8751	8756	8762	8768	8774	8779	8785	8791	8797	8802	1	1	2	2	3	3	4	5	5
76	8808	8814	8820	8825	8831	8837	8842	8848	8854	8859	1	1	2	2	3	3	4	5	5
77	8865	8871	8876	8882	8887	8893	8899	8904	8910	8915	1	1	2	2	3	3	4	4	5
78	8921	8927	8932	8938	8943	8949	8954	8960	8965	8971	1	1	2	2	3	3	4	4	5
79	8976	8982	8987	8993	8998	9004	9009	9015	9020	9025	1	1	2	2	3	3	4	4	5
80	9031	9036	9042	9047	9053	9058	9063	9069	9074	9079	1	1	2	2	3	3	4	4	5
81	9085	9090	9096	9101	9106	9112	9117	9122	9128	9133	1	1	2	2	3	3	4	4	5
82	9138	9143	9149	9154	9159	9165	9170	9175	9180	9186	1	1	2	2	3	3	4	4	5
83	9191	9196	9201	9206	9212	9217	9222	9227	9232	9238	1	1	2	2	3	3	4	4	5
84	9243	9248	9253	9258	9263	9269	9274	9279	9284	9289	1	1	2	2	3	3	4	4	5
85	9294	9299	9304	9309	9315	9320	9325	9330	9335	9340	1	1	2	2	3	3	4	4	5
86	9345	9350	9355	9360	9365	9370	9375	9380	9385	9390	1	1	2	2	3	3	4	4	5
87	9395	9400	9405	9410	9415	9420	9425	9430	9435	9440	0	1	1	2	2	3	3	4	4
88	9445	9450	9455	9460	9465	9469	9474	9479	9484	9489	0	1	1	2	2	3	3	4	4
89	9494	9499	9504	9509	9513	9518	9523	9528	9533	9538	0	1	1	2	2	3	3	4	4
90	9542	9547	9552	9557	9562	9566	9571	9576	9581	9586	0	1	1	2	2	3	3	4	4
91	9590	9595	9600	9605	9609	9614	9619	9624	9628	9633	0	1	1	2	2	3	3	4	4
92	9638	9643	9647	9652	9657	9661	9666	9671	9675	9680	0	1	1	2	2	3	3	4	4
93	9685	9689	9694	9699	9703	9708	9713	9717	9722	9727	0	1	1	2	2	3	3	4	4
94	9731	9736	9741	9745	9750	9754	9759	9763	9768	9773	0	1	1	2	2	3	3	4	4
95	9777	9782	9786	9791	9795	9800	9805	9809	9814	9818	0	1	1	2	2	3	3	4	4
96	9823	9827	9832	9836	9841	9845	9850	9854	9859	9863	0	1	1	2	2	3	3	4	4
97	9868	9872	9877	9881	9886	9890	9894	9899	9903	9908	0	1	1	2	2	3	3	4	4
98	9912	9917	9921	9926	9930	9934	9939	9943	9948	9952	0	1	1	2	2	3	3	4	4
99	9956	9961	9965	9969	9974	9978	9983	9987	9991	9996	0	1	1	2	2	3	3	3	4

## 2

**Sin x**  
(x in degrees and minutes)

<i>x</i>	0'	10'	20'	30'	40'	50'
0°	.0000	.0029	.0058	.0087	.0116	.0145
1	.0175	.0204	.0233	.0262	.0291	.0320
2	.0349	.0378	.0407	.0436	.0465	.0494
3	.0523	.0552	.0581	.0610	.0640	.0669
4	.0698	.0727	.0756	.0785	.0814	.0843
5°	.0872	.0901	.0929	.0958	.0987	.1016
6	.1045	.1074	.1103	.1132	.1161	.1190
7	.1219	.1248	.1276	.1305	.1334	.1363
8	.1392	.1421	.1449	.1478	.1507	.1536
9	.1564	.1593	.1622	.1650	.1679	.1708
10°	.1736	.1765	.1794	.1822	.1851	.1880
11	.1908	.1937	.1965	.1994	.2022	.2051
12	.2079	.2108	.2136	.2164	.2193	.2221
13	.2250	.2278	.2306	.2334	.2363	.2391
14	.2419	.2447	.2476	.2504	.2532	.2560
15°	.2588	.2616	.2644	.2672	.2700	.2728
16	.2756	.2784	.2812	.2840	.2868	.2896
17	.2924	.2952	.2979	.3007	.3035	.3062
18	.3090	.3118	.3145	.3173	.3201	.3228
19	.3256	.3283	.3311	.3338	.3365	.3393
20°	.3420	.3448	.3475	.3502	.3529	.3557
21	.3584	.3611	.3638	.3665	.3692	.3719
22	.3746	.3773	.3800	.3827	.3854	.3881
23	.3907	.3934	.3961	.3987	.4014	.4041
24	.4067	.4094	.4120	.4147	.4173	.4200
25°	.4226	.4253	.4279	.4305	.4331	.4358
26	.4384	.4410	.4436	.4462	.4488	.4514
27	.4540	.4566	.4592	.4617	.4643	.4669
28	.4695	.4720	.4746	.4772	.4797	.4823
29	.4848	.4874	.4899	.4924	.4950	.4975
30°	.5000	.5025	.5050	.5075	.5100	.5125
31	.5150	.5175	.5200	.5225	.5250	.5275
32	.5299	.5324	.5348	.5373	.5398	.5422
33	.5446	.5471	.5495	.5519	.5544	.5568
34	.5592	.5616	.5640	.5664	.5688	.5712
35°	.5736	.5760	.5783	.5807	.5831	.5854
36	.5878	.5901	.5925	.5948	.5972	.5995
37	.6018	.6041	.6065	.6088	.6111	.6134
38	.6157	.6180	.6202	.6225	.6248	.6271
39	.6293	.6316	.6338	.6361	.6383	.6406
40°	.6428	.6450	.6472	.6494	.6517	.6539
41	.6561	.6583	.6604	.6626	.6648	.6670
42	.6691	.6713	.6734	.6756	.6777	.6799
43	.6820	.6841	.6862	.6884	.6905	.6926
44	.6947	.6967	.6988	.7009	.7030	.7050
45°	.7071	.7092	.7112	.7133	.7153	.7173

<i>x</i>	0'	10'	20'	30'	40'	50'
45°	.7071	.7092	.7112	.7133	.7153	.7173
46	.7193	.7214	.7234	.7254	.7274	.7294
47	.7314	.7333	.7353	.7373	.7392	.7412
48	.7431	.7451	.7470	.7490	.7509	.7528
49	.7547	.7566	.7585	.7604	.7623	.7642
50°	.7660	.7679	.7698	.7716	.7735	.7753
51	.7771	.7790	.7808	.7826	.7844	.7862
52	.7880	.7898	.7916	.7934	.7951	.7969
53	.7986	.8004	.8021	.8039	.8056	.8073
54	.8090	.8107	.8124	.8141	.8158	.8175
55°	.8192	.8208	.8225	.8241	.8258	.8274
56	.8290	.8307	.8323	.8339	.8355	.8371
57	.8387	.8403	.8418	.8434	.8450	.8465
58	.8480	.8496	.8511	.8526	.8542	.8557
59	.8572	.8587	.8601	.8616	.8631	.8646
60°	.8660	.8675	.8689	.8704	.8718	.8732
61	.8746	.8760	.8774	.8788	.8802	.8816
62	.8829	.8843	.8857	.8870	.8884	.8897
63	.8910	.8923	.8936	.8949	.8962	.8975
64	.8988	.9001	.9013	.9026	.9038	.9051
65°	.9063	.9075	.9088	.9100	.9112	.9124
66	.9135	.9147	.9159	.9171	.9182	.9194
67	.9205	.9216	.9228	.9239	.9250	.9261
68	.9272	.9283	.9293	.9304	.9315	.9325
69	.9336	.9346	.9356	.9367	.9377	.9387
70°	.9397	.9407	.9417	.9426	.9436	.9446
71	.9455	.9465	.9474	.9483	.9492	.9502
72	.9511	.9520	.9528	.9537	.9546	.9555
73	.9563	.9572	.9580	.9588	.9596	.9605
74	.9613	.9621	.9628	.9636	.9644	.9652
75°	.9659	.9667	.9674	.9681	.9689	.9696
76	.9703	.9710	.9717	.9724	.9730	.9737
77	.9744	.9750	.9757	.9763	.9769	.9775
78	.9781	.9787	.9793	.9799	.9805	.9811
79	.9816	.9822	.9827	.9833	.9838	.9843
80°	.9848	.9853	.9858	.9863	.9868	.9872
81	.9877	.9881	.9886	.9890	.9894	.9899
82	.9903	.9907	.9911	.9914	.9918	.9922
83	.9925	.9929	.9932	.9936	.9939	.9942
84	.9945	.9948	.9951	.9954	.9957	.9959
85°	.9962	.9964	.9967	.9969	.9971	.9974
86	.9976	.9978	.9980	.9981	.9983	.9985
87	.9986	.9988	.9989	.9990	.9992	.9993
88	.9994	.9995	.9996	.9997	.9998	
89	.9998	.9999	.9999	1.0000	1.0000	
90°	1.0000					

# 3

**Cos x**  
(x in degrees and minutes)

<i>x</i>	0'	10'	20'	30'	40'	50'
0°	1.0000	1.0000	1.0000	1.0000	.9999	.9999
1	.9998	.9998	.9997	.9997	.9996	.9995
2	.9994	.9993	.9992	.9990	.9989	.9988
3	.9986	.9985	.9983	.9981	.9980	.9978
4	.9976	.9974	.9971	.9969	.9967	.9964
5°	.9962	.9959	.9957	.9954	.9951	.9948
6	.9945	.9942	.9939	.9936	.9932	.9929
7	.9925	.9922	.9918	.9914	.9911	.9907
8	.9903	.9899	.9894	.9890	.9886	.9881
9	.9877	.9872	.9868	.9863	.9858	.9853
10°	.9848	.9843	.9838	.9833	.9827	.9822
11	.9816	.9811	.9805	.9799	.9793	.9787
12	.9781	.9775	.9769	.9763	.9757	.9750
13	.9744	.9737	.9730	.9724	.9717	.9710
14	.9703	.9696	.9689	.9681	.9674	.9667
15°	.9659	.9652	.9644	.9636	.9628	.9621
16	.9613	.9605	.9596	.9588	.9580	.9572
17	.9563	.9555	.9546	.9537	.9528	.9520
18	.9511	.9502	.9492	.9483	.9474	.9465
19	.9455	.9446	.9436	.9426	.9417	.9407
20°	.9397	.9387	.9377	.9367	.9356	.9346
21	.9336	.9325	.9315	.9304	.9293	.9283
22	.9272	.9261	.9250	.9239	.9228	.9216
23	.9205	.9194	.9182	.9171	.9159	.9147
24	.9135	.9124	.9112	.9100	.9088	.9075
25°	.9063	.9051	.9038	.9026	.9013	.9001
26	.8988	.8975	.8962	.8949	.8936	.8923
27	.8910	.8897	.8884	.8870	.8857	.8843
28	.8829	.8816	.8802	.8788	.8774	.8760
29	.8746	.8732	.8718	.8704	.8689	.8675
30°	.8660	.8646	.8631	.8616	.8601	.8587
31	.8572	.8557	.8542	.8526	.8511	.8496
32	.8480	.8465	.8450	.8434	.8418	.8403
33	.8387	.8371	.8355	.8339	.8323	.8307
34	.8290	.8274	.8258	.8241	.8225	.8208
35°	.8192	.8175	.8158	.8141	.8124	.8107
36	.8090	.8073	.8056	.8039	.8021	.8004
37	.7986	.7969	.7951	.7934	.7916	.7898
38	.7880	.7862	.7844	.7826	.7808	.7790
39	.7771	.7753	.7735	.7716	.7698	.7679
40°	.7660	.7642	.7623	.7604	.7585	.7566
41	.7547	.7528	.7509	.7490	.7470	.7451
42	.7431	.7412	.7392	.7373	.7353	.7333
43	.7314	.7294	.7274	.7254	.7234	.7214
44	.7193	.7173	.7153	.7133	.7112	.7092
45°	.7071	.7050	.7030	.7009	.6988	.6967

<i>x</i>	0'	10'	20'	30'	40'	50'
45°	.7071	.7050	.7030	.7009	.6988	.6967
46	.6947	.6926	.6905	.6884	.6862	.6841
47	.6820	.6799	.6777	.6756	.6734	.6713
48	.6691	.6670	.6648	.6626	.6604	.6583
49	.6561	.6539	.6517	.6494	.6472	.6450
50°	.6428	.6406	.6383	.6361	.6338	.6316
51	.6293	.6271	.6248	.6225	.6202	.6180
52	.6157	.6134	.6111	.6088	.6065	.6041
53	.6018	.5995	.5972	.5948	.5925	.5901
54	.5878	.5854	.5831	.5807	.5783	.5760
55°	.5736	.5712	.5688	.5664	.5640	.5616
56	.5592	.5568	.5544	.5519	.5495	.5471
57	.5446	.5422	.5398	.5373	.5348	.5324
58	.5299	.5275	.5250	.5225	.5200	.5175
59	.5150	.5125	.5100	.5075	.5050	.5025
60°	.5000	.4975	.4950	.4924	.4899	.4874
61	.4848	.4823	.4797	.4772	.4746	.4720
62	.4695	.4669	.4643	.4617	.4592	.4566
63	.4540	.4514	.4488	.4462	.4436	.4410
64	.4384	.4358	.4331	.4305	.4279	.4253
65°	.4226	.4200	.4173	.4147	.4120	.4094
66	.4067	.4041	.4014	.3987	.3961	.3934
67	.3907	.3881	.3854	.3827	.3800	.3773
68	.3746	.3719	.3692	.3665	.3638	.3611
69	.3584	.3557	.3529	.3502	.3475	.3448
70°	.3420	.3393	.3365	.3338	.3311	.3283
71	.3256	.3228	.3201	.3173	.3145	.3118
72	.3090	.3062	.3035	.3007	.2979	.2952
73	.2924	.2896	.2868	.2840	.2812	.2784
74	.2756	.2728	.2700	.2672	.2644	.2616
75°	.2588	.2560	.2532	.2504	.2476	.2447
76	.2419	.2391	.2363	.2334	.2306	.2278
77	.2250	.2221	.2193	.2164	.2136	.2108
78	.2079	.2051	.2022	.1994	.1965	.1937
79	.1908	.1880	.1851	.1822	.1794	.1765
80°	.1736	.1708	.1679	.1650	.1622	.1593
81	.1564	.1536	.1507	.1478	.1449	.1421
82	.1392	.1363	.1334	.1305	.1276	.1248
83	.1219	.1190	.1161	.1132	.1103	.1074
84	.1045	.1016	.0987	.0958	.0929	.0901
85°	.0872	.0843	.0814	.0785	.0756	.0727
86	.0698	.0669	.0640	.0610	.0581	.0552
87	.0523	.0494	.0465	.0436	.0407	.0378
88	.0349	.0320	.0291	.0262	.0233	.0204
89	.0175	.0145	.0116	.0087	.0058	.0029
90°	.0000					

## 4

**Tan x**  
(x in degrees and minutes)

$x$	0'	10'	20'	30'	40'	50'
0°	.0000	.0029	.0058	.0087	.0116	.0145
1	.0175	.0204	.0233	.0262	.0291	.0320
2	.0349	.0378	.0407	.0437	.0466	.0495
3	.0524	.0553	.0582	.0612	.0641	.0670
4	.0699	.0729	.0758	.0787	.0816	.0846
5°	.0875	.0904	.0934	.0963	.0992	.1022
6	.1051	.1080	.1110	.1139	.1169	.1198
7	.1228	.1257	.1287	.1317	.1346	.1376
8	.1405	.1435	.1465	.1495	.1524	.1554
9	.1584	.1614	.1644	.1673	.1703	.1733
10°	.1763	.1793	.1823	.1853	.1883	.1914
11	.1944	.1974	.2004	.2035	.2065	.2095
12	.2126	.2156	.2186	.2217	.2247	.2278
13	.2309	.2339	.2370	.2401	.2432	.2462
14	.2493	.2524	.2555	.2586	.2617	.2648
15°	.2679	.2711	.2742	.2773	.2805	.2836
16	.2867	.2899	.2931	.2962	.2994	.3026
17	.3057	.3089	.3121	.3153	.3185	.3217
18	.3249	.3281	.3314	.3346	.3378	.3411
19	.3443	.3476	.3508	.3541	.3574	.3607
20°	.3640	.3673	.3706	.3739	.3772	.3805
21	.3839	.3872	.3906	.3939	.3973	.4006
22	.4040	.4074	.4108	.4142	.4176	.4210
23	.4245	.4279	.4314	.4348	.4383	.4417
24	.4452	.4487	.4522	.4557	.4592	.4628
25°	.4663	.4699	.4734	.4770	.4806	.4841
26	.4877	.4913	.4950	.4986	.5022	.5059
27	.5095	.5132	.5169	.5206	.5243	.5280
28	.5317	.5354	.5392	.5430	.5467	.5505
29	.5543	.5581	.5619	.5658	.5696	.5735
30°	.5774	.5812	.5851	.5890	.5930	.5969
31	.6009	.6048	.6088	.6128	.6168	.6208
32	.6249	.6289	.6330	.6371	.6412	.6453
33	.6494	.6536	.6577	.6619	.6661	.6703
34	.6745	.6787	.6830	.6873	.6916	.6959
35°	.7002	.7046	.7089	.7133	.7177	.7221
36	.7265	.7310	.7355	.7400	.7445	.7490
37	.7536	.7581	.7627	.7673	.7720	.7766
38	.7813	.7860	.7907	.7954	.8002	.8050
39	.8098	.8146	.8195	.8243	.8292	.8342
40°	.8391	.8441	.8491	.8541	.8591	.8642
41	.8693	.8744	.8796	.8847	.8899	.8952
42	.9004	.9057	.9110	.9163	.9217	.9271
43	.9325	.9380	.9435	.9490	.9545	.9601
44	.9657	.9713	.9770	.9827	.9884	.9942
45°	1.0000	1.0058	1.0117	1.0176	1.0235	1.0295

$x$	0'	10'	20'	30'	40'	50'
45°	1.0000	1.0058	1.0117	1.0176	1.0235	1.0295
46	1.0355	1.0416	1.0477	1.0538	1.0599	1.0661
47	1.0724	1.0786	1.0850	1.0913	1.0977	1.1041
48	1.1106	1.1171	1.1237	1.1303	1.1369	1.1436
49	1.1504	1.1571	1.1640	1.1708	1.1778	1.1847
50°	1.1918	1.1988	1.2059	1.2131	1.2203	1.2276
51	1.2349	1.2423	1.2497	1.2572	1.2647	1.2723
52	1.2799	1.2876	1.2954	1.3032	1.3111	1.3190
53	1.3270	1.3351	1.3432	1.3514	1.3597	1.3680
54	1.3764	1.3848	1.3934	1.4019	1.4106	1.4193
55°	1.4281	1.4370	1.4460	1.4550	1.4641	1.4733
56	1.4826	1.4919	1.5013	1.5108	1.5204	1.5301
57	1.5399	1.5497	1.5597	1.5697	1.5798	1.5900
58	1.6003	1.6107	1.6212	1.6319	1.6426	1.6534
59	1.6643	1.6753	1.6864	1.6977	1.7090	1.7205
60°	1.7321	1.7437	1.7556	1.7675	1.7796	1.7917
61	1.8040	1.8165	1.8291	1.8418	1.8546	1.8676
62	1.8807	1.8940	1.9074	1.9210	1.9347	1.9486
63	1.9626	1.9768	1.9912	2.0057	2.0204	2.0353
64	2.0503	2.0655	2.0809	2.0965	2.1123	2.1283
65°	2.1445	2.1609	2.1775	2.1943	2.2113	2.2286
66	2.2460	2.2637	2.2817	2.2998	2.3183	2.3369
67	2.3559	2.3750	2.3945	2.4142	2.4342	2.4545
68	2.4751	2.4960	2.5172	2.5386	2.5605	2.5826
69	2.6051	2.6279	2.6511	2.6746	2.6985	2.7228
70°	2.7475	2.7725	2.7980	2.8239	2.8502	2.8770
71	2.9042	2.9319	2.9600	2.9887	3.0178	3.0475
72	3.0777	3.1084	3.1397	3.1716	3.2041	3.2371
73	3.2709	3.3052	3.3402	3.3759	3.4124	3.4495
74	3.4874	3.5261	3.5656	3.6059	3.6470	3.6891
75°	3.7321	3.7760	3.8208	3.8667	3.9136	3.9617
76	4.0108	4.0611	4.1126	4.1653	4.2193	4.2747
77	4.3315	4.3897	4.4494	4.5107	4.5736	4.6382
78	4.7046	4.7729	4.8430	4.9152	4.9894	5.0658
79	5.1446	5.2257	5.3093	5.3955	5.4845	5.5764
80°	5.6713	5.7694	5.8708	5.9758	6.0844	6.1970
81	6.3138	6.4348	6.5606	6.6912	6.8269	6.9682
82	7.1154	7.2687	7.4287	7.5958	7.7704	7.9530
83	8.1443	8.3450	8.5555	8.7769	9.0098	9.2553
84	9.5144	9.7882	10.078	10.385	10.712	11.059
85°	11.430	11.826	12.251	12.706	13.197	13.727
86	14.301	14.924	15.605	16.350	17.169	18.075
87	19.081	20.206	21.470	22.904	24.542	26.432
88	28.636	31.242	34.368	38.188	42.964	49.104
89	57.290	68.750	85.940	114.59	171.89	343.77
90°	$\infty$					

# 5

## CONVERSION OF RADIANS TO DEGREES, MINUTES, AND SECONDS OR FRACTIONS OF DEGREES

Radians	Deg.	Min.	Sec.	Fractions of Degrees
1	57°	17'	44.8''	57.2958°
2	114°	35'	29.6''	114.5916°
3	171°	53'	14.4''	171.8873°
4	229°	10'	59.2''	229.1831°
5	286°	28'	44.0''	286.4789°
6	343°	46'	28.8''	343.7747°
7	401°	4'	13.6''	401.0705°
8	458°	21'	58.4''	458.3662°
9	515°	39'	43.3''	515.6620°
10	572°	57'	28.1''	572.9578°
.1	5°	43'	46.5''	
.2	11°	27'	33.0''	
.3	17°	11'	19.4''	
.4	22°	55'	5.9''	
.5	28°	38'	52.4''	
.6	34°	22'	38.9''	
.7	40°	6'	25.4''	
.8	45°	50'	11.8''	
.9	51°	33'	58.3''	
.01	0°	34'	22.6''	
.02	1°	8'	45.3''	
.03	1°	43'	7.9''	
.04	2°	17'	30.6''	
.05	2°	51'	53.2''	
.06	3°	26'	15.9''	
.07	4°	0'	38.5''	
.08	4°	35'	1.2''	
.09	5°	9'	23.8''	
.001	0°	3'	26.3''	
.002	0°	6'	52.5''	
.003	0°	10'	18.8''	
.004	0°	13'	45.1''	
.005	0°	17'	11.3''	
.006	0°	20'	37.6''	
.007	0°	24'	3.9''	
.008	0°	27'	30.1''	
.009	0°	30'	56.4''	
.0001	0°	0'	20.6''	
.0002	0°	0'	41.3''	
.0003	0°	1'	1.9''	
.0004	0°	1'	22.5''	
.0005	0°	1'	43.1''	
.0006	0°	2'	3.8''	
.0007	0°	2'	24.4''	
.0008	0°	2'	45.0''	
.0009	0°	3'	5.6''	

# 6

## CONVERSION OF DEGREES, MINUTES, AND SECONDS TO RADIANS

Degrees	Radians
1°	.0174533
2°	.0349066
3°	.0523599
4°	.0698132
5°	.0872665
6°	.1047198
7°	.1221730
8°	.1396263
9°	.1570796
10°	.1745329

Minutes	Radians
1'	.00029089
2'	.00058178
3'	.00087266
4'	.00116355
5'	.00145444
6'	.00174533
7'	.00203622
8'	.00232711
9'	.00261800
10'	.00290888

Seconds	Radians
1"	.0000048481
2"	.0000096963
3"	.0000145444
4"	.0000193925
5"	.0000242407
6"	.0000290888
7"	.0000339370
8"	.0000387851
9"	.0000436332
10"	.0000484814

## 7

## NATURAL OR NAPIERIAN LOGARITHMS

 $\log_e x$  or  $\ln x$ 

$x$	0	1	2	3	4	5	6	7	8	9
1.0	.00000	.00995	.01980	.02956	.03922	.04879	.05827	.06766	.07696	.08618
1.1	.09531	.10436	.11333	.12222	.13103	.13976	.14842	.15700	.16551	.17395
1.2	.18232	.19062	.19885	.20701	.21511	.22314	.23111	.23902	.24686	.25464
1.3	.26236	.27003	.27763	.28518	.29267	.30010	.30748	.31481	.32208	.32930
1.4	.33647	.34359	.35066	.35767	.36464	.37156	.37844	.38526	.39204	.39878
1.5	.40547	.41211	.41871	.42527	.43178	.43825	.44469	.45108	.45742	.46373
1.6	.47000	.47623	.48243	.48858	.49470	.50078	.50682	.51282	.51879	.52473
1.7	.53063	.53649	.54232	.54812	.55389	.55962	.56531	.57098	.57661	.58222
1.8	.58779	.59333	.59884	.60432	.60977	.61519	.62058	.62594	.63127	.63658
1.9	.64185	.64710	.65233	.65752	.66269	.66783	.67294	.67803	.68310	.68813
2.0	.69315	.69813	.70310	.70804	.71295	.71784	.72271	.72755	.73237	.73716
2.1	.74194	.74669	.75142	.75612	.76081	.76547	.77011	.77473	.77932	.78390
2.2	.78846	.79299	.79751	.80200	.80648	.81093	.81536	.81978	.82418	.82855
2.3	.83291	.83725	.84157	.84587	.85015	.85442	.85866	.86289	.86710	.87129
2.4	.87547	.87963	.88377	.88789	.89200	.89609	.90016	.90422	.90826	.91228
2.5	.91629	.92028	.92426	.92822	.93216	.93609	.94001	.94391	.94779	.95166
2.6	.95551	.95935	.96317	.96698	.97078	.97456	.97833	.98208	.98582	.98954
2.7	.99325	.99695	1.00063	1.00430	1.00796	1.01160	1.01523	1.01885	1.02245	1.02604
2.8	1.02962	1.03318	1.03674	1.04028	1.04380	1.04732	1.05082	1.05431	1.05779	1.06126
2.9	1.06471	1.06815	1.07158	1.07500	1.07841	1.08181	1.08519	1.08856	1.09192	1.09527
3.0	1.09861	1.10194	1.10526	1.10856	1.11186	1.11514	1.11841	1.12168	1.12493	1.12817
3.1	1.13140	1.13462	1.13783	1.14103	1.14422	1.14740	1.15057	1.15373	1.15688	1.16002
3.2	1.16315	1.16627	1.16938	1.17248	1.17557	1.17865	1.18173	1.18479	1.18784	1.19089
3.3	1.19392	1.19695	1.19996	1.20297	1.20597	1.20896	1.21194	1.21491	1.21788	1.22083
3.4	1.22378	1.22671	1.22964	1.23256	1.23547	1.23837	1.24127	1.24415	1.24703	1.24990
3.5	1.25276	1.25562	1.25846	1.26130	1.26413	1.26695	1.26976	1.27257	1.27536	1.27815
3.6	1.28093	1.28371	1.28647	1.28923	1.29198	1.29473	1.29746	1.30019	1.30291	1.30563
3.7	1.30833	1.31103	1.31372	1.31641	1.31909	1.32176	1.32442	1.32708	1.32972	1.33237
3.8	1.33500	1.33763	1.34025	1.34286	1.34547	1.34807	1.35067	1.35325	1.35584	1.35841
3.9	1.36098	1.36354	1.36609	1.36864	1.37118	1.37372	1.37624	1.37877	1.38128	1.38379
4.0	1.38629	1.38879	1.39128	1.39377	1.39624	1.39872	1.40118	1.40364	1.40610	1.40854
4.1	1.41099	1.41342	1.41585	1.41828	1.42070	1.42311	1.42552	1.42792	1.43031	1.43270
4.2	1.43508	1.43746	1.43984	1.44220	1.44456	1.44692	1.44927	1.45161	1.45395	1.45629
4.3	1.45862	1.46094	1.46326	1.46557	1.46787	1.47018	1.47247	1.47476	1.47705	1.47933
4.4	1.48160	1.48387	1.48614	1.48840	1.49065	1.49290	1.49515	1.49739	1.49962	1.50185
4.5	1.50408	1.50630	1.50851	1.51072	1.51293	1.51513	1.51732	1.51951	1.52170	1.52388
4.6	1.52606	1.52823	1.53039	1.53256	1.53471	1.53687	1.53902	1.54116	1.54330	1.54543
4.7	1.54756	1.54969	1.55181	1.55393	1.55604	1.55814	1.56025	1.56235	1.56444	1.56653
4.8	1.56862	1.57070	1.57277	1.57485	1.57691	1.57898	1.58104	1.58309	1.58515	1.58719
4.9	1.58924	1.59127	1.59331	1.59534	1.59737	1.59939	1.60141	1.60342	1.60543	1.60744

$\ln 10 = 2.30259$

$4 \ln 10 = 9.21034$

$7 \ln 10 = 16.11810$

$2 \ln 10 = 4.60517$

$5 \ln 10 = 11.51293$

$8 \ln 10 = 18.42068$

$3 \ln 10 = 6.90776$

$6 \ln 10 = 13.81551$

$9 \ln 10 = 20.72327$

## 7

## NATURAL OR NAPIERIAN LOGARITHMS

 $\log_e x$  or  $\ln x$  (*Continued*)

$x$	0	1	2	3	4	5	6	7	8	9
5.0	1.60944	1.61144	1.61343	1.61542	1.61741	1.61939	1.62137	1.62334	1.62531	1.62728
5.1	1.62924	1.63120	1.63315	1.63511	1.63705	1.63900	1.64094	1.64287	1.64481	1.64673
5.2	1.64866	1.65058	1.65250	1.65441	1.65632	1.65823	1.66013	1.66203	1.66393	1.66582
5.3	1.66771	1.66959	1.67147	1.67335	1.67523	1.67710	1.67896	1.68083	1.68269	1.68455
5.4	1.68640	1.68825	1.69010	1.69194	1.69378	1.69562	1.69745	1.69928	1.70111	1.70293
5.5	1.70475	1.70656	1.70838	1.71019	1.71199	1.71380	1.71560	1.71740	1.71919	1.72098
5.6	1.72277	1.72455	1.72633	1.72811	1.72988	1.73166	1.73342	1.73519	1.73695	1.73871
5.7	1.74047	1.74222	1.74397	1.74572	1.74746	1.74920	1.75094	1.75267	1.75440	1.75613
5.8	1.75786	1.75958	1.76130	1.76302	1.76473	1.76644	1.76815	1.76985	1.77156	1.77326
5.9	1.77495	1.77665	1.77834	1.78002	1.78171	1.78339	1.78507	1.78675	1.78842	1.79009
6.0	1.79176	1.79342	1.79509	1.79675	1.79840	1.80006	1.80171	1.80336	1.80500	1.80665
6.1	1.80829	1.80993	1.81156	1.81319	1.81482	1.81645	1.81808	1.81970	1.82132	1.82294
6.2	1.82455	1.82616	1.82777	1.82938	1.83098	1.83258	1.83418	1.83578	1.83737	1.83896
6.3	1.84055	1.84214	1.84372	1.84530	1.84688	1.84845	1.85003	1.85160	1.85317	1.85473
6.4	1.85630	1.85786	1.85942	1.86097	1.86253	1.86408	1.86563	1.86718	1.86872	1.87026
6.5	1.87180	1.87334	1.87487	1.87641	1.87794	1.87947	1.88099	1.88251	1.88403	1.88555
6.6	1.88707	1.88858	1.89010	1.89160	1.89311	1.89462	1.89612	1.89762	1.89912	1.90061
6.7	1.90211	1.90360	1.90509	1.90658	1.90806	1.90954	1.91102	1.91250	1.91398	1.91545
6.8	1.91692	1.91839	1.91986	1.92132	1.92279	1.92425	1.92571	1.92716	1.92862	1.93007
6.9	1.93152	1.93297	1.93442	1.93586	1.93730	1.93874	1.94018	1.94162	1.94305	1.94448
7.0	1.94591	1.94734	1.94876	1.95019	1.95161	1.95303	1.95445	1.95586	1.95727	1.95869
7.1	1.96009	1.96150	1.96291	1.96431	1.96571	1.96711	1.96851	1.96991	1.97130	1.97269
7.2	1.97408	1.97547	1.97685	1.97824	1.97962	1.98100	1.98238	1.98376	1.98513	1.98650
7.3	1.98787	1.98924	1.99061	1.99198	1.99334	1.99470	1.99606	1.99742	1.99877	2.00013
7.4	2.00148	2.00283	2.00418	2.00553	2.00687	2.00821	2.00956	2.01089	2.01223	2.01357
7.5	2.01490	2.01624	2.01757	2.01890	2.02022	2.02155	2.02287	2.02419	2.02551	2.02683
7.6	2.02815	2.02946	2.03078	2.03209	2.03340	2.03471	2.03601	2.03732	2.03862	2.03992
7.7	2.04122	2.04252	2.04381	2.04511	2.04640	2.04769	2.04898	2.05027	2.05156	2.05284
7.8	2.05412	2.05540	2.05668	2.05796	2.05924	2.06051	2.06179	2.06306	2.06433	2.06560
7.9	2.06686	2.06813	2.06939	2.07065	2.07191	2.07317	2.07443	2.07568	2.07694	2.07819
8.0	2.07944	2.08069	2.08194	2.08318	2.08443	2.08567	2.08691	2.08815	2.08939	2.09063
8.1	2.09186	2.09310	2.09433	2.09556	2.09679	2.09802	2.09924	2.10047	2.10169	2.10291
8.2	2.10413	2.10535	2.10657	2.10779	2.10900	2.11021	2.11142	2.11263	2.11384	2.11505
8.3	2.11626	2.11746	2.11866	2.11986	2.12106	2.12226	2.12346	2.12465	2.12585	2.12704
8.4	2.12823	2.12942	2.13061	2.13180	2.13298	2.13417	2.13535	2.13653	2.13771	2.13889
8.5	2.14007	2.14124	2.14242	2.14359	2.14476	2.14593	2.14710	2.14827	2.14943	2.15060
8.6	2.15176	2.15292	2.15409	2.15524	2.15640	2.15756	2.15871	2.15987	2.16102	2.16217
8.7	2.16332	2.16447	2.16562	2.16677	2.16791	2.16905	2.17020	2.17134	2.17248	2.17361
8.8	2.17475	2.17589	2.17702	2.17816	2.17929	2.18042	2.18155	2.18267	2.18380	2.18493
8.9	2.18605	2.18717	2.18830	2.18942	2.19054	2.19165	2.19277	2.19389	2.19500	2.19611
9.0	2.19722	2.19834	2.19944	2.20055	2.20166	2.20276	2.20387	2.20497	2.20607	2.20717
9.1	2.20827	2.20937	2.21047	2.21157	2.21266	2.21375	2.21485	2.21594	2.21703	2.21812
9.2	2.21920	2.22029	2.22138	2.22246	2.22354	2.22462	2.22570	2.22678	2.22786	2.22894
9.3	2.23001	2.23109	2.23216	2.23324	2.23431	2.23538	2.23645	2.23751	2.23858	2.23965
9.4	2.24071	2.24177	2.24284	2.24390	2.24496	2.24601	2.24707	2.24813	2.24918	2.25024
9.5	2.25129	2.25234	2.25339	2.25444	2.25549	2.25654	2.25759	2.25863	2.25968	2.26072
9.6	2.26176	2.26280	2.26384	2.26488	2.26592	2.26696	2.26799	2.26903	2.27006	2.27109
9.7	2.27213	2.27316	2.27419	2.27521	2.27624	2.27727	2.27829	2.27932	2.28034	2.28136
9.8	2.28238	2.28340	2.28442	2.28544	2.28646	2.28747	2.28849	2.28950	2.29051	2.29152
9.9	2.29253	2.29354	2.29455	2.29556	2.29657	2.29757	2.29858	2.29958	2.30058	2.30158

## 8

## EXPONENTIAL FUNCTIONS

 $e^x$ 

$x$	0	1	2	3	4	5	6	7	8	9
.0	1.0000	1.0101	1.0202	1.0305	1.0408	1.0513	1.0618	1.0725	1.0833	1.0942
.1	1.1052	1.1163	1.1275	1.1388	1.1503	1.1618	1.1735	1.1853	1.1972	1.2092
.2	1.2214	1.2337	1.2461	1.2586	1.2712	1.2840	1.2969	1.3100	1.3231	1.3364
.3	1.3499	1.3634	1.3771	1.3910	1.4049	1.4191	1.4333	1.4477	1.4623	1.4770
.4	1.4918	1.5068	1.5220	1.5373	1.5527	1.5683	1.5841	1.6000	1.6161	1.6323
.5	1.6487	1.6653	1.6820	1.6989	1.7160	1.7333	1.7507	1.7683	1.7860	1.8040
.6	1.8221	1.8404	1.8589	1.8776	1.8965	1.9155	1.9348	1.9542	1.9739	1.9937
.7	2.0138	2.0340	2.0544	2.0751	2.0959	2.1170	2.1383	2.1598	2.1815	2.2034
.8	2.2255	2.2479	2.2705	2.2933	2.3164	2.3396	2.3632	2.3869	2.4109	2.4351
.9	2.4596	2.4843	2.5093	2.5345	2.5600	2.5857	2.6117	2.6379	2.6645	2.6912
1.0	2.7183	2.7456	2.7732	2.8011	2.8292	2.8577	2.8864	2.9154	2.9447	2.9743
1.1	3.0042	3.0344	3.0649	3.0957	3.1268	3.1582	3.1899	3.2220	3.2544	3.2871
1.2	3.3201	3.3535	3.3872	3.4212	3.4556	3.4903	3.5254	3.5609	3.5966	3.6328
1.3	3.6693	3.7062	3.7434	3.7810	3.8190	3.8574	3.8962	3.9354	3.9749	4.0149
1.4	4.0552	4.0960	4.1371	4.1787	4.2207	4.2631	4.3060	4.3492	4.3929	4.4371
1.5	4.4817	4.5267	4.5722	4.6182	4.6646	4.7115	4.7588	4.8066	4.8550	4.9037
1.6	4.9530	5.0028	5.0531	5.1039	5.1552	5.2070	5.2593	5.3122	5.3656	5.4195
1.7	5.4739	5.5290	5.5845	5.6407	5.6973	5.7546	5.8124	5.8709	5.9299	5.9895
1.8	6.0496	6.1104	6.1719	6.2339	6.2965	6.3598	6.4237	6.4883	6.5535	6.6194
1.9	6.6859	6.7531	6.8210	6.8895	6.9588	7.0287	7.0993	7.1707	7.2427	7.3155
2.0	7.3891	7.4633	7.5383	7.6141	7.6906	7.7679	7.8460	7.9248	8.0045	8.0849
2.1	8.1662	8.2482	8.3311	8.4149	8.4994	8.5849	8.6711	8.7583	8.8463	8.9352
2.2	9.0250	9.1157	9.2073	9.2999	9.3933	9.4877	9.5831	9.6794	9.7767	9.8749
2.3	9.9742	10.074	10.176	10.278	10.381	10.486	10.591	10.697	10.805	10.913
2.4	11.023	11.134	11.246	11.359	11.473	11.588	11.705	11.822	11.941	12.061
2.5	12.182	12.305	12.429	12.554	12.680	12.807	12.936	13.066	13.197	13.330
2.6	13.464	13.599	13.736	13.874	14.013	14.154	14.296	14.440	14.585	14.732
2.7	14.880	15.029	15.180	15.333	15.487	15.643	15.800	15.959	16.119	16.281
2.8	16.445	16.610	16.777	16.945	17.116	17.288	17.462	17.637	17.814	17.993
2.9	18.174	18.357	18.541	18.728	18.916	19.106	19.298	19.492	19.688	19.886
3.0	20.086	20.287	20.491	20.697	20.905	21.115	21.328	21.542	21.758	21.977
3.1	22.198	22.421	22.646	22.874	23.104	23.336	23.571	23.807	24.047	24.288
3.2	24.533	24.779	25.028	25.280	25.534	25.790	26.050	26.311	26.576	26.843
3.3	27.113	27.385	27.660	27.938	28.219	28.503	28.789	29.079	29.371	29.666
3.4	29.964	30.265	30.569	30.877	31.187	31.500	31.817	32.137	32.460	32.786
3.5	33.115	33.448	33.784	34.124	34.467	34.813	35.163	35.517	35.874	36.234
3.6	36.598	36.966	37.338	37.713	38.092	38.475	38.861	39.252	39.646	40.045
3.7	40.447	40.854	41.264	41.679	42.098	42.521	42.948	43.380	43.816	44.256
3.8	44.701	45.150	45.604	46.063	46.525	46.993	47.465	47.942	48.424	48.911
3.9	49.402	49.899	50.400	50.907	51.419	51.935	52.457	52.985	53.517	54.055
4.	54.598	60.340	66.686	73.700	81.451	90.017	99.484	109.95	121.51	134.29
5.	148.41	164.02	181.27	200.34	221.41	244.69	270.43	298.87	330.30	365.04
6.	403.43	445.86	492.75	544.57	601.85	665.14	735.10	812.41	897.85	992.27
7.	1096.6	1212.0	1339.4	1480.3	1636.0	1808.0	1998.2	2208.3	2440.6	2697.3
8.	2981.0	3294.5	3641.0	4023.9	4447.1	4914.8	5431.7	6002.9	6634.2	7332.0
9.	8103.1	8955.3	9897.1	10938	12088	13360	14765	16318	18034	19930
10.	22026									

## 9

## EXPONENTIAL FUNCTIONS

$e^{-x}$

$x$	0	1	2	3	4	5	6	7	8	9
.0	1.00000	.99005	.98020	.97045	.96079	.95123	.94176	.93239	.92312	.91393
.1	.90484	.89583	.88692	.87810	.86936	.86071	.85214	.84366	.83527	.82696
.2	.81873	.81058	.80252	.79453	.78663	.77880	.77105	.76338	.75578	.74826
.3	.74082	.73345	.72615	.71892	.71177	.70469	.69768	.69073	.68386	.67706
.4	.67032	.66365	.65705	.65051	.64404	.63763	.63128	.62500	.61878	.61263
.5	.60653	.60050	.59452	.58860	.58275	.57695	.57121	.56553	.55990	.55433
.6	.54881	.54335	.53794	.53259	.52729	.52205	.51685	.51171	.50662	.50158
.7	.49659	.49164	.48675	.48191	.47711	.47237	.46767	.46301	.45841	.45384
.8	.44933	.44486	.44043	.43605	.43171	.42741	.42316	.41895	.41478	.41066
.9	.40657	.40252	.39852	.39455	.39063	.38674	.38289	.37908	.37531	.37158
1.0	.36788	.36422	.36060	.35701	.35345	.34994	.34646	.34301	.33960	.33622
1.1	.33287	.32956	.32628	.32303	.31982	.31664	.31349	.31037	.30728	.30422
1.2	.30119	.29820	.29523	.29229	.28938	.28650	.28365	.28083	.27804	.27527
1.3	.27253	.26982	.26714	.26448	.26185	.25924	.25666	.25411	.25158	.24908
1.4	.24660	.24414	.24171	.23931	.23693	.23457	.23224	.22993	.22764	.22537
1.5	.22313	.22091	.21871	.21654	.21438	.21225	.21014	.20805	.20598	.20393
1.6	.20190	.19989	.19790	.19593	.19398	.19205	.19014	.18825	.18637	.18452
1.7	.18268	.18087	.17907	.17728	.17552	.17377	.17204	.17033	.16864	.16696
1.8	.16530	.16365	.16203	.16041	.15882	.15724	.15567	.15412	.15259	.15107
1.9	.14957	.14808	.14661	.14515	.14370	.14227	.14086	.13946	.13807	.13670
2.0	.13534	.13399	.13266	.13134	.13003	.12873	.12745	.12619	.12493	.12369
2.1	.12246	.12124	.12003	.11884	.11765	.11648	.11533	.11418	.11304	.11192
2.2	.11080	.10970	.10861	.10753	.10646	.10540	.10435	.10331	.10228	.10127
2.3	.10026	.09926	.09827	.09730	.09633	.09537	.09442	.09348	.09255	.09163
2.4	.09072	.08982	.08892	.08804	.08716	.08629	.08543	.08458	.08374	.08291
2.5	.08208	.08127	.08046	.07966	.07887	.07808	.07730	.07654	.07577	.07502
2.6	.07427	.07353	.07280	.07208	.07136	.07065	.06995	.06925	.06856	.06788
2.7	.06721	.06654	.06587	.06522	.06457	.06393	.06329	.06266	.06204	.06142
2.8	.06081	.06020	.05961	.05901	.05843	.05784	.05727	.05670	.05613	.05558
2.9	.05502	.05448	.05393	.05340	.05287	.05234	.05182	.05130	.05079	.05029
3.0	.04979	.04929	.04880	.04832	.04783	.04736	.04689	.04642	.04596	.04550
3.1	.04505	.04460	.04416	.04372	.04328	.04285	.04243	.04200	.04159	.04117
3.2	.04076	.04036	.03996	.03956	.03916	.03877	.03839	.03801	.03763	.03725
3.3	.03688	.03652	.03615	.03579	.03544	.03508	.03474	.03439	.03405	.03371
3.4	.03337	.03304	.03271	.03239	.03206	.03175	.03143	.03112	.03081	.03050
3.5	.03020	.02990	.02960	.02930	.02901	.02872	.02844	.02816	.02788	.02760
3.6	.02732	.02705	.02678	.02652	.02625	.02599	.02573	.02548	.02522	.02497
3.7	.02472	.02448	.02423	.02399	.02375	.02352	.02328	.02305	.02282	.02260
3.8	.02237	.02215	.02193	.02171	.02149	.02128	.02107	.02086	.02065	.02045
3.9	.02024	.02004	.01984	.01964	.01945	.01925	.01906	.01887	.01869	.01850
4.	.018316	.016573	.014996	.013569	.012277	.011109	.010052	.0290953	.0282297	.0274466
5.	.0267379	.0260967	.0255166	.0249916	.0245166	.0240868	.0236979	.0233460	.0230276	.0227394
6.	.0224788	.0222429	.0220294	.0218363	.0216616	.0215034	.0213604	.0212309	.0211138	.0210078
7.	.0391188	.0382510	.0374659	.0367554	.0361125	.0355308	.0350045	.0345283	.0340973	.0337074
8.	.033546	.0330354	.0327465	.0324852	.0322487	.0320347	.0318411	.0316659	.0315073	.0313639
9.	.0312341	.0311167	.0310104	.0491424	.0482724	.0474852	.0467729	.0461283	.0455452	.0450175
10.	.0445400									

# 10

## EXPONENTIAL, SINE, AND COSINE INTEGRALS

$$\text{Ei}(x) = \int_x^{\infty} \frac{e^{-u}}{u} du, \quad \text{Si}(x) = \int_0^x \frac{\sin u}{u} du, \quad \text{Ci}(x) = \int_x^{\infty} \frac{\cos u}{u} du$$

<i>x</i>	Ei( <i>x</i> )	Si( <i>x</i> )	Ci( <i>x</i> )
.0	$\infty$	.0000	$\infty$
.5	.5598	.4931	.1778
1.0	.2194	.9461	-.3374
1.5	.1000	1.3247	-.4704
2.0	.04890	1.6054	-.4230
2.5	.02491	1.7785	-.2859
3.0	.01305	1.8487	-.1196
3.5	.026970	1.8331	.0321
4.0	.023779	1.7582	.1410
4.5	.022073	1.6541	.1935
5.0	.021148	1.5499	.1900
5.5	.036409	1.4687	.1421
6.0	.033601	1.4247	.0681
6.5	.032034	1.4218	-.0111
7.0	.031155	1.4546	-.0767
7.5	.046583	1.5107	-.1156
8.0	.043767	1.5742	-.1224
8.5	.042162	1.6296	-.09943
9.0	.041245	1.6650	-.05535
9.5	.057185	1.6745	-.022678
10.0	.054157	1.6583	.04546

## Section II: Factorial and Gamma Function, Binomial Coefficients

# 11

### FACTORIAL $n$

$$n! = 1 \cdot 2 \cdot 3 \cdots \cdot n$$

$n$	$n!$
0	1 (by definition)
1	1
2	2
3	6
4	24
5	120
6	720
7	5040
8	40,320
9	362,880
10	3,628,800
11	39,916,800
12	479,001,600
13	6,227,020,800
14	87,178,291,200
15	1,307,674,368,000
16	20,922,789,888,000
17	355,687,428,096,000
18	6,402,373,705,728,000
19	121,645,100,408,832,000
20	2,432,902,008,176,640,000
21	51,090,942,171,709,440,000
22	1,124,000,727,777,607,680,000
23	25,852,016,738,884,976,640,000
24	620,448,401,733,239,439,360,000
25	15,511,210,043,330,985,984,000,000
26	403,291,461,126,605,635,584,000,000
27	10,888,869,450,418,352,160,768,000,000
28	304,888,344,611,713,860,501,504,000,000
29	8,841,761,993,739,701,954,543,616,000,000
30	265,252,859,812,191,058,636,308,480,000,000
31	$8.22284 \times 10^{33}$
32	$2.63131 \times 10^{35}$
33	$8.68332 \times 10^{36}$
34	$2.95233 \times 10^{38}$
35	$1.03331 \times 10^{40}$
36	$3.71993 \times 10^{41}$
37	$1.37638 \times 10^{43}$
38	$5.23023 \times 10^{44}$
39	$2.03979 \times 10^{46}$

$n$	$n!$
40	$8.15915 \times 10^{47}$
41	$3.34525 \times 10^{49}$
42	$1.40501 \times 10^{51}$
43	$6.04153 \times 10^{52}$
44	$2.65827 \times 10^{54}$
45	$1.19622 \times 10^{56}$
46	$5.50262 \times 10^{57}$
47	$2.58623 \times 10^{59}$
48	$1.24139 \times 10^{61}$
49	$6.08282 \times 10^{62}$
50	$3.04141 \times 10^{64}$
51	$1.55112 \times 10^{66}$
52	$8.06582 \times 10^{67}$
53	$4.27488 \times 10^{69}$
54	$2.30844 \times 10^{71}$
55	$1.26964 \times 10^{73}$
56	$7.10999 \times 10^{74}$
57	$4.05269 \times 10^{76}$
58	$2.35056 \times 10^{78}$
59	$1.38683 \times 10^{80}$
60	$8.32099 \times 10^{81}$
61	$5.07580 \times 10^{83}$
62	$3.14700 \times 10^{85}$
63	$1.98261 \times 10^{87}$
64	$1.26887 \times 10^{89}$
65	$8.24765 \times 10^{90}$
66	$5.44345 \times 10^{92}$
67	$3.64711 \times 10^{94}$
68	$2.48004 \times 10^{96}$
69	$1.71122 \times 10^{98}$
70	$1.19786 \times 10^{100}$
71	$8.50479 \times 10^{101}$
72	$6.12345 \times 10^{103}$
73	$4.47012 \times 10^{105}$
74	$3.30789 \times 10^{107}$
75	$2.48091 \times 10^{109}$
76	$1.88549 \times 10^{111}$
77	$1.45183 \times 10^{113}$
78	$1.13243 \times 10^{115}$
79	$8.94618 \times 10^{116}$

$n$	$n!$
80	$7.15695 \times 10^{118}$
81	$5.79713 \times 10^{120}$
82	$4.75364 \times 10^{122}$
83	$3.94552 \times 10^{124}$
84	$3.31424 \times 10^{126}$
85	$2.81710 \times 10^{128}$
86	$2.42271 \times 10^{130}$
87	$2.10776 \times 10^{132}$
88	$1.85483 \times 10^{134}$
89	$1.65080 \times 10^{136}$
90	$1.48572 \times 10^{138}$
91	$1.35200 \times 10^{140}$
92	$1.24384 \times 10^{142}$
93	$1.15677 \times 10^{144}$
94	$1.08737 \times 10^{146}$
95	$1.03300 \times 10^{148}$
96	$9.91678 \times 10^{149}$
97	$9.61928 \times 10^{151}$
98	$9.42689 \times 10^{153}$
99	$9.33262 \times 10^{155}$
100	$9.33262 \times 10^{157}$

# 12

## GAMMA FUNCTION

$$\Gamma(x) = \int_x^{\infty} t^{x-1} e^{-t} dt \quad \text{for } 1 \leq x \leq 2$$

[For other values use the formula  $\Gamma(x + 1) = x \Gamma(x)$ ]

$x$	$\Gamma(x)$	$x$	$\Gamma(x)$
1.00	1.00000	1.50	.88623
1.01	.99433	1.51	.88659
1.02	.98884	1.52	.88704
1.03	.98355	1.53	.88757
1.04	.97844	1.54	.88818
1.05	.97350	1.55	.88887
1.06	.96874	1.56	.88964
1.07	.96415	1.57	.89049
1.08	.95973	1.58	.89142
1.09	.95546	1.59	.89243
1.10	.95135	1.60	.89352
1.11	.94740	1.61	.89468
1.12	.94359	1.62	.89592
1.13	.93993	1.63	.89724
1.14	.93642	1.64	.89864
1.15	.93304	1.65	.90012
1.16	.92980	1.66	.90167
1.17	.92670	1.67	.90330
1.18	.92373	1.68	.90500
1.19	.92089	1.69	.90678
1.20	.91817	1.70	.90864
1.21	.91558	1.71	.91057
1.22	.91311	1.72	.91258
1.23	.91075	1.73	.91467
1.24	.90852	1.74	.91683
1.25	.90640	1.75	.91906
1.26	.90440	1.76	.92137
1.27	.90250	1.77	.92376
1.28	.90072	1.78	.92623
1.29	.89904	1.79	.92877
1.30	.89747	1.80	.93138
1.31	.89600	1.81	.93408
1.32	.89464	1.82	.93685
1.33	.89338	1.83	.93969
1.34	.89222	1.84	.94261
1.35	.89115	1.85	.94561
1.36	.89018	1.86	.94869
1.37	.88931	1.87	.95184
1.38	.88854	1.88	.95507
1.39	.88785	1.89	.95838
1.40	.88726	1.90	.96177
1.41	.88676	1.91	.96523
1.42	.88636	1.92	.96877
1.43	.88604	1.93	.97240
1.44	.88581	1.94	.97610
1.45	.88566	1.95	.97988
1.46	.88560	1.96	.98374
1.47	.88563	1.97	.98768
1.48	.88575	1.98	.99171
1.49	.88595	1.99	.99581
1.50	.88623	2.00	1.00000

# 13

## BINOMIAL COEFFICIENTS

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1)\cdots(n-k+1)}{k!} = \binom{n}{n-k}, \quad 0! = 1$$

Note that each number is the sum of two numbers in the row above; one of these numbers is in the same column and the other is in the preceding column (e.g.,  $56 = 35 + 21$ ). The arrangement is often called *Pascal's triangle* (see 3.6, page 8).

$n \backslash k$	0	1	2	3	4	5	6	7	8	9
1	1	1								
2	1	2	1							
3	1	3	3	1						
4	1	4	6	4	1					
5	1	5	10	10	5	1				
6	1	6	15	20	15	6	1			
7	1	7	21	35	35	21	7	1		
8	1	8	28	56	70	56	28	8	1	
9	1	9	36	84	126	126	84	36	9	1
10	1	10	45	120	210	252	210	120	45	10
11	1	11	55	165	330	462	462	330	165	55
12	1	12	66	220	495	792	924	792	495	220
13	1	13	78	286	715	1287	1716	1716	1287	715
14	1	14	91	364	1001	2002	3003	3432	3003	2002
15	1	15	105	455	1365	3003	5005	6435	6435	5005
16	1	16	120	560	1820	4368	8008	11440	12870	11440
17	1	17	136	680	2380	6188	12376	19448	24310	24310
18	1	18	153	816	3060	8568	18564	31824	43758	48620
19	1	19	171	969	3876	11628	27132	50388	75582	92378
20	1	20	190	1140	4845	15504	38760	77520	125970	167960
21	1	21	210	1330	5985	20349	54264	116280	203490	293930
22	1	22	231	1540	7315	26334	74613	170544	319770	497420
23	1	23	253	1771	8855	33649	100947	245157	490314	817190
24	1	24	276	2024	10626	42504	134596	346104	735471	1307504
25	1	25	300	2300	12650	53130	177100	480700	1081575	2042975
26	1	26	325	2600	14950	65780	230230	657800	1562275	3124550
27	1	27	351	2925	17550	80730	296010	888030	2220075	4686825
28	1	28	378	3276	20475	98280	376740	1184040	3108105	6906900
29	1	29	406	3654	23751	118755	475020	1560780	4292145	10015005
30	1	30	435	4060	27405	142506	593775	2035800	5852925	14307150

# 13

## BINOMIAL COEFFICIENTS

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1)\cdots(n-k+1)}{k!} = \binom{n}{n-k}, \quad 0! = 1$$

*(Continued)*

$\backslash$	$k$	10	11	12	13	14	15
$n$							
10	1						
11	11	1					
12	66	12	1				
13	286	78	13				
14	1001	364	91	14			
15	3003	1365	455	105	15		1
16	8008	4368	1820	560	120	16	
17	19448	12376	6188	2380	680	136	
18	48758	31824	18564	8568	3060	816	
19	92378	75582	50388	27132	11628	3876	
20	184756	167960	125970	77520	38760	15504	
21	352716	352716	293930	203490	116280	54264	
22	646646	705432	646646	497420	319770	170544	
23	1144066	1352078	1352078	1144066	817190	490314	
24	1961256	2496144	2704156	2496144	1961256	1307504	
25	3268760	4457400	5200300	5200300	4457400	3268760	
26	5311735	7726160	9657700	10400600	9657700	7726160	
27	8436285	13037895	17383860	20058300	20058300	17383860	
28	13123110	21474180	30421755	37442160	40116600	37442160	
29	20030010	34597290	51895935	67863915	77558760	77558760	
30	30045015	54627300	86493225	119759850	145422675	155117520	

For  $k > 15$  use the fact that  $\binom{n}{k} = \binom{n}{n-k}$ .

### Section III: Bessel Functions

# 14

### BESSEL FUNCTIONS

$$J_0(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	1.0000	.9975	.9900	.9776	.9604	.9385	.9120	.8812	.8463	.8075
1.	.7652	.7196	.6711	.6201	.5669	.5118	.4554	.3980	.3400	.2818
2.	.2239	.1666	.1104	.0555	.0025	-.0484	-.0968	-.1424	-.1850	-.2243
3.	-.2601	-.2921	-.3202	-.3443	-.3643	-.3801	-.3918	-.3992	-.4026	-.4018
4.	-.3971	-.3887	-.3766	-.3610	-.3423	-.3205	-.2961	-.2693	-.2404	-.2097
5.	-.1776	-.1443	-.1103	-.0758	-.0412	-.0068	.0270	.0599	.0917	.1220
6.	.1506	.1773	.2017	.2238	.2433	.2601	.2740	.2851	.2931	.2981
7.	.3001	.2991	.2951	.2882	.2786	.2663	.2516	.2346	.2154	.1944
8.	.1717	.1475	.1222	.0960	.0692	.0419	.0146	-.0125	-.0392	-.0653
9.	-.0903	-.1142	-.1367	-.1577	-.1768	-.1939	-.2090	-.2218	-.2323	-.2403

# 15

### BESSEL FUNCTIONS

$$J_1(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	.0000	.0499	.0995	.1483	.1960	.2423	.2867	.3290	.3688	.4059
1.	.4401	.4709	.4983	.5220	.5419	.5579	.5699	.5778	.5815	.5812
2.	.5767	.5683	.5560	.5399	.5202	.4971	.4708	.4416	.4097	.3754
3.	.3391	.3009	.2613	.2207	.1792	.1374	.0955	.0538	.0128	-.0272
4.	-.0660	-.1033	-.1386	-.1719	-.2028	-.2311	-.2566	-.2791	-.2985	-.3147
5.	-.3276	-.3371	-.3432	-.3460	-.3453	-.3414	-.3343	-.3241	-.3110	-.2951
6.	-.2767	-.2559	-.2329	-.2081	-.1816	-.1538	-.1250	-.0953	-.0652	-.0349
7.	-.0047	.0252	.0543	.0826	.1096	.1352	.1592	.1813	.2014	.2192
8.	.2346	.2476	.2580	.2657	.2708	.2731	.2728	.2697	.2641	.2559
9.	.2453	.2324	.2174	.2004	.1816	.1613	.1395	.1166	.0928	.0684

# 16

## BESSEL FUNCTIONS

$$Y_0(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	$-\infty$	-1.5342	-1.0811	-.8073	-.6060	-.4445	-.3085	-.1907	-.0868	.0056
1.	.0883	.1622	.2281	.2865	.3379	.3824	.4204	.4520	.4774	.4968
2.	.5104	.5183	.5208	.5181	.5104	.4981	.4813	.4605	.4359	.4079
3.	.3769	.3431	.3071	.2691	.2296	.1890	.1477	.1061	.0645	.0234
4.	-.0169	-.0561	-.0938	-.1296	-.1633	-.1947	-.2235	-.2494	-.2723	-.2921
5.	-.3085	-.3216	-.3313	-.3374	-.3402	-.3395	-.3354	-.3282	-.3177	-.3044
6.	-.2882	-.2694	-.2483	-.2251	-.1999	-.1732	-.1452	-.1162	-.0864	-.0563
7.	-.0259	.0042	.0339	.0628	.0907	.1173	.1424	.1658	.1872	.2065
8.	.2235	.2381	.2501	.2595	.2662	.2702	.2715	.2700	.2659	.2592
9.	.2499	.2383	.2245	.2086	.1907	.1712	.1502	.1279	.1045	.0804

# 17

## BESSEL FUNCTIONS

$$Y_1(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	$-\infty$	-6.4590	-3.3238	-2.2931	-1.7809	-1.4715	-1.2604	-1.1032	-.9781	-.8731
1.	-.7812	-.6981	-.6211	-.5485	-.4791	-.4123	-.3476	-.2847	-.2237	-.1644
2.	-.1070	-.0517	.0015	.0523	.1005	.1459	.1884	.2276	.2635	.2959
3.	.3247	.3496	.3707	.3879	.4010	.4102	.4154	.4167	.4141	.4078
4.	.3979	.3846	.3680	.3484	.3260	.3010	.2737	.2445	.2136	.1812
5.	.1479	.1137	.0792	.0445	.0101	-.0238	-.0568	-.0887	-.1192	-.1481
6.	-.1750	-.1998	-.2223	-.2422	-.2596	-.2741	-.2857	-.2945	-.3002	-.3029
7.	-.3027	-.2995	-.2934	-.2846	-.2731	-.2591	-.2428	-.2243	-.2039	-.1817
8.	-.1581	-.1331	-.1072	-.0806	-.0535	-.0262	.0011	.0280	.0544	.0799
9.	.1043	.1275	.1491	.1691	.1871	.2032	.2171	.2287	.2379	.2447

# 18

## BESSEL FUNCTIONS

$$I_0(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	1.000	1.003	1.010	1.023	1.040	1.063	1.092	1.126	1.167	1.213
1.	1.266	1.326	1.394	1.469	1.553	1.647	1.750	1.864	1.990	2.128
2.	2.280	2.446	2.629	2.830	3.049	3.290	3.553	3.842	4.157	4.503
3.	4.881	5.294	5.747	6.243	6.785	7.378	8.028	8.739	9.517	10.37
4.	11.30	12.32	13.44	14.67	16.01	17.48	19.09	20.86	22.79	24.91
5.	27.24	29.79	32.58	35.65	39.01	42.69	46.74	51.17	56.04	61.38
6.	67.23	73.66	80.72	88.46	96.96	106.3	116.5	127.8	140.1	153.7
7.	168.6	185.0	202.9	222.7	244.3	268.2	294.3	323.1	354.7	389.4
8.	427.6	469.5	515.6	566.3	621.9	683.2	750.5	824.4	905.8	995.2
9.	1094	1202	1321	1451	1595	1753	1927	2119	2329	2561

# 19

## BESSEL FUNCTIONS

$$I_1(x)$$

$x$	0	1	2	3	4	5	6	7	8	9
0.	.0000	.0501	.1005	.1517	.2040	.2579	.3137	.3719	.4329	.4971
1.	.5652	.6375	.7147	.7973	.8861	.9817	1.085	1.196	1.317	1.448
2.	1.591	1.745	1.914	2.098	2.298	2.517	2.755	3.016	3.301	3.613
3.	3.953	4.326	4.734	5.181	5.670	6.206	6.793	7.436	8.140	8.913
4.	9.759	10.69	11.71	12.82	14.05	15.39	16.86	18.48	20.25	22.20
5.	24.34	26.68	29.25	32.08	35.18	38.59	42.33	46.44	50.95	55.90
6.	61.34	67.32	73.89	81.10	89.03	97.74	107.3	117.8	129.4	142.1
7.	156.0	171.4	188.3	206.8	227.2	249.6	274.2	301.3	331.1	363.9
8.	399.9	439.5	483.0	531.0	583.7	641.6	705.4	775.5	852.7	937.5
9.	1031	1134	1247	1371	1508	1658	1824	2006	2207	2428

**20****BESSEL FUNCTIONS**

$K_0(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	$\infty$	2.4271	1.7527	1.3725	1.1145	.9244	.7775	.6605	.5653	.4867
1.	.4210	.3656	.3185	.2782	.2437	.2138	.1880	.1655	.1459	.1288
2.	.1139	.1008	.08927	.07914	.07022	.06235	.05540	.04926	.04382	.03901
3.	.03474	.03095	.02759	.02461	.02196	.01960	.01750	.01563	.01397	.01248
4.	.01116	.029980	.028927	.027988	.027149	.026400	.025730	.025132	.024597	.024119
5.	.023691	.023308	.022966	.022659	.022385	.022139	.021918	.021721	.021544	.021386
6.	.021244	.021117	.021003	.039001	.038083	.037259	.036520	.035857	.035262	.034728
7.	.034248	.033817	.033431	.033084	.032772	.032492	.032240	.032014	.031811	.031629
8.	.031465	.031317	.031185	.031066	.049588	.048626	.047761	.046983	.046283	.045654
9.	.045088	.044579	.044121	.043710	.043339	.043006	.042706	.042436	.042193	.041975

**21****BESSEL FUNCTIONS**

$K_1(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	$\infty$	9.8538	4.7760	3.0560	2.1844	1.6564	1.3028	1.0503	.8618	.7165
1.	.6019	.5098	.4346	.3725	.3208	.2774	.2406	.2094	.1826	.1597
2.	.1399	.1227	.1079	.09498	.08372	.07389	.06528	.05774	.05111	.04529
3.	.04016	.03563	.03164	.02812	.02500	.02224	.01979	.01763	.01571	.01400
4.	.01248	.01114	.029938	.028872	.027923	.027078	.026325	.025654	.025055	.024521
5.	.024045	.023619	.023239	.022900	.022597	.022326	.022083	.021866	.021673	.021499
6.	.021344	.021205	.021081	.039691	.038693	.037799	.036998	.036280	.035636	.035059
7.	.034542	.034078	.033662	.033288	.032953	.032653	.032383	.032141	.031924	.031729
8.	.031554	.031396	.031255	.031128	.031014	.049120	.048200	.047374	.046631	.045964
9.	.045364	.044825	.044340	.043904	.043512	.043160	.042843	.042559	.042302	.042072

# 22

## BESSEL FUNCTIONS $\text{Ber}(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	1.0000	1.0000	1.0000	.9999	.9996	.9990	.9980	.9962	.9936	.9898
1.	.9844	.9771	.9676	.9554	.9401	.9211	.8979	.8700	.8367	.7975
2.	.7517	.6987	.6377	.5680	.4890	.4000	.3001	.1887	.06511	-.07137
3.	-.2214	-.3855	-.5644	-.7584	-.9680	-1.1936	-1.4353	-1.6933	-1.9674	-2.2576
4.	-2.5634	-2.8843	-3.2195	-3.5679	-3.9283	-4.2991	-4.6784	-5.0639	-5.4531	-5.8429
5.	-6.2301	-6.6107	-6.9803	-7.3344	-7.6674	-7.9736	-8.2466	-8.4794	-8.6644	-8.7937
6.	-8.8583	-8.8491	-8.7561	-8.5688	-8.2762	-7.8669	-7.3287	-6.6492	-5.8155	-4.8146
7.	-3.6329	-2.2571	-0.6737	1.1308	3.1695	5.4550	7.9994	10.814	13.909	17.293
8.	20.974	24.957	29.245	33.840	38.738	43.936	49.423	55.187	61.210	67.469
9.	73.936	80.576	87.350	94.208	101.10	107.95	114.70	121.26	127.54	133.43

# 23

## BESSEL FUNCTIONS $\text{Bei}(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	.0000	.022500	.01000	.02250	.04000	.06249	.08998	.1224	.1599	.2023
1.	.2496	.3017	.3587	.4204	.4867	.5576	.6327	.7120	.7953	.8821
2.	.9723	1.0654	1.1610	1.2585	1.3575	1.4572	1.5569	1.6557	1.7529	1.8472
3.	1.9376	2.0228	2.1016	2.1723	2.2334	2.2832	2.3199	2.3413	2.3454	2.3300
4.	2.2927	2.2309	2.1422	2.0236	1.8726	1.6860	1.4610	1.1946	.8837	.5251
5.	.1160	-.3467	-.8658	-1.4443	-2.0845	-2.7890	-3.5597	-4.3986	-5.3068	-6.2854
6.	-.73347	-8.4545	-9.6437	-10.901	-12.223	-13.607	-15.047	-16.538	-18.074	-19.644
7.	-21.239	-22.848	-24.456	-26.049	-27.609	-29.116	-30.548	-31.882	-33.092	-34.147
8.	-35.017	-35.667	-36.061	-36.159	-35.920	-35.298	-34.246	-32.714	-30.651	-28.003
9.	-24.713	-20.724	-15.976	-10.412	-3.9693	3.4106	11.787	21.218	31.758	43.459

# 24

## BESSEL FUNCTIONS

$\text{Ker}(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	$\infty$	2.4205	1.7331	1.3372	1.0626	.8559	.6931	.5614	.4529	.3625
1.	.2867	.2228	.1689	.1235	.08513	.05293	.02603	.023691	-.01470	-.02966
2.	-.04166	-.05111	-.05834	-.06367	-.06737	-.06969	-.07083	-.07097	-.07030	-.06894
3.	-.06703	-.06468	-.06198	-.05903	-.05590	-.05264	-.04932	-.04597	-.04265	-.03937
4.	-.03618	-.03308	-.03011	-.02726	-.02456	-.02200	-.01960	-.01734	-.01525	-.01330
5.	-.01151	-.029865	-.028359	-.026989	-.025749	-.024632	-.023632	-.022740	-.021952	-.021258
6.	-.0 <sup>3</sup> 6530	-.0 <sup>3</sup> 1295	.0 <sup>3</sup> 3191	.0 <sup>3</sup> 6991	.0 <sup>2</sup> 1017	.0 <sup>2</sup> 1278	.0 <sup>2</sup> 1488	.0 <sup>2</sup> 1653	.0 <sup>2</sup> 1777	.0 <sup>2</sup> 1866
7.	.0 <sup>2</sup> 1922	.0 <sup>2</sup> 1951	.0 <sup>2</sup> 1956	.0 <sup>2</sup> 1940	.0 <sup>2</sup> 1907	.0 <sup>2</sup> 1860	.0 <sup>2</sup> 1800	.0 <sup>2</sup> 1731	.0 <sup>2</sup> 1655	.0 <sup>2</sup> 1572
8.	.0 <sup>2</sup> 1486	.0 <sup>2</sup> 1397	.0 <sup>2</sup> 1306	.0 <sup>2</sup> 1216	.0 <sup>2</sup> 1126	.0 <sup>2</sup> 1037	.0 <sup>3</sup> 9511	.0 <sup>3</sup> 8675	.0 <sup>3</sup> 7871	.0 <sup>3</sup> 7102
9.	.0 <sup>3</sup> 6372	.0 <sup>3</sup> 5681	.0 <sup>3</sup> 5030	.0 <sup>3</sup> 4422	.0 <sup>3</sup> 3855	.0 <sup>3</sup> 3330	.0 <sup>3</sup> 2846	.0 <sup>3</sup> 2402	.0 <sup>3</sup> 1996	.0 <sup>3</sup> 1628

# 25

## BESSEL FUNCTIONS

$\text{Kei}(x)$

$x$	0	1	2	3	4	5	6	7	8	9
0.	-.7854	-.7769	-.7581	-.7331	-.7038	-.6716	-.6374	-.6022	-.5664	-.5305
1.	-.4950	-.4601	-.4262	-.3933	-.3617	-.3314	-.3026	-.2752	-.2494	-.2251
2.	-.2024	-.1812	-.1614	-.1431	-.1262	-.1107	-.09644	-.08342	-.07157	-.06083
3.	-.05112	-.04240	-.03458	-.02762	-.02145	-.01600	-.01123	-.027077	-.023487	-.0 <sup>3</sup> 4108
4.	.0 <sup>2</sup> 2198	.0 <sup>2</sup> 4386	.0 <sup>2</sup> 6194	.0 <sup>2</sup> 7661	.0 <sup>2</sup> 8826	.0 <sup>2</sup> 9721	.01038	.01083	.01110	.01121
5.	.01119	.01105	.01082	.01051	.01014	.0 <sup>2</sup> 9716	.0 <sup>2</sup> 9255	.0 <sup>2</sup> 8766	.0 <sup>2</sup> 8258	.0 <sup>2</sup> 7739
6.	.0 <sup>2</sup> 7216	.0 <sup>2</sup> 6696	.0 <sup>2</sup> 6183	.0 <sup>2</sup> 5681	.0 <sup>2</sup> 5194	.0 <sup>2</sup> 4724	.0 <sup>2</sup> 4274	.0 <sup>2</sup> 3846	.0 <sup>2</sup> 3440	.0 <sup>2</sup> 3058
7.	.0 <sup>2</sup> 2700	.0 <sup>2</sup> 2366	.0 <sup>2</sup> 2057	.0 <sup>2</sup> 1770	.0 <sup>2</sup> 1507	.0 <sup>2</sup> 1267	.0 <sup>2</sup> 1048	.0 <sup>3</sup> 8498	.0 <sup>3</sup> 6714	.0 <sup>3</sup> 5117
8.	.0 <sup>3</sup> 696	.0 <sup>3</sup> 2440	.0 <sup>3</sup> 1339	.0 <sup>4</sup> 3809	-.0 <sup>4</sup> 4449	-.0 <sup>3</sup> 1149	-.0 <sup>3</sup> 1742	-.0 <sup>3</sup> 2233	-.0 <sup>3</sup> 2632	-.0 <sup>3</sup> 2949
9.	-.0 <sup>3</sup> 3192	-.0 <sup>3</sup> 3368	-.0 <sup>3</sup> 3486	-.0 <sup>3</sup> 3552	-.0 <sup>3</sup> 3574	-.0 <sup>3</sup> 3557	-.0 <sup>3</sup> 3508	-.0 <sup>3</sup> 3430	-.0 <sup>3</sup> 3329	-.0 <sup>3</sup> 3210

# 26

## VALUES FOR APPROXIMATE ZEROS OF BESSEL FUNCTIONS

The following table lists the first few positive roots of various equations. Note that for all cases listed the successive large roots differ approximately by  $\pi = 3.14159\dots$

	$n = 0$	$n = 1$	$n = 2$	$n = 3$	$n = 4$	$n = 5$	$n = 6$
$J_n(x) = 0$	2.4048	3.8317	5.1356	6.3802	7.5883	8.7715	9.9361
	5.5201	7.0156	8.4172	9.7610	11.0647	12.3386	13.5893
	8.6537	10.1735	11.6198	13.0152	14.3725	15.7002	17.0038
	11.7915	13.3237	14.7960	16.2235	17.6160	18.9801	20.3208
	14.9309	16.4706	17.9598	19.4094	20.8269	22.2178	23.5861
	18.0711	19.6159	21.1170	22.5827	24.0190	25.4303	26.8202
$Y_n(x) = 0$	0.8936	2.1971	3.3842	4.5270	5.6452	6.7472	7.8377
	3.9577	5.4297	6.7938	8.0976	9.3616	10.5972	11.8110
	7.0861	8.5960	10.0235	11.3965	12.7301	14.0338	15.3136
	10.2223	11.7492	13.2100	14.6231	15.9996	17.3471	18.6707
	13.3611	14.8974	16.3790	17.8185	19.2244	20.6029	21.9583
	16.5009	18.0434	19.5390	20.9973	22.4248	23.8265	25.2062
$J'_n(x) = 0$	0.0000	1.8412	3.0542	4.2012	5.3176	6.4156	7.5013
	3.8317	5.3314	6.7061	8.0152	9.2824	10.5199	11.7349
	7.0156	8.5363	9.9695	11.3459	12.6819	13.9872	15.2682
	10.1735	11.7060	13.1704	14.5859	15.9641	17.3128	18.6374
	13.3237	14.8636	16.3475	17.7888	19.1960	20.5755	21.9317
	16.4706	18.0155	19.5129	20.9725	22.4010	23.8036	25.1839
$Y'_n(x) = 0$	2.1971	3.6830	5.0026	6.2536	7.4649	8.6496	9.8148
	5.4297	6.9415	8.3507	9.6988	11.0052	12.2809	13.5328
	8.5960	10.1234	11.5742	12.9724	14.3317	15.6608	16.9655
	11.7492	13.2858	14.7609	16.1905	17.5844	18.9497	20.2913
	14.8974	16.4401	17.9313	19.3824	20.8011	22.1928	23.5619
	18.0434	19.5902	21.0929	22.5598	23.9970	25.4091	26.7995

## Section IV: Legendre Polynomials

**27**

### LEGENDRE POLYNOMIALS $P_n(x)$

$$[P_0(x)=1, P_1(x)=x]$$

$x$	$P_2(x)$	$P_3(x)$	$P_4(x)$	$P_5(x)$
.00	-.5000	.0000	.3750	.0000
.05	-.4963	-.0747	.3657	.0927
.10	-.4850	-.1475	.3379	.1788
.15	-.4663	-.2166	.2928	.2523
.20	-.4400	-.2800	.2320	.3075
.25	-.4063	-.3359	.1577	.3397
.30	-.3650	-.3825	.0729	.3454
.35	-.3163	-.4178	-.0187	.3225
.40	-.2600	-.4400	-.1130	.2706
.45	-.1963	-.4472	-.2050	.1917
.50	-.1250	-.4375	-.2891	.0898
.55	-.0463	-.4091	-.3590	-.0282
.60	.0400	-.3600	-.4080	-.1526
.65	.1338	-.2884	-.4284	-.2705
.70	.2350	-.1925	-.4121	-.3652
.75	.3438	-.0703	-.3501	-.4164
.80	.4600	.0800	-.2330	-.3995
.85	.5838	.2603	-.0506	-.2857
.90	.7150	.4725	.2079	-.0411
.95	.8538	.7184	.5541	.3727
1.00	1.0000	1.0000	1.0000	1.0000

# 28

**LEGENDRE POLYNOMIALS  $P_n(\cos \theta)$**   
 $[P_0(\cos \theta)=1]$

$\theta$	$P_1(\cos \theta)$	$P_2(\cos \theta)$	$P_3(\cos \theta)$	$P_4(\cos \theta)$	$P_5(\cos \theta)$
0°	1.0000	1.0000	1.0000	1.0000	1.0000
5°	.9962	.9886	.9773	.9623	.9437
10°	.9848	.9548	.9106	.8532	.7840
15°	.9659	.8995	.8042	.6847	.5471
20°	.9397	.8245	.6649	.4750	.2715
25°	.9063	.7321	.5016	.2465	.0009
30°	.8660	.6250	.3248	.0234	-.2233
35°	.8192	.5065	.1454	-.1714	-.3691
40°	.7660	.3802	-.0252	-.3190	-.4197
45°	.7071	.2500	-.1768	-.4063	-.3757
50°	.6428	.1198	-.3002	-.4275	-.2545
55°	.5736	-.0065	-.3886	-.3852	-.0868
60°	.5000	-.1250	-.4375	-.2891	.0898
65°	.4226	-.2321	-.4452	-.1552	.2381
70°	.3420	-.3245	-.4130	-.0038	.3281
75°	.2588	-.3995	-.3449	.1434	.3427
80°	.1737	-.4548	-.2474	.2659	.2810
85°	.0872	-.4886	-.1291	.3468	.1577
90°	.0000	-.5000	.0000	.3750	.0000

## Section V: Elliptic Integrals

**29**

### COMPLETE ELLIPTIC INTEGRALS OF FIRST AND SECOND KINDS

$$K = \int_0^{\pi/2} \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}, \quad E = \int_0^{\pi/2} \sqrt{1 - k^2 \sin^2 \theta} d\theta, \quad k = \sin \psi$$

$\psi$	$K$	$E$
0°	1.5708	1.5708
1	1.5709	1.5707
2	1.5713	1.5703
3	1.5719	1.5697
4	1.5727	1.5689
5	1.5738	1.5678
6	1.5751	1.5665
7	1.5767	1.5649
8	1.5785	1.5632
9	1.5805	1.5611
10	1.5828	1.5589
11	1.5854	1.5564
12	1.5882	1.5537
13	1.5913	1.5507
14	1.5946	1.5476
15	1.5981	1.5442
16	1.6020	1.5405
17	1.6061	1.5367
18	1.6105	1.5326
19	1.6151	1.5283
20	1.6200	1.5238
21	1.6252	1.5191
22	1.6307	1.5141
23	1.6365	1.5090
24	1.6426	1.5037
25	1.6490	1.4981
26	1.6557	1.4924
27	1.6627	1.4864
28	1.6701	1.4803
29	1.6777	1.4740
30	1.6858	1.4675

$\psi$	$K$	$E$
30°	1.6858	1.4675
31	1.6941	1.4608
32	1.7028	1.4539
33	1.7119	1.4469
34	1.7214	1.4397
35	1.7312	1.4323
36	1.7415	1.4248
37	1.7522	1.4171
38	1.7633	1.4092
39	1.7748	1.4013
40	1.7868	1.3931
41	1.7992	1.3849
42	1.8122	1.3765
43	1.8256	1.3680
44	1.8396	1.3594
45	1.8541	1.3506
46	1.8691	1.3418
47	1.8848	1.3329
48	1.9011	1.3238
49	1.9180	1.3147
50	1.9356	1.3055
51	1.9539	1.2963
52	1.9729	1.2870
53	1.9927	1.2776
54	2.0133	1.2681
55	2.0347	1.2587
56	2.0571	1.2492
57	2.0804	1.2397
58	2.1047	1.2301
59	2.1300	1.2206
60	2.1565	1.2111

$\psi$	$K$	$E$
60°	2.1565	1.2111
61	2.1842	1.2015
62	2.2132	1.1920
63	2.2435	1.1826
64	2.2754	1.1732
65	2.3088	1.1638
66	2.3439	1.1545
67	2.3809	1.1453
68	2.4198	1.1362
69	2.4610	1.1272
70	2.5046	1.1184
71	2.5507	1.1096
72	2.5998	1.1011
73	2.6521	1.0927
74	2.7081	1.0844
75	2.7681	1.0764
76	2.8327	1.0686
77	2.9026	1.0611
78	2.9786	1.0538
79	3.0617	1.0468
80	3.1534	1.0401
81	3.2553	1.0338
82	3.3699	1.0278
83	3.5004	1.0223
84	3.6519	1.0172
85	3.8317	1.0127
86	4.0528	1.0086
87	4.3387	1.0053
88	4.7427	1.0026
89	5.4349	1.0008
90	$\infty$	1.0000

# 30

**INCOMPLETE ELLIPTIC INTEGRAL  
OF THE FIRST KIND**

$$F(k, \phi) = \int_0^\phi \frac{d\theta}{\sqrt{1 - k^2 \sin^2 \theta}}, \quad k = \sin \psi$$

$\frac{\psi}{\phi}$	0°	10°	20°	30°	40°	50°	60°	70°	80°	90°
0°	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
10°	0.1745	0.1746	0.1746	0.1748	0.1749	0.1751	0.1752	0.1753	0.1754	0.1754
20°	0.3491	0.3493	0.3499	0.3508	0.3520	0.3533	0.3545	0.3555	0.3561	0.3564
30°	0.5236	0.5243	0.5263	0.5294	0.5334	0.5379	0.5422	0.5459	0.5484	0.5493
40°	0.6981	0.6997	0.7043	0.7116	0.7213	0.7323	0.7436	0.7535	0.7604	0.7629
50°	0.8727	0.8756	0.8842	0.8982	0.9173	0.9401	0.9647	0.9876	1.0044	1.0107
60°	1.0472	1.0519	1.0660	1.0896	1.1226	1.1643	1.2126	1.2619	1.3014	1.3170
70°	1.2217	1.2286	1.2495	1.2853	1.3372	1.4068	1.4944	1.5959	1.6918	1.7354
80°	1.3963	1.4056	1.4344	1.4846	1.5597	1.6660	1.8125	2.0119	2.2653	2.4362
90°	1.5708	1.5828	1.6200	1.6858	1.7868	1.9356	2.1565	2.5046	3.1534	∞

# 31

**INCOMPLETE ELLIPTIC INTEGRAL  
OF THE SECOND KIND**

$$E(k, \phi) = \int_0^\phi \sqrt{1 - k^2 \sin^2 \theta} d\theta, \quad k = \sin \psi$$

$\frac{\psi}{\phi}$	0°	10°	20°	30°	40°	50°	60°	70°	80°	90°
0°	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
10°	0.1745	0.1745	0.1744	0.1743	0.1742	0.1740	0.1739	0.1738	0.1737	0.1736
20°	0.3491	0.3489	0.3483	0.3473	0.3462	0.3450	0.3438	0.3429	0.3422	0.3420
30°	0.5236	0.5229	0.5209	0.5179	0.5141	0.5100	0.5061	0.5029	0.5007	0.5000
40°	0.6981	0.6966	0.6921	0.6851	0.6763	0.6667	0.6575	0.6497	0.6446	0.6428
50°	0.8727	0.8698	0.8614	0.8483	0.8317	0.8134	0.7954	0.7801	0.7697	0.7660
60°	1.0472	1.0426	1.0290	1.0076	0.9801	0.9493	0.9184	0.8914	0.8728	0.8660
70°	1.2217	1.2149	1.1949	1.1632	1.1221	1.0750	1.0266	0.9830	0.9514	0.9397
80°	1.3963	1.3870	1.3597	1.3161	1.2590	1.1926	1.1225	1.0565	1.0054	0.9848
90°	1.5708	1.5589	1.5238	1.4675	1.3931	1.3055	1.2111	1.1184	1.0401	1.0000

## Section VI: Financial Tables

# 32

### COMPOUND AMOUNT: $(1 + r)^n$

If a principal  $P$  is deposited at interest rate  $r$  (in decimals) compounded annually, then at the end of  $n$  years the accumulated amount  $A = P(1 + r)^n$ .

$n \backslash r$	1%	$1\frac{1}{4}\%$	$1\frac{1}{2}\%$	2%	$2\frac{1}{2}\%$	3%	4%	5%	6%
1	1.0100	1.0125	1.0150	1.0200	1.0250	1.0300	1.0400	1.0500	1.0600
2	1.0201	1.0252	1.0302	1.0404	1.0506	1.0609	1.0816	1.1025	1.1236
3	1.0303	1.0380	1.0457	1.0612	1.0769	1.0927	1.1249	1.1576	1.1910
4	1.0406	1.0509	1.0614	1.0824	1.1038	1.1255	1.1699	1.2155	1.2625
5	1.0510	1.0641	1.0773	1.1041	1.1314	1.1593	1.2167	1.2763	1.3382
6	1.0615	1.0774	1.0934	1.1262	1.1597	1.1941	1.2653	1.3401	1.4185
7	1.0721	1.0909	1.1098	1.1487	1.1887	1.2299	1.3159	1.4071	1.5036
8	1.0829	1.1045	1.1265	1.1717	1.2184	1.2668	1.3688	1.4775	1.5938
9	1.0937	1.1183	1.1434	1.1951	1.2489	1.3048	1.4233	1.5513	1.6895
10	1.1046	1.1323	1.1605	1.2190	1.2801	1.3439	1.4802	1.6289	1.7908
11	1.1157	1.1464	1.1779	1.2434	1.3121	1.3842	1.5395	1.7103	1.8983
12	1.1268	1.1608	1.1956	1.2682	1.3449	1.4258	1.6010	1.7959	2.0122
13	1.1381	1.1753	1.2136	1.2936	1.3785	1.4685	1.6651	1.8856	2.1329
14	1.1495	1.1900	1.2318	1.3195	1.4130	1.5126	1.7317	1.9799	2.2609
15	1.1610	1.2048	1.2502	1.3459	1.4483	1.5580	1.8009	2.0789	2.3966
16	1.1726	1.2199	1.2690	1.3728	1.4845	1.6047	1.8730	2.1829	2.5404
17	1.1843	1.2351	1.2880	1.4002	1.5216	1.6528	1.9479	2.2920	2.6928
18	1.1961	1.2506	1.3073	1.4282	1.5597	1.7024	2.0258	2.4066	2.8543
19	1.2081	1.2662	1.3270	1.4568	1.5987	1.7535	2.1068	2.5270	3.0256
20	1.2202	1.2820	1.3469	1.4859	1.6386	1.8061	2.1911	2.6533	3.2071
21	1.2324	1.2981	1.3671	1.5157	1.6796	1.8603	2.2788	2.7860	3.3996
22	1.2447	1.3143	1.3876	1.5460	1.7216	1.9161	2.3699	2.9253	3.6035
23	1.2572	1.3307	1.4084	1.5769	1.7646	1.9736	2.4647	3.0715	3.8197
24	1.2697	1.3474	1.4295	1.6084	1.8087	2.0328	2.5633	3.2251	4.0489
25	1.2824	1.3642	1.4509	1.6406	1.8539	2.0938	2.6658	3.3864	4.2919
26	1.2953	1.3812	1.4727	1.6734	1.9003	2.1566	2.7725	3.5557	4.5494
27	1.3082	1.3985	1.4948	1.7069	1.9478	2.2213	2.8834	3.7335	4.8223
28	1.3213	1.4160	1.5172	1.7410	1.9965	2.2879	2.9987	3.9201	5.1117
29	1.3345	1.4337	1.5400	1.7758	2.0464	2.3566	3.1187	4.1161	5.4184
30	1.3478	1.4516	1.5631	1.8114	2.0976	2.4273	3.2434	4.3219	5.7435
31	1.3613	1.4698	1.5865	1.8476	2.1500	2.5001	3.3731	4.5380	6.0881
32	1.3749	1.4881	1.6103	1.8845	2.2038	2.5751	3.5081	4.7649	6.4534
33	1.3887	1.5067	1.6345	1.9222	2.2589	2.6523	3.6484	5.0032	6.8406
34	1.4026	1.5256	1.6590	1.9607	2.3153	2.7319	3.7943	5.2533	7.2510
35	1.4166	1.5446	1.6839	1.9999	2.3732	2.8139	3.9461	5.5160	7.6861
36	1.4308	1.5639	1.7091	2.0399	2.4325	2.8983	4.1039	5.7918	8.1473
37	1.4451	1.5835	1.7348	2.0807	2.4933	2.9852	4.2681	6.0814	8.6361
38	1.4595	1.6033	1.7608	2.1223	2.5557	3.0748	4.4388	6.3855	9.1543
39	1.4741	1.6233	1.7872	2.1647	2.6196	3.1670	4.6164	6.7048	9.7035
40	1.4889	1.6436	1.8140	2.2080	2.6851	3.2620	4.8010	7.0400	10.2857
41	1.5038	1.6642	1.8412	2.2522	2.7522	3.3599	4.9931	7.3920	10.9029
42	1.5188	1.6850	1.8688	2.2972	2.8210	3.4607	5.1928	7.7616	11.5570
43	1.5340	1.7060	1.8969	2.3432	2.8915	3.5645	5.4005	8.1497	12.2505
44	1.5493	1.7274	1.9253	2.3901	2.9638	3.6715	5.6165	8.5572	12.9855
45	1.5648	1.7489	1.9542	2.4379	3.0379	3.7816	5.8412	8.9850	13.7646
46	1.5805	1.7708	1.9835	2.4866	3.1139	3.8950	6.0748	9.4343	14.5905
47	1.5963	1.7929	2.0133	2.5363	3.1917	4.0119	6.3178	9.9060	15.4659
48	1.6122	1.8154	2.0435	2.5871	3.2715	4.1323	6.5705	10.4013	16.3939
49	1.6283	1.8380	2.0741	2.6388	3.3533	4.2562	6.8333	10.9213	17.3775
50	1.6446	1.8610	2.1052	2.6916	3.4371	4.3839	7.1067	11.4674	18.4202

# 33

## PRESENT VALUE OF AN AMOUNT: $(1 + r)^{-n}$

The present value  $P$  which will amount to  $A$  in  $n$  years at an interest rate of  $r$  (in decimals) compounded annually is  $P = A(1 + r)^{-n}$ .

$\frac{r}{n}$	1%	$1\frac{1}{4}\%$	$1\frac{1}{2}\%$	2%	$2\frac{1}{2}\%$	3%	4%	5%	6%
1	.99010	.98765	.98522	.98039	.97561	.97087	.96154	.95238	.94340
2	.98030	.97546	.97066	.96117	.95181	.94260	.92456	.90703	.89000
3	.97059	.96342	.95632	.94232	.92860	.91514	.88900	.86384	.83962
4	.96098	.95152	.94218	.92385	.90595	.88849	.85480	.82270	.79209
5	.95147	.93978	.92826	.90573	.88385	.86261	.82193	.78353	.74726
6	.94205	.92817	.91454	.88797	.86230	.83748	.79031	.74622	.70496
7	.93272	.91672	.90103	.87056	.84127	.81309	.75992	.71068	.66506
8	.92348	.90540	.88771	.85349	.82075	.78941	.73069	.67684	.62741
9	.91434	.89422	.87459	.83676	.80073	.76642	.70259	.64461	.59190
10	.90529	.88318	.86167	.82035	.78120	.74409	.67556	.61391	.55839
11	.89632	.87228	.84893	.80426	.76214	.72242	.64958	.58468	.52679
12	.88745	.86151	.83639	.78849	.74356	.70138	.62460	.55684	.49697
13	.87866	.85087	.82403	.77303	.72542	.68095	.60057	.53032	.46884
14	.86996	.84037	.81185	.75788	.70773	.66112	.57748	.50507	.44230
15	.86135	.82999	.79985	.74301	.69047	.64186	.55526	.48102	.41727
16	.85282	.81975	.78803	.72845	.67362	.62317	.53391	.45811	.39365
17	.84438	.80963	.77639	.71416	.65720	.60502	.51337	.43630	.37136
18	.83602	.79963	.76491	.70016	.64117	.58739	.49363	.41552	.35034
19	.82774	.78976	.75361	.68643	.62553	.57029	.47464	.39573	.33051
20	.81954	.78001	.74247	.67297	.61027	.55368	.45639	.37689	.31180
21	.81143	.77038	.73150	.65978	.59539	.53755	.43883	.35894	.29416
22	.80340	.76087	.72069	.64684	.58086	.52189	.42196	.34185	.27751
23	.79544	.75147	.71004	.63416	.56670	.50669	.40573	.32557	.26180
24	.78757	.74220	.69954	.62172	.55288	.49193	.39012	.31007	.24698
25	.77977	.73303	.68921	.60953	.53939	.47761	.37512	.29530	.23300
26	.77205	.72398	.67902	.59758	.52623	.46369	.36069	.28124	.21981
27	.76440	.71505	.66899	.58586	.51340	.45019	.34682	.26785	.20737
28	.75684	.70622	.65910	.57437	.50088	.43708	.33348	.25509	.19563
29	.74934	.69750	.64936	.56311	.48866	.42435	.32065	.24295	.18456
30	.74192	.68889	.63976	.55207	.47674	.41199	.30832	.23138	.17411
31	.73458	.68038	.63031	.54125	.46511	.39999	.29646	.22036	.16425
32	.72730	.67198	.62099	.53063	.45377	.38834	.28506	.20987	.15496
33	.72010	.66369	.61182	.52023	.44270	.37703	.27409	.19987	.14619
34	.71297	.65549	.60277	.51003	.43191	.36604	.26355	.19035	.13791
35	.70591	.64740	.59387	.50003	.42137	.35538	.25342	.18129	.13011
36	.69892	.63941	.58509	.49022	.41109	.34503	.24367	.17266	.12274
37	.69200	.63152	.57644	.48061	.40107	.33498	.23430	.16444	.11579
38	.68515	.62372	.56792	.47119	.39128	.32523	.22529	.15661	.10924
39	.67837	.61602	.55953	.46195	.38174	.31575	.21662	.14915	.10306
40	.67165	.60841	.55126	.45289	.37243	.30656	.20829	.14205	.09722
41	.66500	.60090	.54312	.44401	.36335	.29763	.20028	.13528	.09172
42	.65842	.59348	.53509	.43530	.35448	.28896	.19257	.12884	.08653
43	.65190	.58616	.52718	.42677	.34584	.28054	.18517	.12270	.08163
44	.64545	.57892	.51939	.41840	.33740	.27237	.17805	.11686	.07701
45	.63905	.57177	.51171	.41020	.32917	.26444	.17120	.11130	.07265
46	.63273	.56471	.50415	.40215	.32115	.25674	.16461	.10600	.06854
47	.62646	.55774	.49670	.39427	.31331	.24926	.15828	.10095	.06466
48	.62026	.55086	.48936	.38654	.30567	.24200	.15219	.09614	.06100
49	.61412	.54406	.48213	.37896	.29822	.23495	.14634	.09156	.05755
50	.60804	.53734	.47500	.37153	.29094	.22811	.14071	.08720	.05429

# 34

## AMOUNT OF AN ANNUITY: $\frac{(1+r)^n - 1}{r}$

If a principal  $P$  is deposited at the end of each year at interest rate  $r$  (in decimals) compounded annually, then at the end of  $n$  years the accumulated amount is  $P \left[ \frac{(1-r)^n - 1}{r} \right]$ . The process is often called an *annuity*.

$n \backslash r$	1%	1 1/4%	1 1/2%	2%	2 1/2%	3%	4%	5%	6%
1	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000
2	2.0100	2.0125	2.0150	2.0200	2.0250	2.0300	2.0400	2.0500	2.0600
3	3.0301	3.0377	3.0452	3.0604	3.0756	3.0909	3.1216	3.1525	3.1836
4	4.0604	4.0756	4.0909	4.1216	4.1525	4.1836	4.2465	4.3101	4.3746
5	5.1010	5.1266	5.1523	5.2040	5.2563	5.3091	5.4163	5.5256	5.6371
6	6.1520	6.1907	6.2296	6.3081	6.3877	6.4684	6.6330	6.8019	6.9753
7	7.2135	7.2680	7.3230	7.4343	7.5474	7.6625	7.8983	8.1420	8.3938
8	8.2857	8.3589	8.4328	8.5830	8.7361	8.8923	9.2142	9.5491	9.8975
9	9.3685	9.4634	9.5593	9.7546	9.9545	10.1591	10.5828	11.0266	11.4913
10	10.4622	10.5817	10.7027	10.9497	11.2034	11.4639	12.0061	12.5779	13.1808
11	11.5668	11.7139	11.8633	12.1687	12.4835	12.8078	13.4864	14.2068	14.9716
12	12.6825	12.8604	13.0412	13.4121	13.7956	14.1920	15.0258	15.9171	16.8699
13	13.8093	14.0211	14.2368	14.6803	15.1404	15.6178	16.6268	17.7130	18.8821
14	14.9474	15.1964	15.4504	15.9739	16.5190	17.0863	18.2919	19.5986	21.0151
15	16.0969	16.3863	16.6821	17.2934	17.9319	18.5989	20.0236	21.5786	23.2760
16	17.2579	17.5912	17.9324	18.6393	19.3802	20.1569	21.8245	23.6575	25.6725
17	18.4304	18.8111	19.2014	20.0121	20.8647	21.7616	23.6975	25.8404	28.2129
18	19.6147	20.0462	20.4894	21.4123	22.3863	23.4144	25.6454	28.1324	30.9057
19	20.8109	21.2968	21.7967	22.8406	23.9460	25.1169	27.6712	30.5390	33.7600
20	22.0190	22.5630	23.1237	24.2974	25.5447	26.8704	29.7781	33.0660	36.7856
21	23.2392	23.8450	24.4705	25.7833	27.1833	28.6765	31.9692	35.7193	39.9927
22	24.4716	25.1431	25.8376	27.2990	28.8629	30.5368	34.2480	38.5052	43.3923
23	25.7163	26.4574	27.2251	28.8450	30.5844	32.4529	36.6179	41.4305	46.9958
24	26.9735	27.7881	28.6335	30.4219	32.3490	34.4265	39.0826	44.5020	50.8156
25	28.2432	29.1354	30.0630	32.0303	34.1578	36.4593	41.6459	47.7271	54.8645
26	29.5256	30.4996	31.5140	33.6709	36.0117	38.5530	44.3117	51.1135	59.1564
27	30.8209	31.8809	32.9867	35.3443	37.9120	40.7096	47.0842	54.6691	63.7058
28	32.1291	33.2794	34.4815	37.0512	39.8598	42.9309	49.9676	58.4026	68.5281
29	33.4504	34.6954	35.9987	38.7922	41.8563	45.2189	52.9663	62.3227	73.6398
30	34.7849	36.1291	37.5387	40.5681	43.9027	47.5754	56.0849	66.4388	79.0582
31	36.1327	37.5807	39.1018	42.3794	46.0003	50.0027	59.3283	70.7608	84.8017
32	37.4941	39.0504	40.6883	44.2270	48.1503	52.5028	62.7015	75.2988	90.8898
33	38.8690	40.5386	42.2986	46.1116	50.3540	55.0778	66.2095	80.0638	97.3432
34	40.2577	42.0453	43.9331	48.0338	52.6129	57.7302	69.8579	85.0670	104.1838
35	41.6603	43.5709	45.5921	49.9945	54.9282	60.4621	73.6522	90.3203	111.4348
36	43.0769	45.1155	47.2760	51.9944	57.3014	63.2759	77.5983	95.8363	119.1209
37	44.5076	46.6794	48.9851	54.0343	59.7339	66.1742	81.7022	101.6281	127.2681
38	45.9527	48.2629	50.7199	56.1149	62.2273	69.1594	85.9703	107.7095	135.9042
39	47.4123	49.8662	52.4807	58.2372	64.7830	72.2342	90.4091	114.0950	145.0585
40	48.8864	51.4896	54.2679	60.4020	67.4026	75.4013	95.0255	120.7998	154.7620
41	50.3752	53.1332	56.0819	62.6100	70.0876	78.6633	99.8265	127.8398	165.0477
42	51.8790	54.7973	57.9231	64.8622	72.8398	82.0232	104.8196	135.2318	175.9505
43	53.3978	56.4823	59.7920	67.1595	75.6608	85.4839	110.0124	142.9933	187.5076
44	54.9318	58.1883	61.6889	69.5027	78.5523	89.0484	115.4129	151.1430	199.7580
45	56.4811	59.9157	63.6142	71.8927	81.5161	92.7199	121.0294	159.7002	212.7435
46	58.0459	61.6646	65.5684	74.3306	84.5540	96.5015	126.8706	168.6852	226.5081
47	59.6263	63.4354	67.5519	76.8172	87.6679	100.3965	132.9454	178.1194	241.0986
48	61.2226	65.2284	69.5652	79.3535	90.8596	104.4084	139.2632	188.0254	256.5645
49	62.8348	67.0437	71.6087	81.9406	94.1311	108.5406	145.8337	198.4267	272.9584
50	64.4632	68.8818	73.6828	84.5794	97.4843	112.7969	152.6671	209.3480	290.3359

# 35

**PRESENT VALUE OF AN ANNUITY:**  $\frac{1 - (1+r)^{-n}}{r}$

An annuity in which the yearly payment at the end of each of  $n$  years is  $A$  at an interest rate  $r$  (in decimals) compounded annually has present value

$$A \left[ \frac{1 - (1+r)^{-n}}{r} \right].$$

$n \backslash r$	1%	1 1/4%	1 1/2%	2%	2 1/2%	3%	4%	5%	6%
1	0.9901	0.9877	0.9852	0.9804	0.9756	0.9709	0.9615	0.9524	0.9434
2	1.9704	1.9631	1.9559	1.9416	1.9274	1.9135	1.8861	1.8594	1.8334
3	2.9410	2.9265	2.9122	2.8839	2.8560	2.8286	2.7751	2.7232	2.6730
4	3.9020	3.8781	3.8544	3.8077	3.7620	3.7171	3.6299	3.5460	3.4651
5	4.8534	4.8178	4.7826	4.7135	4.6458	4.5797	4.4518	4.3295	4.2124
6	5.7955	5.7460	5.6972	5.6014	5.5081	5.4172	5.2421	5.0757	4.9173
7	6.7282	6.6627	6.5982	6.4720	6.3494	6.2303	6.0021	5.7864	5.5824
8	7.6517	7.5681	7.4859	7.3255	7.1701	7.0197	6.7327	6.4632	6.2098
9	8.5660	8.4623	8.3605	8.1622	7.9709	7.7861	7.4353	7.1078	6.8017
10	9.4713	9.3455	9.2222	8.9826	8.7521	8.5302	8.1109	7.7217	7.3601
11	10.3676	10.2178	10.0711	9.7868	9.5142	9.2526	8.7605	8.3064	7.8869
12	11.2551	11.0793	10.9075	10.5753	10.2578	9.9540	9.3851	8.8633	8.3838
13	12.1337	11.9302	11.7315	11.3484	10.9832	10.6350	9.9856	9.3936	8.8527
14	13.0037	12.7706	12.5434	12.1062	11.6909	11.2961	10.5631	9.8986	9.2950
15	13.8651	13.6005	13.3432	12.8493	12.3814	11.9379	11.1184	10.3797	9.7122
16	14.7179	14.4203	14.1313	13.5777	13.0550	12.5611	11.6523	10.8378	10.1059
17	15.5623	15.2299	14.9076	14.2919	13.7122	13.1661	12.1657	11.2741	10.4773
18	16.3983	16.0295	15.6726	14.9920	14.3534	13.7535	12.6593	11.6896	10.8276
19	17.2260	16.8193	16.4262	15.6785	14.9789	14.3238	13.1339	12.0853	11.1581
20	18.0456	17.5993	17.1686	16.3514	15.5892	14.8775	13.5903	12.4622	11.4699
21	18.8570	18.3697	17.9001	17.0112	16.1845	15.4150	14.0292	12.8212	11.7641
22	19.6604	19.1306	18.6208	17.6580	16.7654	15.9369	14.4511	13.1630	12.0416
23	20.4558	19.8820	19.3309	18.2922	17.321	16.4436	14.8568	13.4886	12.3034
24	21.2434	20.6242	20.0304	18.9139	17.8850	16.9355	15.2470	13.7986	12.5504
25	22.0232	21.3573	20.7196	19.5235	18.4244	17.4131	15.6221	14.0939	12.7834
26	22.7952	22.0813	21.3986	20.1210	18.9506	17.8768	15.9828	14.3752	13.0032
27	23.5596	22.7963	22.0676	20.7069	19.4640	18.3270	16.3296	14.6430	13.2105
28	24.3164	23.5025	22.7267	21.2813	19.9649	18.7641	16.6631	14.8981	13.4062
29	25.0658	24.2000	23.3761	21.8444	20.4535	19.1885	16.9837	15.1411	13.5907
30	25.8077	24.8889	24.0158	22.3965	20.9303	19.6004	17.2920	15.3725	13.7648
31	26.5423	25.5693	24.6461	22.9377	21.3954	20.0004	17.5885	15.5928	13.9291
32	27.2696	26.2413	25.2671	23.4683	21.8492	20.3888	17.8736	15.8027	14.0840
33	27.9897	26.9050	25.8790	23.9886	22.2919	20.7658	18.1476	16.0025	14.2302
34	28.7027	27.5605	26.4817	24.4986	22.7238	21.1318	18.4112	16.1929	14.3681
35	29.4086	28.2079	27.0756	24.9986	23.1452	21.4872	18.6646	16.3742	14.4982
36	30.1075	28.8473	27.6607	25.4888	23.5563	21.8323	18.9083	16.5469	14.6210
37	30.7995	29.4788	28.2371	25.9695	23.9573	22.1672	19.1426	16.7113	14.7368
38	31.4847	30.1025	28.8051	26.4406	24.3486	22.4925	19.3679	16.8679	14.8460
39	32.1630	30.7185	29.3646	26.9026	24.7303	22.8082	19.5845	17.0170	14.9491
40	32.8347	31.3269	29.9158	27.3555	25.1028	23.1148	19.7928	17.1591	15.0463
41	33.4997	31.9278	30.4590	27.7995	25.4661	23.4124	19.9931	17.2944	15.1380
42	34.1581	32.5213	30.9941	28.2348	25.8206	23.7014	20.1856	17.4232	15.2245
43	34.8100	33.1075	31.5212	28.6616	26.1664	23.9819	20.3708	17.5459	15.3062
44	35.4555	33.6864	32.0406	29.0800	26.5038	24.2543	20.5488	17.6628	15.3832
45	36.0945	34.2582	32.5523	29.4902	26.8330	24.5187	20.7200	17.7741	15.4558
46	36.7272	34.8229	33.0565	29.8923	27.1542	24.7754	20.8847	17.8801	15.5244
47	37.3597	35.3806	33.5532	30.2866	27.4675	25.0247	21.0429	17.9810	15.5890
48	37.9740	35.9815	34.0426	30.6731	27.7732	25.2667	21.1951	18.0772	15.6500
49	38.5881	36.4755	34.5247	31.0521	28.0714	25.5017	21.3415	18.1687	15.7076
50	39.1961	37.0129	34.9997	31.4236	28.3623	25.7298	21.4822	18.2559	15.7619


## Section VII: Probability and Statistics

# 36

### AREAS UNDER THE STANDARD NORMAL CURVE

from  $-\infty$  to  $x$

$$\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-t^2/2} dt$$




NOTE:  $\text{erf}(x) = 2\Phi(x\sqrt{2}) - 1$

$x$	0	1	2	3	4	5	6	7	8	9
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5754
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7258	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7518	.7549
0.7	.7580	.7612	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7996	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998
3.5	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998	.9998
3.6	.9998	.9998	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999
3.7	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999
3.8	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999	.9999
3.9	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000	1.0000

# 37

## ORDINATES OF THE STANDARD NORMAL CURVE

$$y = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$$




$x$	0	1	2	3	4	5	6	7	8	9
0.0	.3989	.3989	.3989	.3988	.3986	.3984	.3982	.3980	.3977	.3973
0.1	.3970	.3965	.3961	.3956	.3951	.3945	.3939	.3932	.3925	.3918
0.2	.3910	.3902	.3894	.3885	.3876	.3867	.3857	.3847	.3836	.3825
0.3	.3814	.3802	.3790	.3778	.3765	.3752	.3739	.3725	.3712	.3697
0.4	.3683	.3668	.3653	.3637	.3621	.3605	.3589	.3572	.3555	.3538
0.5	.3521	.3503	.3485	.3467	.3448	.3429	.3410	.3391	.3372	.3352
0.6	.3332	.3312	.3292	.3271	.3251	.3230	.3209	.3187	.3166	.3144
0.7	.3123	.3101	.3079	.3056	.3034	.3011	.2989	.2966	.2943	.2920
0.8	.2897	.2874	.2850	.2827	.2803	.2780	.2756	.2732	.2709	.2685
0.9	.2661	.2637	.2613	.2589	.2565	.2541	.2516	.2492	.2468	.2444
1.0	.2420	.2396	.2371	.2347	.2323	.2299	.2275	.2251	.2227	.2203
1.1	.2179	.2155	.2131	.2107	.2083	.2059	.2036	.2012	.1989	.1965
1.2	.1942	.1919	.1895	.1872	.1849	.1826	.1804	.1781	.1758	.1736
1.3	.1714	.1691	.1669	.1647	.1626	.1604	.1582	.1561	.1539	.1518
1.4	.1497	.1476	.1456	.1435	.1415	.1394	.1374	.1354	.1334	.1315
1.5	.1295	.1276	.1257	.1238	.1219	.1200	.1182	.1163	.1145	.1127
1.6	.1109	.1092	.1074	.1057	.1040	.1023	.1006	.0989	.0973	.0957
1.7	.0940	.0925	.0909	.0893	.0878	.0863	.0848	.0833	.0818	.0804
1.8	.0790	.0775	.0761	.0748	.0734	.0721	.0707	.0694	.0681	.0669
1.9	.0656	.0644	.0632	.0620	.0608	.0596	.0584	.0573	.0562	.0551
2.0	.0540	.0529	.0519	.0508	.0498	.0488	.0478	.0468	.0459	.0449
2.1	.0440	.0431	.0422	.0413	.0404	.0396	.0387	.0379	.0371	.0363
2.2	.0355	.0347	.0339	.0332	.0325	.0317	.0310	.0303	.0297	.0290
2.3	.0283	.0277	.0270	.0264	.0258	.0252	.0246	.0241	.0235	.0229
2.4	.0224	.0219	.0213	.0208	.0203	.0198	.0194	.0189	.0184	.0180
2.5	.0175	.0171	.0167	.0163	.0158	.0154	.0151	.0147	.0143	.0139
2.6	.0136	.0132	.0129	.0126	.0122	.0119	.0116	.0113	.0110	.0107
2.7	.0104	.0101	.0099	.0096	.0093	.0091	.0088	.0086	.0084	.0081
2.8	.0079	.0077	.0075	.0073	.0071	.0069	.0067	.0065	.0063	.0061
2.9	.0060	.0058	.0056	.0055	.0053	.0051	.0050	.0048	.0047	.0046
3.0	.0044	.0043	.0042	.0040	.0039	.0038	.0037	.0036	.0035	.0034
3.1	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026	.0025	.0025
3.2	.0024	.0023	.0022	.0022	.0021	.0020	.0020	.0019	.0018	.0018
3.3	.0017	.0017	.0016	.0016	.0015	.0015	.0014	.0014	.0013	.0013
3.4	.0012	.0012	.0012	.0011	.0011	.0010	.0010	.0010	.0009	.0009
3.5	.0009	.0008	.0008	.0008	.0008	.0007	.0007	.0007	.0007	.0006
3.6	.0006	.0006	.0006	.0005	.0005	.0005	.0005	.0005	.0005	.0004
3.7	.0004	.0004	.0004	.0004	.0004	.0004	.0003	.0003	.0003	.0003
3.8	.0003	.0003	.0003	.0003	.0003	.0002	.0002	.0002	.0002	.0002
3.9	.0002	.0002	.0002	.0002	.0002	.0002	.0002	.0002	.0001	.0001

# 38

**PERCENTILE VALUES ( $t_p$ )  
FOR STUDENT'S t  
DISTRIBUTION**

with  $n$  degrees of freedom (shaded area =  $p$ )




$n$	$t_{.995}$	$t_{.99}$	$t_{.975}$	$t_{.95}$	$t_{.90}$	$t_{.80}$	$t_{.75}$	$t_{.70}$	$t_{.60}$	$t_{.55}$
1	63.66	31.82	12.71	6.31	3.08	1.376	1.000	.727	.325	.158
2	9.92	6.96	4.30	2.92	1.89	1.061	.816	.617	.289	.142
3	5.84	4.54	3.18	2.35	1.64	.978	.765	.584	.277	.137
4	4.60	3.75	2.78	2.13	1.53	.941	.741	.569	.271	.134
5	4.03	3.36	2.57	2.02	1.48	.920	.727	.559	.267	.132
6	3.71	3.14	2.45	1.94	1.44	.906	.718	.553	.265	.131
7	3.50	3.00	2.36	1.90	1.42	.896	.711	.549	.263	.130
8	3.36	2.90	2.31	1.86	1.40	.889	.706	.546	.262	.130
9	3.25	2.82	2.26	1.83	1.38	.883	.703	.543	.261	.129
10	3.17	2.76	2.23	1.81	1.37	.879	.700	.542	.260	.129
11	3.11	2.72	2.20	1.80	1.36	.876	.697	.540	.260	.129
12	3.06	2.68	2.18	1.78	1.36	.873	.695	.539	.259	.128
13	3.01	2.65	2.16	1.77	1.35	.870	.694	.538	.259	.128
14	2.98	2.62	2.14	1.76	1.34	.868	.692	.537	.258	.128
15	2.95	2.60	2.13	1.75	1.34	.866	.691	.536	.258	.128
16	2.92	2.58	2.12	1.75	1.34	.865	.690	.535	.258	.128
17	2.90	2.57	2.11	1.74	1.33	.863	.689	.534	.257	.128
18	2.88	2.55	2.10	1.73	1.33	.862	.688	.534	.257	.127
19	2.86	2.54	2.09	1.73	1.33	.861	.688	.533	.257	.127
20	2.84	2.53	2.09	1.72	1.32	.860	.687	.533	.257	.127
21	2.83	2.52	2.08	1.72	1.32	.859	.686	.532	.257	.127
22	2.82	2.51	2.07	1.72	1.32	.858	.686	.532	.256	.127
23	2.81	2.50	2.07	1.71	1.32	.858	.685	.532	.256	.127
24	2.80	2.49	2.06	1.71	1.32	.857	.685	.531	.256	.127
25	2.79	2.48	2.06	1.71	1.32	.856	.684	.531	.256	.127
26	2.78	2.48	2.06	1.71	1.32	.856	.684	.531	.256	.127
27	2.77	2.47	2.05	1.70	1.31	.855	.684	.531	.256	.127
28	2.76	2.47	2.05	1.70	1.31	.855	.683	.530	.256	.127
29	2.76	2.46	2.04	1.70	1.31	.854	.683	.530	.256	.127
30	2.75	2.46	2.04	1.70	1.31	.854	.683	.530	.256	.127
40	2.70	2.42	2.02	1.68	1.30	.851	.681	.529	.255	.126
60	2.66	2.39	2.00	1.67	1.30	.848	.679	.527	.254	.126
120	2.62	2.36	1.98	1.66	1.29	.845	.677	.526	.254	.126
$\infty$	2.58	2.33	1.96	1.645	1.28	.842	.674	.524	.253	.126

Source: R. A. Fisher and F. Yates, *Statistical Tables for Biological, Agricultural and Medical Research* (6th edition, 1963), Table III, Oliver and Boyd Ltd., Edinburgh, by permission of the authors and publishers.

# 39

**PERCENTILE VALUES ( $\chi^2_p$ )  
FOR  $\chi^2$  (CHI-SQUARE)  
DISTRIBUTION**

with  $n$  degrees of freedom (shaded area =  $p$ )




$n$	$\chi^2_{.995}$	$\chi^2_{.99}$	$\chi^2_{.975}$	$\chi^2_{.95}$	$\chi^2_{.90}$	$\chi^2_{.75}$	$\chi^2_{.50}$	$\chi^2_{.25}$	$\chi^2_{.10}$	$\chi^2_{.05}$	$\chi^2_{.025}$	$\chi^2_{.01}$	$\chi^2_{.005}$
1	7.88	6.63	5.02	3.84	2.71	1.32	.455	.102	.0158	.0039	.0010	.0002	.0000
2	10.6	9.21	7.38	5.99	4.61	2.77	1.39	.575	.211	.103	.0506	.0201	.0100
3	12.8	11.3	9.35	7.81	6.25	4.11	2.37	1.21	.584	.352	.216	.115	.072
4	14.9	13.3	11.1	9.49	7.78	5.39	3.36	1.92	1.06	.711	.484	.297	.207
5	16.7	15.1	12.8	11.1	9.24	6.63	4.35	2.67	1.61	1.15	.831	.554	.412
6	18.5	16.8	14.4	12.6	10.6	7.84	5.35	3.45	2.20	1.64	1.24	.872	.676
7	20.3	18.5	16.0	14.1	12.0	9.04	6.35	4.25	2.83	2.17	1.69	1.24	.989
8	22.0	20.1	17.5	15.5	13.4	10.2	7.34	5.07	3.49	2.73	2.18	1.65	1.34
9	23.6	21.7	19.0	16.9	14.7	11.4	8.34	5.90	4.17	3.33	2.70	2.09	1.73
10	25.2	23.2	20.5	18.3	16.0	12.5	9.34	6.74	4.87	3.94	3.25	2.56	2.16
11	26.8	24.7	21.9	19.7	17.3	13.7	10.3	7.58	5.58	4.57	3.82	3.05	2.60
12	28.3	26.2	23.3	21.0	18.5	14.8	11.3	8.44	6.30	5.23	4.40	3.57	3.07
13	29.8	27.7	24.7	22.4	19.8	16.0	12.3	9.30	7.04	5.89	5.01	4.11	3.57
14	31.3	29.1	26.1	23.7	21.1	17.1	13.3	10.2	7.79	6.57	5.63	4.66	4.07
15	32.8	30.6	27.5	25.0	22.3	18.2	14.3	11.0	8.55	7.26	6.26	5.23	4.60
16	34.3	32.0	28.8	26.3	23.5	19.4	15.3	11.9	9.31	7.96	6.91	5.81	5.14
17	35.7	33.4	30.2	27.6	24.8	20.5	16.3	12.8	10.1	8.67	7.56	6.41	5.70
18	37.2	34.8	31.5	28.9	26.0	21.6	17.3	13.7	10.9	9.39	8.23	7.01	6.26
19	38.6	36.2	32.9	30.1	27.2	22.7	18.3	14.6	11.7	10.1	8.91	7.63	6.84
20	40.0	37.6	34.2	31.4	28.4	23.8	19.3	15.5	12.4	10.9	9.59	8.26	7.43
21	41.4	38.9	35.5	32.7	29.6	24.9	20.3	16.3	13.2	11.6	10.3	8.90	8.03
22	42.8	40.3	36.8	33.9	30.8	26.0	21.3	17.2	14.0	12.3	11.0	9.54	8.64
23	44.2	41.6	38.1	35.2	32.0	27.1	22.3	18.1	14.8	13.1	11.7	10.2	9.26
24	45.6	43.0	39.4	36.4	33.2	28.2	23.3	19.0	15.7	13.8	12.4	10.9	9.89
25	46.9	44.3	40.6	37.7	34.4	29.3	24.3	19.9	16.5	14.6	13.1	11.5	10.5
26	48.3	45.6	41.9	38.9	35.6	30.4	25.3	20.8	17.3	15.4	13.8	12.2	11.2
27	49.6	47.0	43.2	40.1	36.7	31.5	26.3	21.7	18.1	16.2	14.6	12.9	11.8
28	51.0	48.3	44.5	41.3	37.9	32.6	27.3	22.7	18.9	16.9	15.3	13.6	12.5
29	52.3	49.6	45.7	42.6	39.1	33.7	28.3	23.6	19.8	17.7	16.0	14.3	13.1
30	53.7	50.9	47.0	43.8	40.3	34.8	29.3	24.5	20.6	18.5	16.8	15.0	13.8
40	66.8	63.7	59.3	55.8	51.8	45.6	39.3	33.7	29.1	26.5	24.4	22.2	20.7
50	79.5	76.2	71.4	67.5	63.2	56.3	49.3	42.9	37.7	34.8	32.4	29.7	28.0
60	92.0	88.4	83.3	79.1	74.4	67.0	59.3	52.3	46.5	43.2	40.5	37.5	35.5
70	104.2	100.4	95.0	90.5	85.5	77.6	69.3	61.7	55.3	51.7	48.8	45.4	43.3
80	116.3	112.3	106.6	101.9	96.6	88.1	79.3	71.1	64.3	60.4	57.2	53.5	51.2
90	128.3	124.1	118.1	113.1	107.6	98.6	89.3	80.6	73.3	69.1	65.6	61.8	59.2
100	140.2	135.8	129.6	124.3	118.5	109.1	99.3	90.1	82.4	77.9	74.2	70.1	67.3

Source: Catherine M. Thompson, *Table of percentage points of the  $\chi^2$  distribution*, Biometrika, Vol. 32 (1941), by permission of the author and publisher.

# 40

## 95th PERCENTILE VALUES FOR THE F DISTRIBUTION

$n_1$  = degrees of freedom for numerator  
 $n_2$  = degrees of freedom for denominator  
 (shaded area = .95)



$\frac{n_1}{n_2}$	1	2	3	4	5	6	8	12	16	20	30	40	50	100	$\infty$
1	161.4	199.5	215.7	224.6	230.2	234.0	238.9	243.9	246.3	248.0	250.1	251.1	252.2	253.0	254.3
2	18.51	19.00	19.16	19.25	19.30	19.33	19.37	19.41	19.43	19.45	19.46	19.46	19.47	19.49	19.50
3	10.13	9.55	9.28	9.12	9.01	8.94	8.85	8.74	8.69	8.66	8.62	8.60	8.58	8.56	8.53
4	7.71	6.94	6.59	6.39	6.26	6.16	6.04	5.91	5.84	5.80	5.75	5.71	5.70	5.66	5.63
5	6.61	5.79	5.41	5.19	5.05	4.95	4.82	4.68	4.60	4.56	4.50	4.46	4.44	4.40	4.36
6	5.99	5.14	4.76	4.53	4.39	4.28	4.15	4.00	3.92	3.87	3.81	3.77	3.75	3.71	3.67
7	5.59	4.74	4.35	4.12	3.97	3.87	3.73	3.57	3.49	3.44	3.38	3.34	3.32	3.28	3.23
8	5.32	4.46	4.07	3.84	3.69	3.58	3.44	3.28	3.20	3.15	3.08	3.05	3.03	2.98	2.93
9	5.12	4.26	3.86	3.63	3.48	3.37	3.23	3.07	2.98	2.93	2.86	2.82	2.80	2.76	2.71
10	4.96	4.10	3.71	3.48	3.33	3.22	3.07	2.91	2.82	2.77	2.70	2.67	2.64	2.59	2.54
11	4.84	3.98	3.59	3.36	3.20	3.09	2.95	2.79	2.70	2.65	2.57	2.53	2.50	2.45	2.40
12	4.75	3.89	3.49	3.26	3.11	3.00	2.85	2.69	2.60	2.54	2.46	2.42	2.40	2.35	2.30
13	4.67	3.81	3.41	3.18	3.03	2.92	2.77	2.60	2.51	2.46	2.38	2.34	2.32	2.26	2.21
14	4.60	3.74	3.34	3.11	2.96	2.85	2.70	2.53	2.44	2.39	2.31	2.27	2.24	2.19	2.13
15	4.54	3.68	3.29	3.06	2.90	2.79	2.64	2.48	2.39	2.33	2.25	2.21	2.18	2.12	2.07
16	4.49	3.63	3.24	3.01	2.85	2.74	2.59	2.42	2.33	2.28	2.20	2.16	2.13	2.07	2.01
17	4.45	3.59	3.20	2.96	2.81	2.70	2.55	2.38	2.29	2.23	2.15	2.11	2.08	2.02	1.96
18	4.41	3.55	3.16	2.93	2.77	2.66	2.51	2.34	2.25	2.19	2.11	2.07	2.04	1.98	1.92
19	4.38	3.52	3.13	2.90	2.74	2.63	2.48	2.31	2.21	2.15	2.07	2.02	2.00	1.94	1.88
20	4.35	3.49	3.10	2.87	2.71	2.60	2.45	2.28	2.18	2.12	2.04	1.99	1.96	1.90	1.84
22	4.30	3.44	3.05	2.82	2.66	2.55	2.40	2.23	2.13	2.07	1.98	1.93	1.91	1.84	1.78
24	4.26	3.40	3.01	2.78	2.62	2.51	2.36	2.18	2.09	2.03	1.94	1.89	1.86	1.80	1.73
26	4.23	3.37	2.98	2.74	2.59	2.47	2.32	2.15	2.05	1.99	1.90	1.85	1.82	1.76	1.69
28	4.20	3.34	2.95	2.71	2.56	2.45	2.29	2.12	2.02	1.96	1.87	1.81	1.78	1.72	1.65
30	4.17	3.32	2.92	2.69	2.53	2.42	2.27	2.09	1.99	1.93	1.84	1.79	1.76	1.69	1.62
40	4.08	3.23	2.84	2.61	2.45	2.34	2.18	2.00	1.90	1.84	1.74	1.69	1.66	1.59	1.51
50	4.03	3.18	2.79	2.56	2.40	2.29	2.13	1.95	1.85	1.78	1.69	1.63	1.60	1.52	1.44
60	4.00	3.15	2.76	2.53	2.37	2.25	2.10	1.92	1.81	1.75	1.65	1.59	1.56	1.48	1.39
70	3.98	3.13	2.74	2.50	2.35	2.23	2.07	1.89	1.79	1.72	1.62	1.56	1.53	1.45	1.35
80	3.96	3.11	2.72	2.48	2.33	2.21	2.05	1.88	1.77	1.70	1.60	1.54	1.51	1.42	1.32
100	3.94	3.09	2.70	2.46	2.30	2.19	2.03	1.85	1.75	1.68	1.57	1.51	1.48	1.39	1.28
150	3.91	3.06	2.67	2.43	2.27	2.16	2.00	1.82	1.71	1.64	1.54	1.47	1.44	1.34	1.22
200	3.89	3.04	2.65	2.41	2.26	2.14	1.98	1.80	1.69	1.62	1.52	1.45	1.42	1.32	1.19
400	3.86	3.02	2.62	2.39	2.23	2.12	1.96	1.78	1.67	1.60	1.49	1.42	1.38	1.28	1.13
$\infty$	3.84	2.99	2.60	2.37	2.21	2.09	1.94	1.75	1.64	1.57	1.46	1.40	1.32	1.24	1.00

Source: G. W. Snedecor and W. G. Cochran, *Statistical Methods* (6th edition, 1967), Iowa State University Press, Ames, Iowa, by permission of the authors and publisher.


# 41

## 99th PERCENTILE VALUES

### FOR THE F DISTRIBUTION

$n_1$  = degrees of freedom for numerator

$n_2$  = degrees of freedom for denominator  
(shaded area = .99)



$\frac{n_1}{n_2}$	1	2	3	4	5	6	8	12	16	20	30	40	50	100	$\infty$
1	4052	4999	5403	5625	5764	5859	5981	6106	6169	6208	6258	6286	6302	6334	6366
2	98.49	99.01	99.17	99.25	99.30	99.33	99.36	99.42	99.44	99.45	99.47	99.48	99.48	99.49	99.50
3	34.12	30.81	29.46	28.71	28.24	27.41	27.49	27.05	28.63	26.69	26.50	26.41	26.35	26.23	26.12
4	21.20	18.00	16.69	15.98	15.52	15.21	14.80	14.37	14.15	14.02	13.83	13.74	13.69	13.57	13.46
5	16.26	13.27	12.06	11.39	10.97	10.67	10.27	9.89	9.68	9.55	9.38	9.29	9.24	9.13	9.02
6	13.74	10.92	9.78	9.15	8.75	8.47	8.10	7.72	7.52	7.39	7.23	7.14	7.09	6.99	6.88
7	12.25	9.55	8.45	7.85	7.46	7.19	6.84	6.47	6.27	6.15	5.98	5.90	5.85	5.75	5.65
8	11.26	8.65	7.59	7.01	6.63	6.37	6.03	5.67	5.48	5.36	5.20	5.11	5.06	4.96	4.86
9	10.56	8.02	6.99	6.42	6.06	5.80	5.47	5.11	4.92	4.80	4.64	4.56	4.51	4.41	4.31
10	10.04	7.56	6.55	5.99	5.64	5.39	5.06	4.71	4.52	4.41	4.25	4.17	4.12	4.01	3.91
11	9.05	7.20	6.22	5.67	5.32	5.07	4.74	4.40	4.21	4.10	3.94	3.86	3.80	3.70	3.60
12	9.33	6.93	5.95	5.41	5.06	4.82	4.50	4.16	3.98	3.86	3.70	3.61	3.56	3.46	3.36
13	9.07	6.70	5.74	5.20	4.86	4.62	4.30	3.96	3.78	3.67	3.51	3.42	3.37	3.27	3.16
14	8.86	6.51	5.56	5.03	4.69	4.46	4.14	3.80	3.62	3.51	3.34	3.26	3.21	3.11	3.00
15	8.68	6.36	5.42	4.89	4.56	4.32	4.00	3.67	3.48	3.36	3.20	3.12	3.07	2.97	2.87
16	8.53	6.23	5.29	4.77	4.44	4.20	3.89	3.55	3.37	3.25	3.10	3.01	2.96	2.86	2.75
17	8.40	6.11	5.18	4.67	4.34	4.10	3.79	3.45	3.27	3.16	3.00	2.92	2.86	2.76	2.65
18	8.28	6.01	5.09	4.58	4.25	4.01	3.71	3.37	3.19	3.07	2.91	2.83	2.78	2.68	2.57
19	8.18	5.93	5.01	4.50	4.17	3.94	3.63	3.30	3.12	3.00	2.84	2.76	2.70	2.60	2.49
20	8.10	5.85	4.94	4.43	4.10	3.87	3.56	3.23	3.05	2.94	2.77	2.69	2.63	2.53	2.42
22	7.94	5.72	4.82	4.31	3.99	3.76	3.45	3.12	2.94	2.83	2.67	2.58	2.53	2.42	2.31
24	7.82	5.61	4.72	4.22	3.90	3.67	3.36	3.03	2.85	2.74	2.58	2.49	2.44	2.33	2.21
26	7.72	5.53	4.64	4.14	3.82	3.59	3.29	2.96	2.77	2.66	2.50	2.41	2.36	2.25	2.13
28	7.64	5.45	4.57	4.07	3.76	3.53	3.23	2.90	2.71	2.60	2.44	2.35	2.30	2.18	2.06
30	7.56	5.39	4.51	4.02	3.70	3.47	3.17	2.84	2.66	2.55	2.38	2.29	2.24	2.13	2.01
40	7.31	5.18	4.31	3.83	3.51	3.29	2.99	2.66	2.49	2.37	2.20	2.11	2.05	1.94	1.81
50	7.17	5.06	4.20	3.72	3.41	3.18	2.88	2.56	2.39	2.26	2.10	2.00	1.94	1.82	1.68
60	7.08	4.98	4.13	3.65	3.34	3.12	2.82	2.50	2.32	2.20	2.03	1.93	1.87	1.74	1.60
70	7.01	4.92	4.08	3.60	3.29	3.07	2.77	2.45	2.28	2.15	1.98	1.88	1.82	1.69	1.53
80	6.96	4.88	4.04	3.56	3.25	3.04	2.74	2.41	2.24	2.11	1.94	1.84	1.78	1.65	1.49
100	6.90	4.82	3.98	3.51	3.20	2.99	2.69	2.36	2.19	2.06	1.89	1.79	1.73	1.59	1.43
150	6.81	4.75	3.91	3.44	3.14	2.92	2.62	2.30	2.12	2.00	1.83	1.72	1.66	1.51	1.33
200	6.76	4.71	3.88	3.41	3.11	2.90	2.60	2.28	2.09	1.97	1.79	1.69	1.62	1.48	1.28
400	6.70	4.66	3.83	3.36	3.06	2.85	2.55	2.23	2.04	1.92	1.74	1.64	1.57	1.42	1.19
$\infty$	6.64	4.60	3.78	3.32	3.02	2.80	2.51	2.18	1.99	1.87	1.69	1.59	1.52	1.36	1.00

Source: G. W. Snedecor and W. G. Cochran, *Statistical Methods* (6th edition, 1967), Iowa State University Press, Ames, Iowa, by permission of the authors and publisher.

# 42

## RANDOM NUMBERS

51772	74640	42331	29044	46621	62898	93582	04186	19640	87056
24033	23491	83587	06568	21960	21387	76105	10863	97453	90581
45939	60173	52078	25424	11645	55870	56974	37428	93507	94271
30586	02133	75797	45406	31041	86707	12973	17169	88116	42187
03585	79353	81938	82322	96799	85659	36081	50884	14070	74950
64937	03355	95863	20790	65304	55189	00745	65253	11822	15804
15630	64759	51135	98527	62586	41889	25439	88036	24034	67283
09448	56301	57683	30277	94623	85418	68829	06652	41982	49159
21631	91157	77331	60710	52290	16835	48653	71590	16159	14676
91097	17480	29414	06829	87843	28195	27279	47152	35683	47280
50532	25496	95652	42457	73547	76552	50020	24819	52984	76168
07136	40876	79971	54195	25708	51817	36732	72484	94923	75936
27989	64728	10744	08396	56242	90985	28868	99431	50995	20507
85184	73949	36601	46253	00477	25234	09908	36574	72139	70185
54398	21154	97810	36764	32869	11785	55261	59009	38714	38723
65544	34371	09591	07839	58892	92843	72828	91341	84821	63886
08263	65952	85762	64236	39238	18776	84303	99247	46149	03229
39817	67906	48236	16057	81812	15815	63700	85915	19219	45943
62257	04077	79443	95203	02479	30763	92486	54083	23631	05825
53298	90276	62545	21944	16530	03878	07516	95715	02526	33537

# Index of Special Symbols and Notations

The following list show special symbols and notations together with pages on which they are defined or first appear. Cases where a symbol has more than one meaning will be clear from the context.

## Symbols

---

$\text{Ber}_n(x)$ , $\text{Bei}_n(x)$	Ber and Bei functions, 157
$B(m, n)$	beta function, 152
$B_b$	Bernoulli numbers, 142
$C(x)$	Fresnel cosine integral, 204
$C_i(x)$	cosine integral, 204
$e_1, e_2, e_3$	unit vectors in curvilinear coordinates, 127
$\text{erf}(x)$	error function, 203
$\text{erfc}(x)$	complementary error function, 203
$E = E(k, \pi/2)$	complete elliptic integral of the second kind, 198
$E = E(k, \phi)$	incomplete elliptic integral of the second kind, 198
$\text{Ei}(x)$	exponential integral, 203
$E_n$	Euler number, 142
$E(X)$	mean or expectation of random variable $X$ , 223
$f[x_0, x_1, \dots, x_k]$	divided distance formula, 287, 288
$F(a), F(x)$	cumulative distribution function, 209
$F(a, b; c; x)$	hypergeometric function, 178
$F(k, \phi)$	incomplete elliptic integral of the first kind, 198
$\mathcal{G}, \mathcal{G}^{-1}$	Fourier transform and inverse Fourier transform, 194
G. M.	geometric mean, 209
$h_1, h_2, h_3$	scale factors in curvilinear coordinates, 127
$H_n(x)$	Hermite polynomial, 169
$H_n^{(1)}(x), H_n^{(2)}(x)$	Hankel functions of the first and second kind, 155
H. M.	harmonic mean, 210
i, j, k	unit vectors in rectangular coordinates, 120
$I_n(x)$	modified Bessel function of the first kind, 155
$J_n(x)$	Bessel function of the first kind, 153
$K = F(k, \pi/2)$	complete elliptic integral of the first kind, 198
$\text{Ker}_n(x), \text{Kei}_n(x)$	Ker and Kei functions, 158
$K_n(x)$	modified Bessel function of the second kind, 156
$\ln x$ or $\log_e x$	natural logarithm of $x$ , 53
$\log x$ or $\log_{10} x$	common logarithm of $x$ , 53
$L_n(x)$	Laguerre polynomials, 171
$L_n^m(x)$	associated Laguerre polynomials, 173
$\mathcal{L}, \mathcal{L}^{-1}$	Laplace transform and inverse Laplace transform, 180
M.D.	mean deviation
$P(A/E)$	conditional probability of $A$ given $E$ , 219
$P_n(x)$	Legendre polynomials, 164
$P_n^m(x)$	associated Legendre polynomials, 173
$Q_u, M, Q_L$	quartiles, 211
$Q_n(x)$	Legendre functions of second kind, 167
$Q_n^m(x)$	associated Legendre functions of second kind, 168
$r$	sample correlation coefficient, 213
R.M.S.	root-mean-square, 211
$s$	sample standard deviation, 208
$s^2$	sample variance, 210
$s_{xy}$	sample covariance, 213
$\text{Si}(x)$	Sine integral, 203
$S(x)$	Fresnel sine integral, 204
$T_n(x)$	Chebyshev polynomials of first kind, 175

$U_n(x)$	Chebyshev polynomials of second kind, 176
$\text{Var}(X)$	variance of random variable $X$ , 224
$\bar{x}, \bar{\bar{x}}$	sample mean, grand mean, 208, 209
$x_k^{(n)}$	$k$ th zero of Legendre polynomial $P_n(x)$ , 232
$Y_n(x)$	Bessel function of second kind, 153
$Z$	standardized random variable, 226

## Greek Symbols

---

$\alpha_r$	$r$ th moment in standard units, 212	$\pi$	pi, 3
$\gamma$	Euler's constant, 4	$\phi$	spherical coordinate, 38
$\Gamma(x)$	gamma function, 149	$\Phi(p)$	$\sum 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{p}, \Phi(0) = 0, 154$
$\zeta(x)$	Riemann zeta function, 204	$\Phi(x)$	probability distribution function, 226
$\mu$	population mean, 208	$\sigma$	population standard deviation, 223
$\theta$	coordinate: cylindrical 37, polar, 11, 24; spherical, 38	$\sigma^2$	population variance, 223

## Notations

---

$A \sim B$	$A$ is asymptotic to $B$ or $A/B$ approaches 1, 151
$ A $	absolute value of $A = \begin{cases} A & \text{if } A \geq 0 \\ -A & \text{if } A < 0 \end{cases}$
$n!$	factorial $n$ , 7
$\binom{n}{k}$	binomial coefficients, 8
$y' = \frac{dy}{dx} = f'(x)$ $y'' = \frac{d^2y}{dx^2} = f''(x), \text{etc.}$	derivatives of $y$ or $f(x)$ with respect to $x$ , 62
$D^p = \frac{d^p}{dx^p}$	$p$ th derivative with respect to $x$ , 64
$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial x}, \frac{\partial^2 f}{\partial x \partial y}, \text{etc.}$	partial derivatives, 65
$\frac{\partial(x,y,z)}{\partial(u_1,u_2,u_3)}$	Jacobian, 128
$\int f(x) dx$	indefinite integral, 67
$\int_a^b f(x) dx$	definite integral, 108
$\int_C \mathbf{A} \bullet d\mathbf{r}$	line integral of $\mathbf{A}$ along $C$ , 124
$\mathbf{A} \bullet \mathbf{B}$	dot product of $\mathbf{A}$ and $\mathbf{B}$ , 120
$\mathbf{A} \times \mathbf{B}$	cross product of $\mathbf{A}$ and $\mathbf{B}$ , 121
$\nabla$	del operator, 122
$\nabla^2 = \nabla \bullet \nabla$	Laplacian operator, 123
$\nabla^4 = \nabla^2(\nabla^2)$	biharmonic operator, 123

# Index

- Adams-Basforth methods, 236  
Adams-Moulton methods, 236  
Addition formula:  
    Bessel functions, 163  
    Hermite polynomials, 170  
Addition rule (probability) 208  
Addition of vectors, 119  
Algebra of sets, 217  
Algebraic equations, solutions of, 13  
Alphabet, Greek, 3  
Analytic geometry, plane, 22–33  
    solid, 34–40  
Annuity table, 294  
Anti-derivative, 67  
Anti-logarithms, 53  
Arbitrage, 257  
Arithmetic:  
    mean, 208  
    series, 134  
Arithmetic-geometric series, 134  
Associated Laguerre polynomials, 173  
    (See also Laguerre polynomials)  
Associated Legendre functions, 164  
    of the first kind, 168  
    of the second kind, 168  
    (See also Legendre functions)  
Asymptotic expansions or formulas:  
  
    Bernoulli numbers, 143  
    Bessel functions, 160  
    Backward difference formulas, 228  
        Her and Bei functions, 157  
    Bayes formula, 220  
    Bernoulli numbers, 142  
        asymptotic formula, 143  
        series, 143  
    Bernoulli's differential equation, 116  
    Bessel functions, 153–164  
        graphs, 159  
        integral representation, 161  
        modified, 155  
        recurrence formulas, 154, 157  
        series, orthogonal, 161  
        tables, 281–289  
    Bessel's differential equation, 118, 153  
        general solution, 154  
        modified differential equation, 155  
Best fit, line of, 214  
Beta function, 152  
Biharmonic operator, 123  
Binomial:  
    coefficients, 7, 228, 279  
    distribution, 226  
    formula, 7  
    series, 136  
Bipolar coordinates, 131  
Bisection method, 223  
Bivariate data, 212  
Black-Sholes formula, 259  
Brownian Motion, 254  
  
Carioid, 29  
Cassini, ovals of, 32  
Catalan's constant, 200  
Catenary, 29  
Cauchy or Euler differential equation, 117  
Cauchy's form of remainder in Taylor series, 134  
Cauchy-Schwarz inequality, 205  
    for integrals, 206  
Central tendency, 208  
Chain rule for derivatives, 67  
Chebyshev polynomials, 175  
    of the first kind, 175  
    of the second kind, 176  
    recurrence formula, 175  
Chebyshev's differential equation, 175  
    general solution, 177  
Chebyshev's inequality, 206  
Chi-Square distribution, 226  
    table of values, 299  
Circle, 17, 25  
Coefficient:  
    of excess (kurtosis), 212  
    of skewness, 212  
Coefficients:  
    binomial, 7  
    multinomial, 9  
Complementary error function, 203  
Complex:  
    conjugate, 10  
    numbers, 10  
    logarithm of, 55  
    plane, 10  
Components of a vector, 120  
Compound amount, 282  
Computable functions, 250  
Computing, 250  
Confocal:  
    ellipsoidal coordinates, 133  
    paraboloidal coordinates, 133  
Conical coordinates, 129  
Conics, 25 (See also Ellipse, Parabola, Hyperbola)  
Conjugate, complex, 10  
Constant of integration, 67  
Constants, 3  
    series of, 134  
Continuous random variable, 224  
Convergence, interval of, 138  
Conversion factors, 15  
Convolution theorem, Fourier transform, 194  
Coordinates, 127  
    bipolar, 131  
    confocal ellipsoidal, 133  
    confocal paraboloidal, 133  
    conical, 132  
    curvilinear, 127  
    cylindrical, 129  
    elliptic cylindrical, 130  
    oblate spheroidal, 131  
    paraboloidal, 130  
    prolate spheroidal, 131  
    spherical, 129  
    toroidal, 132  
Correlation coefficient, 213  
Cosine, 43  
    graph of, 46  
    table of values, 245  
Cosine integral, 203, 276  
Cosines, law of, 51  
Covariance, 213  
Cross or vector product, 121  
Cubic equation, solution of, 13

- Cumulative distribution function, 225  
 Curl, 123  
 Curve fitting, 215  
 Curvilinear coordinates, 134  
 Cycloid, 28  
 Cylindrical coordinates, 37, 129
- Definite integrals, 108–116  
 approximate formula, 109  
 definition of, 108  
 Degrees, conversion to radians, 271  
 Del operator, 122  
 Delta hedging, 260  
 DeMoivre’s theorem, 11  
 Derivatives, 62–66  
 chain rule for, 62  
 higher, 64  
 Leibniz’s rule, 64  
 of vectors, 122  
 Deviation:  
 mean, 210  
 standard, 210  
 Differential equations, numerical methods for solution:  
 ordinary, 235–236  
 partial, 237–240  
 Differentials, 65, 66  
 Differentiation, 62–66 (*See also* Derivatives)  
 Direction numbers, 34  
 cosines, 34  
 Direction set, 246  
 Discrete random variable, 223  
 Distributions, probability, 226  
 Divergence, 122, 128  
 theorem, 126  
 Divided-difference formula (general), 228  
 Dot or scalar product, 120  
 Double integrals, 125
- Eccentricity, 25  
 Ellipse, 18, 25  
 Ellipsoid, 39  
 Elliptic cylinder, 41  
 Elliptic cylindrical coordinates, 130  
 Elliptic functions, 198–202  
 Jacobi’s, 199  
 series expansion, 200  
 Elliptic integrals, 198–199  
 table of values, 290–291  
 Epicycloid, 30  
 Equality of vectors, 119  
 Equations, algebraic, 13  
 Error functions, 203  
 Euler:  
 constant, 4  
 differential equation, 117  
 methods, 235  
 numbers, 142  
 Euler-Maclaurin summation formula, 137  
 Exact differential equation, 116  
 Excess, coefficient of kurtosis, 212  
 Exponential curve (least-squares), 215  
 Exponential function, 53–54  
 series for, 139  
 table of values, 274–275  
 Exponential integral, 203, 276  
 Exponents, 53
- F* distribution, 226  
 table of values, 300–301  
 Factorial n, 7  
 table of values, 277  
 Factors, special, 5  
 Finance, 254
- Financial tables, 292–295  
 Finite-difference methods for solution of:  
 heat equation, 237  
 Poisson equation, 237  
 wave equation, 238  
 First-order divided-difference formula, 227  
 Five number summary [ $L, Q_L, M, Q_H, H$ ], 211  
 Fixed-point iteration, 234  
 Folium of Descartes, 31  
 Forward difference formulas, 228  
 Fourier series, 144–146  
 Fourier transform, 193  
 convolution of, 194  
 cosine, 194, 197  
 Parseval’s identity for, 193  
 sine, 194, 196  
 tables, 195–199  
 Fourier’s integral theorem, 193  
 Fresnel sine and cosine integral, 204  
 Frullani’s integral, 115
- Gamma function, 149, 150  
 relation to beta function, 152  
 table of values, 258  
 Gauss’ theorem, 126  
 Gauss-Legendre formula, 232  
 Gauss-Seidel method, 230  
 Gaussian quadrature formula, 231  
 Generating functions, 157, 165, 168, 169, 171, 173, 175, 176  
 Geometric:  
 mean (G.M.), 209  
 series, 134  
 Geometry, 16–21  
 analytic, 22–40  
 Gradient, 122, 128  
 Grand mean, 209  
 Greek alphabet, 3  
 Green’s theorem, 126  
 Griggsian logarithms, 53
- Half angle formulas, 48  
 Half rectified sine wave function, 191  
 Hankel functions, 155  
 Harmonic mean, 209  
 Heat equation, 237  
 Heaviside’s unit function, 192  
 Hermite:  
 interpolation, 229  
 polynomials, 169–170  
 Hermite’s differential equation, 169  
 Heun’s method, 235  
 Holder’s inequality, 205  
 for integrals, 206  
 Homogeneous differential equation, 116  
 linear second order, 117  
 Hyperbola, 25  
 Hyperbolic functions, 56–61  
 graphs of, 59  
 inverse, 59–61  
 series for, 140  
 Hyperboloid, 39  
 Hypgeometric:  
 differential equation, 178  
 distribution, 226  
 functions, 178  
 Hypocycloid, 28, 30
- Imaginary part of a complex number, 10  
 Indefinite integrals, 67–107  
 definition of, 67  
 tables of, 71–107  
 transformation of, 69  
 Independent events, 221

- Inequalities, 205  
 Infinite products, 207  
 Integral calculus, fundamental theorem, 108  
 Integrals:  
   definite (*see* Definite integrals)  
   improper, 108  
   indefinite (*see* Indefinite integrals)  
   line, 124  
   multiple, 125  
   surface, 125  
 Integration, 64  
   constant of, 67  
   general rules, 67–69  
   (*See also* Integrals)  
 Integration by parts, 67  
   generalized, 69  
 Intercepts, 22  
 Interest, 292–295  
 Interest rates, 256  
 Intermediate Value Theorem, 233  
 Interpolation, 227  
   Hermite, 229  
 Interpolatory formula (general), 228  
 Interquartile range, 211  
 Interval of convergence, 138  
 Inverse:  
   hyperbolic functions, 59–61  
   Laplace transforms, 180  
   trigonometric functions, 49–51  
 Iteration methods, 240  
   for general linear systems, 240  
   for Poisson equation, 240  
 Jacobi method, 240  
 Jacobi's elliptic functions, 199  
 Jacobian, 128  
 Ker and Kei functions, 158–159  
 Kurtosis, 212  
 Lagrange:  
   form of remainder, 138  
   interpolation, 227  
 Laguerre polynomials, 172  
   generating function for, 173  
   recurrence formula, 192  
 Laguerre's associated differential equation, 170  
 Laguerre's differential equation, 172  
 Landen's transformation, 199  
 Laplace transform, 180–192  
   complex inversion formula for, 180  
   definition of, 180  
   inverse, 180  
   tables of, 181–192  
 Laplacian, 123, 128  
 Least-squares:  
   curve, 215  
   line, 214  
 Legendre functions, 164–168  
   of the second kind, 166  
 Legendre polynomial, 164–165, 232  
   generating function for, 164  
   recurrence formula for, 166  
   tables of values for, 289  
 Legendre's associated differential equation, 168  
 Legendre's differential equation, 118, 164  
 Leibniz's rule, 64  
 Lemniscate, 28  
 Limaçon of Pascal, 32  
 Line, 22, 35  
   of best fit, 214  
   regression, 214  
 Line integral, 124  
 Logarithmic functions, 53–55  
   series for, 139  
   table of values, 245–246, 272–273  
   (*See also* Logarithms)  
 Logarithms, 53–55  
   of complex numbers, 55  
   Griggsian, 53  
 Maclaurin series, 138  
 Mathematical finance, 254  
 Mean, 208  
   continuous random variable, 224  
   deviation (M.D.), 211  
   discrete random variable, 223  
   geometric, 209  
   grand, 209  
   harmonic, 209  
   population, 212  
   weighted, 209  
 Mean value theorem,  
   for definite integrals, 108  
   generalized, 109  
 Median, 208  
 Midpoint rule, 231, 235  
 Midrange, 210  
 Milne's method, 236  
 Minkowski's inequality, 206  
   for integrals, 206  
 Mode, 209  
 Modified Bessel functions, 155–157  
   generating function for, 157  
   graphs of, 159  
   recurrence formulas for, 157  
 Modulus of a complex number, 11  
 Moment, rth, 212  
 Momental skewness, 212  
 Moments of inertia, 41  
 Monotonicity Rule (Probability), 218  
 Mutinomial coefficients, 9  
 Multiple integrals, 125  
 Napier's rules, 52  
 Natural logarithms and antilogarithms, 53  
   tables of, 272–273  
 Neumann's function, 153  
 Newton's:  
   backward-difference formula, 228  
   forward-difference formula, 228  
   interpolation, 227  
   method, 233  
 Nonhomogeneous differential equation, linear  
   second order, 117  
 Nonlinear equations, solution of, 233  
 Normal curve, 296–297  
   distribution, 226  
 Normal equations for least-squares line, 214  
 Normal random variable, 254  
 Null function, 189  
 Numbers:  
   Bernoulli, 142  
   Euler, 142  
 Numerical methods for partial differential equations, 237–239  
 Oblate spheroidal coordinates, 131  
 Orthogonal curvilinear coordinates, 127–128  
   formulas involving, 128  
 Orthogonality:  
   Chebyshev's polynomials, 176  
   Laguerre polynomials, 172  
   Legendre polynomials, 165  
 Ovals of Cassini, 32  
 Parabola, 25  
   segment of, 18

- Parabolic cylindrical coordinates, 129  
 Paraboloid, 40  
 Paraboloidal coordinates, 130  
 Parallelepiped, 19  
 Parallelogram, 7  
 Parameter, 208  
 Parseval's identity for:  
 Fourier series, 144  
 Fourier transform, 194  
 Partial:  
 derivatives, 65  
 differential equations, numerical methods, 237  
 Pascal's triangle, 8  
 Percentile,  $k$ th, 211  
 Periods of elliptic functions, 200  
 Pictures, 247  
 Plane analytic geometry, formulas from, 22–27  
 Plane, complex, 10  
 Poisson:  
 distribution, 226  
 equation, 237  
 summation formula, 137  
 Polar:  
 coordinates, 24  
 form of a complex number, 11  
 Polygon, regular, 17  
 Polynomial function (least-squares), 214  
 Polynomials:  
 Chebyshev's, 175  
 Laguerre, 171  
 Legendre, 164  
 Population, 208  
 mean 210  
 standard deviation, 212  
 variance, 212  
 Power function (least-squares), 214  
 Power series, 138–141  
 reversion of 141  
 Powers, sums of, 134  
 Present value, of an amount, 293  
 of an annuity, 294  
 Probability, 217  
 distribution, 223  
 function, 218  
 tables, 296  
 Products, infinite, 207  
 special, 5  
 Pulse function, 192  
 Pyramid, volume of, 20  
  
 Quadrants, 43  
 Quadratic convergence, 233  
 Quadratic equation, solution of, 103  
 Quadrature, 231–232  
 Quartic equation, solution of, 13  
 Quartile coefficient of skewness, 212  
 Quartiles  $[Q_L, M, Q_U]$ , 211  
 Quintuple, 248  
  
 Radians, 4, 44  
 table of conversion to degrees, 270  
 Random numbers table, 302  
 Random variable, 223–226  
 standardized, 226  
 Range, sample, 210  
 Real part of a complex number, 10  
 Reciprocals of powers, sums of, 135  
 Rectangle, 13  
 Rectangular coordinate system, 120  
 Rectangular coordinates, 24  
 transformation to polar coordinates, 24  
 Rectangular formula, 109  
 Rectified sine wave function, 191  
  
 Recurrence or recursion formulas:  
 Bessel functions, 154  
 Chebyshev's polynomials, 175  
 gamma function, 149  
 Hermite polynomials, 169  
 Laguerre polynomials, 171  
 Legendre polynomials, 165  
 Regression line, 214  
 Regular polygon, 17  
 Remainder:  
 Cauchy's form, 13  
 Lagrange form, 138  
 Remainder formula:  
 Gauss-Legendre interpolation, 232  
 Hermite interpolation, 230  
 Lagrange interpolation, 227  
 Reversion of power series, 141  
 Richardson method, 240  
 Riemann zeta function, 204  
 Right circular cone, 20  
 Rochigue's formula:  
 Laguerre polynomials, 171  
 Legendre's polynomials, 164  
 Root mean square (R.M.S.), 211  
 Roots of complex numbers, 11  
 Rose, 29  
 Rotation, 24, 37  
 Runge-Kutta method, 236  
  
 Sample, 208  
 covariance, 213  
 Saw tooth wave function, 191  
 Scalar, 119  
 multiplication of vectors, 119  
 Scalar or dot product, 120  
 Scale factors, 127  
 Scatterplot, 212  
 Schwarz (Cauchy-Schwarz) inequality, 205  
 for integrals, 206  
 Secant method, 233  
 Second-order differential equation, 117  
 Second-order divided-difference formula, 228  
 Sector of a circle, 17  
 Segment:  
 of circle, 18  
 of parabola, 18  
 Semi-interquartile range, 211  
 Separation of variables, 116  
 Series, arithmetic, 134  
 arithmetic-geometric, 134  
 binomial, 188  
 of constants, 134  
 Fourier, 144–148  
 geometric, 134  
 power, 138  
 of sums of powers, 134  
 Taylor, 138–141  
 Short selling, 257  
 Simpson's formula, 109, 231  
 Sine, 43  
 graph of, 46  
 table of values, 267  
 Sine integral, 88  
 table of values, 284  
 Sines, law of, 51  
 Skewness, 212  
 Solid analytic geometry, 34–40  
 Solutions of algebraic equations, 13–14  
 SOR (successive-overrelaxation) method, 240  
 Sphere, equations of, 38  
 surface area, 19  
 volume, 21

- Spherical coordinates, 38, 129  
Spherical triangle, 51  
Spiral of Archimedes, 33  
Square wave function, 191  
Squares error, 215  
Standard deviation, 210  
continuous random variable, 225  
discrete random variable, 224  
population, 212  
sample, 210  
Standardized random variable, 215  
Statistics, 208–216  
tables, 296–301  
Step function, 192  
Stirling's formula, 150  
Stochastic process, 219  
Stokes' theorem, 126  
Student's  $t$  distribution, 226  
table of, 308  
Successive-overrelaxation (SOR) method, 240  
Summation formula:  
Euler-Maclaurin, 137  
Poisson, 137  
Surface integrals, 125  
Tangent function, 43  
graph of, 46  
table of values, 269  
Tangents, law of, 51, 52  
Taylor series, 138–141  
two variables, 141  
Three-point interpolatory formula, 228  
Toroidal coordinates, 132  
Torus, surface area, volume, 18  
Total probability, Law of, 220  
Tractrix, 31  
Transformation:  
Jacobian of, 128  
of coordinates, 24, 36–37, 128  
of integrals, 70, 128  
Translation of coordinates:  
in a plane, 24  
in space, 36  
Trapezoid, area, perimeter, 16  
Trapezoidal rule (formula), 109, 231, 235  
Tree diagrams, Probability, 219  
Triangle inequality, 205  
Triangular wave function, 191  
Trigonometric functions, 43–52  
definition of, 43  
graphs of, 46  
inverse, 49–50  
series for, 139  
tables of, 267–269  
Triple integrals, 125  
Trochoid, 30  
Turing Machine, 246, 248  
Two-point formula, 228  
Two-point interpolatory formula, 228  
Unit function, Heaviside's, 192  
Unit normal to the surface, 125  
Unit vector, 120  
Variance, 210  
continuous random variable, 225  
discrete random variable, 224  
population, 210  
sample, 210  
Vector analysis, 119–133  
Vector or cross-product, 121  
Vectors, 119  
derivatives of, 122  
integrals involving, 124  
unit, 119  
Volume integrals, 125  
Wallis' product, 207  
Wave equation, 238  
Weber's function, 153  
Weighted mean, 209  
Witch of Agnesi, 31  
 $x$ -intercept, 22  
 $y$ -intercept, 22  
Zero vector, 119  
Zeros of Bessel functions, 287  
Zeta function of Riemann, 204